



**CONTROLS SNAILS, SLUGS, ANTS, EARWIGS, FALL ARMY-  
WORMS, AND SOD WEBWORMS (LAWN MOTHS) IN ORNA-  
MENTAL GARDENS AND DICHONDRA AND TURF LAWNS**

**GRANULAR FORMULATION**

In Convenient Shaker Applicator

For Faster, More Effective Kill

Blends into Lawn and Ornamental Areas

239-2314

**ACCEPTED**  
*Feb. 9, 1970*  
UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTER-  
ED UNDER NO. .... SUBJECT  
~~IN ALL COMMENTS~~

Active Ingredients	By Wt.
Carbaryl (1-naphthyl N-methylcarbamate)* .....	5%
Metalddehyde .....	3%
Inert Ingredients .....	92%
U. S. Pat. Nos. 2,903,478 - 3,009,855	

\*SEVIN—Trade Mark of Union Carbide Corporation for the active ingredient,  
1-naphthyl N-methylcarbamate

**WARNING: KEEP OUT OF REACH OF CHILDREN.**

SEE BACK PANEL FOR ADDITIONAL WARNINGS.

**NET WEIGHT 1 POUND**

NOTICE: Buyer assumes all responsibility for safety and use not in accord-  
ance with directions.



## ORTHO® BUG-GETA® Granules

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER.

ORTHO BUG-GETA Granules will control Snails, Slugs, Ants, Earwigs, Sowbugs, Fall Armyworms and Sod Webworms (Lawn Moths).

### DIRECTIONS FOR USE

WATER AREA TO BE TREATED WELL—PRIOR TO APPLICATION. This product can be applied anytime during the day but late afternoon or evening is preferred.

Simply shake on the proper amount of material from the convenient shaker top can. May also be applied from a fertilizer broadcasting unit. **DO NOT WATER TREATED AREA FOR 48 HOURS AFTER APPLICATION.**

**FOR ORNAMENTAL GARDENS, PLANTED BORDERS AND GROUND COVERS:** To control Snails and Slugs—Water area to be treated thoroughly before application. Withhold watering for 48 hours after treatment. Apply 1 pound to 1500 square feet. Apply over the infested area and around the edges of border or ground cover plants. Apply also to the bases of fences and walls and in shady protected areas. Repeat as necessary.

### **FOR DICHONDRA AND GRASS LAWNS:**

To Control Snails and Slugs—Water area to be treated thoroughly before application. Withhold watering for 48 hours after treatment. Apply 1 pound to 1500 square feet.

To Control Ants, Earwigs, Fall Armyworms, Sod Webworms (Lawn Moths) and Sowbugs—Apply 1 pound to 250 square feet. Follow watering practice as outlined above. Repeat treatment as necessary.

**WARNING:** Harmful if swallowed or inhaled. Do not take internally. Harmful dust. Avoid breathing dust. **Skin contact may be harmful.** Avoid contact with skin, eyes and clothing. Wash hands and face after handling. Wear clean clothing. **Note for Physician:** Carbaryl is a moderate, reversible Cholinesterase inhibitor. Atropine is antidotal. Avoid contamination of feed and foodstuffs. Baits should be spread in such a way that children, pets or farm animals cannot reach them. Do not store in direct sunlight. Dispose of container promptly when empty.

This product is toxic to wildlife. Birds feeding in treated areas may be killed. Keep out of any body of water. Do not apply to any area not specified on the label.