


BUX[®] 2 Emulsive



ORTHO

(INSECTICIDE)

Active Ingredients	By Wt.
m-(1-methylbutyl) phenyl methylcarbamate	17.4%
m-(1-ethylpropyl) phenyl methylcarbamate	5.8%
Aromatic Petroleum Derivative Solvent	42.0%
Inert Ingredients	34.8%

Contains 2 pounds BUX Insecticide per gallon.
U. S. Pat. Nos. 3,062,864—3,062,867—3,553,328

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS

WASH AND DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.

NET CONTENTS 5 GALS.

Chevron Chemical Company

Ortho Division/San Francisco, Calif. 94119

Richmond, California Fresno, California

Des Moines, Iowa Perth Amboy, New Jersey Orlando, Florida

Form 7924 G EPA Reg. No. 239-2322 Product 5036 Made in U.S.A.

DIRECTIONS

CORN (Field) (Popcorn Sweet)—For Control of Corn Rootworm Larvae, ROW TREATMENT—Use 3 to 4 pints of BUX 2 Emulsive in 10 gallons of spray mix per 13,080 linear feet of row. This is equivalent to one acre of corn on a 40-inch row spacing. To apply either the 3 or 4 pint rate of BUX 2 Emulsive regardless of row spacing, one pint of the desired spray mix should be delivered per 164 feet of row at normal planting speed. Apply the spray mix at the time of planting in a 7 to 10-inch band to the soil surface over the covered seed row. Incorporate the spray mix into the top ½ to 1 inch of soil using press wheels, covering disks, or trailing chains. **DOUBLE BAND (SPLIT BOOT) TREATMENT**—

Apply the row treatment rate per acre in combination with the required amount of liquid fertilizer in a band 1 to 2 inches to each side of the seed, 1 to 2 inches deep and 1 inch wide. Do not allow BUX 2 Emulsive to contact the seed as this may result in germination damage. Do not apply BUX 2 Emulsive to only one side of the seed as this may result in poor control. **BROADCAST TREATMENT**—Use 3 to 4 gallons of BUX 2 Emulsive per acre in 15 to 20 gallons of spray mix with ground equipment. Apply as a uniform broadcast treatment prior to or at the time of planting, incorporate into the top one-half to one inch of soil using a harrow or disk.

CORN (Field)—For Tank Mixing With Atrazine 80W: Use 6 pints of BUX 2 Emulsive in 10 to 20 gallons of spray mix as an at-planting treatment in a 14 to 20 inch band to the soil surface over the covered seed row per 13,080 linear feet of row. Use in combination with Atrazine 80W, atrazine, applied at the rate of 1.5 to 2.0 lbs. per acre in 10 to 20 gallons of spray mix. Do not incorporate into the soil using press wheels, covering disks, or trailing chains. For a complete listing of approved combinations and application rates, consult the Atrazine 80W label.

CONDITIONS OF SALE: 1. Chevron Chemical Company (Chevron) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Critical and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability, except those assumed by Chevron under 1 above, in the use of this material.



DO NOT TAKE INTERNALLY
DO NOT INHALE
DO NOT GET ON SKIN

KEEP OUT OF REACH OF CHILDREN

WARNING: Harmful if swallowed. Do not get on skin or in eyes. Do not inhale vapor. Wash skin thoroughly with soap and water. For eyes flush with water, get medical attention. Wash thoroughly after using. Launder contaminated clothing before re-use. Do not store or transport near food or food. **DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.**

Note to Physicians. Atropine is Antidotal.

This product is poisonous to fish and wildlife. Keep out of lakes, ponds, and streams. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply in any manner not specified on the label!