



# PHALTAN<sup>®</sup> Rose Black Spot Control

NET CONTENTS  
1 QUART LIQUID

**Prevents Rose Black Spot,  
Powdery Mildew and Certain  
Other Plant Diseases**

**Makes up to 64 Gallons Diluted Spray  
Easy-To-Use Liquid**

|                   |        |
|-------------------|--------|
| Active Ingredient | By Wt. |
| Folpet*           | 27%    |
| Inert Ingredients | 73%    |

\*N-(trichloromethylthio)phthalimide

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS.

#### DIRECTIONS

ORTHO PHALTAN Rose Black Spot Control is an easy-to-use liquid fungicide designed to control Black Spot, Powdery Mildew and other listed diseases on ornamentals. Refer to table below for recommended rates and timing. For best results, apply with an ORTHO SPRAYETTE or power sprayer. Do not store diluted spray.



READ THE LABEL

KEEP OUT OF REACH OF CHILDREN

#### CAUTION:


Avoid inhalation of spray mist. Avoid contact with skin. Wash hands after use. Avoid contamination of feed or foodstuffs. ORTHO PHALTAN Rose Black Spot Control should not be used in combination with or closely following oil sprays. Under conditions of high temperature and humidity, slight unobjectionable injury to immature rose foliage may occur. The addition of spreading and sticking agents may result in injury. Dispose of container promptly when empty. This product is toxic to fish. Keep out of lakes, streams or ponds. Apply this product only as specified on this label.

| PLANT                                      | DISEASES  | COMMENTS   |
|--|---|--|
| Roses                                      | Black spot,<br>Powdery<br>Mildew  | Use 1 Tablespoonful (1 1/2 fl. oz.) to 1 gal. water. Apply when young leaves first appear. Repeat at 7 to 10 day intervals or as necessary throughout season.  |
| Zinnias,<br>Marigolds,<br>Snap-<br>dragons | Leaf Spot<br>Alternaria,<br>Powdery<br>Mildew<br>Erysiphe,<br>Anthracnose,<br>Powdery<br>Mildew | Use 1 1/2 Tablespoonfuls (1 1/2 fl. oz.) per 1 gal. water. Treat at least weekly during periods of rapid new growth and severe infection and twice weekly during periods of rapid new growth and severe infection caused by continuous rains or high humidity. |

COMBINATION INSECT AND DISEASE CONTROL SPRAY

**FOR INSECT CONTROL AND LISTED DISEASES:** Add 1 Tablespoonful ISOTOX Insect Spray for each gallon of diluted spray. **FOR ROSES:** Apply ORTHO PHALTAN Rose Black Spot Control at regular intervals for disease control; add ISOTOX Insect Spray only when necessary for insect control.

**NOTICE:** Buyer assumes all responsibility for safety and use not in accordance with directions.



6239-2329

## CHEVRON CHEMICAL COMPANY

FORM 8150 A

EPA Reg. No. 239-2329

ORTHO DIVISION — San Francisco, California 94119

Product 5039

Made in U.S.A.

RICHMOND, CALIF.

LAFAYETTE, CALIF.



DES PLAINES, ILL.

ATLANTA, GA.

PERTH AMBOY, N.J.



# PHALTAN<sup>®</sup> Rose Black Spot Control

NET CONTENTS  
1 QUART LIQUID

**Prevents Rose Black Spot,  
Powdery Mildew and Certain  
Other Plant Diseases**

**Makes up to 64 Gallons Diluted Spray  
Easy-To-Use Liquid**

|                   |        |
|-------------------|--------|
| Active Ingredient | By Wt. |
| Folpet*           | 27%    |
| Inert Ingredients | 73%    |

\*N-(trichloromethylthio)phthalimide

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS.

#### DIRECTIONS

ORTHO PHALTAN Rose Black Spot Control is an easy-to-use liquid fungicide designed to control Black Spot, Powdery Mildew and other listed diseases on ornamentals. Refer to table below for recommended rates and timing. For best results, apply with an ORTHO SPRAYETTE or power sprayer. Do not store diluted spray



**READ THE LABEL**

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION:** Avoid inhalation of spray mist. Avoid contact with skin. Wash hands after use. Avoid contamination of feed or foodstuffs. ORTHO PHALTAN Rose Black Spot Control should not be used in combination with or closely following oil sprays. Under conditions of high temperature and humidity, slight unobjectionable injury to immature rose foliage may occur. The addition of spreading and sticking agents may result in injury. Dispose of container promptly when empty.


This product is toxic to fish. Keep out of lakes, streams or ponds. Apply this product only as specified on this label.

| PLANT                                      | DISEASES   | COMMENTS  |
|--|--|---|
| Roses                                      | Black Spot,<br>Powdery<br>Mildew   | Use 1 Tablespoonful (15 ml, oz) to 1 gal. water. Apply when young leaves first appear. Repeat at 7 to 10 day intervals or as necessary throughout season.   |
| Zinnias,<br>Marigolds,<br>Snap-<br>dragons | Leaf Spot<br>Alternaria<br>Powdery<br>Mildew<br>(Erysiphe),<br>Anthracnose,<br>Powdery<br>Mildew | Use 1½ Tablespoonful (22.5 ml, oz) per 1 gal. water. Treat at least weekly during periods of rapid new growth and severe infection and twice weekly during periods of rapid new growth and severe infection caused by continuous rain or high humidity. |

COMBINATION INSECT AND DISEASE CONTROL SPRAY

**FOR INSECT CONTROL AND LISTED DISEASES:** Add 1 Tablespoonful ISOTOX Insect Spray for each gallon of diluted spray. **FOR ROSES:** Apply ORTHO PHALTAN Rose Black Spot Control at regular intervals for disease control; add ISOTOX Insect Spray only when necessary for insect control.

**NOTICE:** Buyer assumes all responsibility for safety and use not in accordance with directions.



6662-2329  
2329-662

## CHEVRON CHEMICAL COMPANY

FORM 8150 A

EPA Reg. No. 239-2329

ORTHO DIVISION — San Francisco, California 94119

Product 5039

Made in U. S. A.

RICHMOND, CALIF.

LAFAYETTE, CALIF.



DES PLAINES, ILL.

ATLANTA, GA.

PERTH AMBOY, N.J.