

P
10-19-71

FEDERAL REGISTER
239-2383

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS.

DIRECTIONS

Repeat treatment as necessary. Do not apply to food crops within 30 days of harvest unless otherwise specified. Remove residues remaining at harvest by washing, wiping or other effective means. ORTHO Chlordane 8 Emulsive XB may be used in airplane or power sprayers.

USE DESIGNATED AMOUNT OF PRODUCT IN THE FOLLOWING GALLONS WATER PER ACRE UNLESS OTHERWISE NOTED:
GROUND APPLICATION—Vegetable and Field Crops—30 to 250 gals.
Where Recommended for AIR APPLICATION—Field Crops—3 to 10 gals.

ANTS (Outside Use): Saturate nests and spot treat infested areas with 1 pint to 50 gallons water. Do not treat food and feed crops.

CABBAGE, CAULIFLOWER, BROCCOLI, BRUSSELS SPROUTS, TURNIPS: Cabbage Maggot (Make pre-planting soil applications at time of planting or transplanting only): Use 1½ pints to 300 gallons of water per acre.

FIELD AND VEGETABLE CROPS (Peanuts, Bean, Beet, Cantaloupe, Corn, Cucumber, Eggplant, Lettuce, Melons, Potato, Tomato, Cabbage): Pre-Plant Soil Treatment — BROADCAST — Wireworms, White Grubs, Cutworms, Seed Corn Maggots, Diabrotica Larvae—2 qts. per acre in water to cover. Work into upper few inches of soil.

BEANS, CABBAGE, CORN, KALE, ONIONS, OKRA, PEAS, STRAWBERRIES, TOMATOES: Grasshoppers and Crickets—Use 1 pint in sufficient water to cover 1 acre. **FOR GRASSHOPPERS:** Set spray high enough to spray top of the foliage. Do not apply to green or snap Beans after pods start to form. Do not apply to Cabbage, Kale, Okra, Strawberries or Tomatoes after the edible parts start to form. Do not use on green or spring Onions. Do not apply to Peas within 15 days of harvest. Apply primarily to silks of sweet Corn. Do not feed treated forage to dairy animals or animals being finished for slaughter.

BEETS, PEAS: Leaf Miners—Use 1 pint in 100 to 200 gals. water per acre as soon as insects appear. Do not apply to Beets after seedling stage if tops are to be used for food. Do not apply to Peas within 15 days of harvest. Do not feed treated forage to dairy animals or animals being finished for slaughter.

ONIONS: Onion Maggots—(Make pre-planting soil applications at time of planting or transplanting only): Use 1 quart in 50 to 100 gals. water per acre as a spray in a 6 inch band. Do not use on green or spring onions.

TERMITES: 1 gallon in 100 gals. water. Apply in trench 30 inches deep, but not below the top of the footing if this is shallower. Trench must extend all the way around the building being treated. Use 1 gal. of the mixture per lineal foot of trench including treatment of backfill soil. Also soak soil under and around porches and around each of the piers under houses.

ONIONS AND GLADIOLUS: Thrips—Use 1 pint in 100 to 200 gals. water per acre as soon as insects appear. Make 2 to 3 applications at 7 to 10-day intervals if necessary. Do not use on green or spring onions.

WARNING

Contact with skin wash with soap and water. Do not store or transport OR STORE NEAR this material should be finished for slaughter. drift to food, for crops thereof

Toxic to Fish and Aquatic Life. Do not contaminate water where runoff is likely. Do not use on equipment, or discharges exposed to direct sunlight may be obtained.

CONDITIONS OF WARRANTY: That this label and is rec neither makes, nor other warranty of representation, express or implied, and prevent it from being liable. Such risks in crops to which the damage caused by though the product even though label and assume all risk under 1 above material.

FIELD AND VEGETABLE CROPS (Peanuts, Bean, Beet, Cantaloupe, Corn, Cucumber, Eggplant, Lettuce, Melons, Potato, Tomato, Cabbage): **Pre-Plant Soil Treatment — BROADCAST — Wireworms, White Grubs, Cutworms, Seed Corn Maggots, Diabrotica Larvae**—2 qts. per acre in water to cover. Work into upper few inches of soil.

BEANS, CABBAGE, CORN, KALE, ONIONS, OKRA, PEAS, STRAWBERRIES, TOMATOES: Grasshoppers and Crickets—Use 1 pint in sufficient water to cover 1 acre. **FOR GRASSHOPPERS:** Set spray high enough to spray top of the foliage. Do not apply to green or snap Beans after pods start to form. Do not apply to Cabbage, Kale, Okra, Strawberries or Tomatoes after the edible parts start to form. Do not use on green or spring Onions. Do not apply to Peas within 15 days of harvest. Apply primarily to silks of sweet Corn. Do not feed treated forage to dairy animals or animals being finished for slaughter.

BEETS, PEAS: Leaf Miners—Use 1 pint in 100 to 200 gals. water per acre as soon as insects appear. Do not apply to Beets after seedling stage if tops are to be used for food. Do not apply to Peas within 15 days of harvest. Do not feed treated forage to dairy animals or animals being finished for slaughter.

ONIONS: Onion Maggots—(Make pre-planting soil applications at time of planting or transplanting only): Use 1 quart in 50 to 100 gals. water per acre as a spray in a 6 inch band. Do not use on green or spring onions.

TERMITES: 1 gallon in 100 gals. water. Apply in trench 30 inches deep, but not below the top of the footing if this is shallower. Trench must extend all the way around the building being treated. Use 1 gal. of the mixture per lineal foot of trench including treatment of backfill soil. Also soak soil under and around porches and around each of the piers under houses.

ONIONS AND GLADIOLUS: Thrips—Use 1 pint in 100 to 200 gals. water per acre as soon as insects appear. Make 2 to 3 applications at 7 to 10-day intervals if necessary. Do not use on green or spring onions.

WARNING: Harmful if swallowed. Avoid breathing spray mist. Contact with skin can cause toxic symptoms. In case of contact with skin, wash with soap and water.

Do not store or transport near feed or food. **DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.** Forage crops treated with this material should not be fed to dairy animals or to animals being finished for slaughter. **Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.**

Toxic to Fish and Wildlife. Birds feeding on treated areas may be killed. Do not contaminate any body of water or apply to areas or crops other than those for which directions are given on the label. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

CONDITIONS OF SALE: 1. Chevron Chemical Company (Chevron) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty of **FITNESS** or of **MERCHANTABILITY**, guarantee or representation, express or implied, concerning this material.

2. Critical and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by Chevron under 1 above) resulting from handling, storage, and use of this material.



hful if swallowed. Avoid breathing spray mist.
toxic symptoms. In case of contact with skin,

ar feed or food. DO NOT USE, POUR, SPILL
R OPEN FLAME. Forage crops treated with
fed to dairy animals or to animals being
ot apply under conditions involving possible
er plantings that might be damaged or the
for sale, use or consumption.

Birds feeding on treated areas may be killed.
dy of water or apply to areas or crops other
ions are given on the label. Do not apply
ur. Do not contaminate water by cleaning of
wastes. This product is highly toxic to bees
or residues on crops. Protective information
r Cooperative Agricultural Extension Service.


1. Chevron Chemical Company (Chevron)
conforms to the chemical description on the
fit for use as directed hereon. Chevron
es any agent or representative to make, any
or of MERCHANTABILITY, guarantee or rep-
plied, concerning this material.

ossible factors beyond Chevron's control
all risks in connection with the use of chemi-
ut are not limited to, damage to plants and
al is applied, lack of complete control, and
o other plants or crops. Such risks occur even
sonably fit for the uses stated hereon and
as are followed. Buyer and user acknowledge
liability (except those assumed by Chevron
from handling, storage, and use of this

[INSECTICIDE] XYLENE BASE SOLVENT

Active Ingredients	By Wt.
Technical Chlordane*	71%
Xylene Base Aromatic Petroleum Solvent	23%
Inert Ingredients	6%
*Equivalent to 42.6% Octachloro-4,7-Methanotetrahydroindane and 28.4% related compounds. Contains 8 lbs. Chlordane per gallon.	

ACCEPTED
15-19-71
UNDER THE FEDERAL INSECTICIDE
REGISTRATION ACT
FOR THE CONTROL OF PESTS
REGISTERED UNDER THE
FEDERAL INSECTICIDE
REGISTRATION ACT



READ THE LABEL

WARNING: KEEP OUT OF REACH OF CHILDREN

SEE BACK PANEL FOR ADDITIONAL WARNINGS.

WHEN CONTAINER IS EMPTY, IMMEDIATELY WASH THOROUGHLY AND DESTROY. NEVER RE-USE.

NET CONTENTS 1 GALLON LIQUID



REGISTRATION NO. 246-142

ACTIVE INGREDIENTS
Cyfluthrin 1.0%

INERT INGREDIENTS

NET WT

Equivalent to 5% Cyfluthrin in 100% water-soluble formulation
and its related compounds

246-142

[CAUTION] 18 point type size

[KEEP OUT OF THE REACH OF CHILDREN] 12 point type size

INSTRUCTIONS FOR USE

FRESH POTATOES AND SWEET POTATOES: For control of Wireworms, White Grubs and Flea Beetles ^{larvae} apply 40 to 60 pounds per acre to the soil then harrow or disc after applying.

For control of White Grubs, Wireworms, White Fringed Beetle Larvae, Japanese Beetle Larvae, Green June Beetle Larvae, 50 pounds per acre work into the top few inches of the soil. For control of White Grubs and Wireworms apply 40 to 60 pounds per acre. Keep children and pets off treated areas until material has been washed into soil and foliage is thoroughly dry.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be harmed. Keep out of lakes, ponds and streams. Do not apply to any area not specified on the label.

U.S. D.H. REG No 246-

Net Weight 50 lbs
Inert Ingredients:
Glycolic acid 56.5%
Pyrophosphate 33.5%

Net weight 50 lbs