PLICTRAN 50 Wettable Miticide EPA Reg. No. 239-2414 Submitted: March 16, 1987

COPY A

Harmful

COPY D

reversible

#### COPY C

When mixing, loading and applying, wear chemically resistant gloves.

## COPY D

į

This product is toxic to fish ar.: wildlife. Do not apply directly to water. Do not apply where run off is likely to occur. Cover or incorporate spills. Do not apply when weather conditions favor drift from areas treated.

EPA Reg. No.

## COPY E

IF SWALLOWED, give water or milk to drink and telephone for medical advice. Consult medical personnel before inducing vomiting. If medical advice cannot be obtained, then take person and product container to the nearest medical emergency treatment center or hospital.

IF IN EYES, wash eyes immediately with fresh water for at least 15 minutes and see a doctor.

TF ON SKIN, wash skin with plenty of soap and water. See a doctor if irritation persists.

IF INHALED, remove person from exposure area.

#### COPA £

500

# PRECAUTIONARY STATEMENTS IAZARDS TO HUMANS & DOMESTIC **NIMALS**

#### VARNING

ay-be-harmful it shallowed. May cause skin and respiratory irritation. Causest to damage. Do not get in eyes, on skin or on clothing. Wear goggles or a use mask when pouring product from bag. Do not breathe dust or spray mist, uting commercial application or prolonged exposure as in spray mixing and adding operations use goggles and a face mask. Cytash thoroughly after andting and spraying. Do not wear contaminated clothing.

#### ENVIRONMENTAL HAZARDS

his product is toxic to lish. Do not apply directly to water. Do not apply where until is likely to occur. Do not apply when weather conditions favor drift from teas freaten.

<u>-0294</u>

#### RECTIONS FOR USE

is a violation of Federal law to use this product in a manner inconsistent with

AD ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL ULATIONS.

#### REENTRY STATEMENT

sp all unprotected persons out of operating areas or vicinity where there may be

net enter treated areas without protective clothing until sprays have dried. clain states may require more restrictive reentry intervals; consult your State De-timent of Agriculture for further information.

itten or oral warmings regarding use of protective clothing and accidental ex-sure must be given to workers who are expected to be in treated areas or in areas out to be treated.

#### STORAGE AND DISPOSAL

#### **PROHIBITIONS**

Do not contaminate water, food, or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited.

#### STORAGE

Keep pesticide in original container.
Do not store or transport near feed or food.
Do not store in areas accessible to children.
For help with any soill, leak, fire or exposure involving this material, call day or night (415) 233-3737.

#### PESTICIDE DISPOSAL

This resticide is acutely hazardous. Improper disposal of excess posticide, spray mixture, or rinsate is a signation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

#### CONTAINER DISPOSAL

Completely empty bag into application equipment. Do not reuse container. Dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

#### IRECTIONS

INCEPTIONS

(ICITAN 50 Wellable Miticide is recommended for control of motile forms of lant teeding mittes, including strains both susceptible and tesistant to other siticides. A wide range of species are controlled as indicated for each erop in the Table. PLICTRAN 50W is not highly injurious to certain predatory mites and sects and is essentially nontoric to hone-bees, PLICTRAN 50W is compatible the Insecticides and Jungacides commonly recommended orchard and ornasental spray schedules except for oil. (See Application Rotes.)

sental spray schedules except for oil. (See Application Roles.)

own to Use: PLICIRAN 50 Weltsble Mittelde mixes readily with water to form
suspension that can be applied with conventional spray equipment. PLICIRAN
OW contains a metting agent. Additional welling agent is not recommended,
ther dilute or concentrate spray, can be used. Follow the dosage recommendons in the Vable. Thorough and complete coverage of foliage and fruit is essential
if good mite control. Do not graze or feed brestock on cover crops growing in
reated areas. Not for use in or around the home.



# PLICTRAN® 50 Wettable

Active Ingredient	By Wt.
Cyhexatin (tricyclohexylhydroxystannane)	50%
Inert Ingredients	50%

WETTABLE POWDER FORMULATION FOR CONTROL OF PLANT FEEDING MITES

PLICTRAN® - Reg. TM of The Dow Chemical Company for Cyhexatin Miticide.

# KEEP OUT OF REACH OF CHILDREN WARNING-AVISO

#### STATEMENT OF PRACTICAL TREATMENT

T SWALLOWED give a large amount of water to drink, make parson romit

IF-IN-EYES immediately flush-eyes with fresh-water for 15 minutes and see

IF-ON-SKIN-wach-skin-with-plenty of-soap and water, See & doctor If irritation

Note to Physicians: Emergency Information - call (415) 233-3737.

FOR AGRICULTURAL USE ONLY SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

Manulactured for

### Chevron Chemical Company 5 1984

Ortho Agricultural Chemicals Division San Francisco CA 94129-7144 Richmond CA 94804-0036

Product 1000 Made in U.S.A.

R Form 8761-L EPA Reg. No. 239-2414-AA

BEST AVAILABLE COPY

_Crop	Miles Controlled
Apples	European Red Twospotted Spider Schoena Spider Pacilic Spider Vellow Spider Apple Rust McDaniel
Pears	European Red Twospotted Spider Schoene Spider Pacific Spider Yellow Spider Apple Rust McDaniel Pear Rust
Citrus	Citrus Red Texas Citrus Citrus Rust Pacific Spider
Orna- mentals	Thospotted Spider
Peaches Plums (Prunes) Ne: Larines	European Red Twospotted Spider Peach Silver McDaniel Pacific Spider
Alinonds Walnuts	Pacific Spider European Red Twospotted Spider
Straw berries	Twospotted Spider
Hops	Twospolted Spider

Application Notes:

1. Make the first application at or soon after petal fall, of water per acce and app dittons to provide thorough 2. On not apply a total "" as "" on apples by groun season on pears by ground to not "hardened of!" since will occur. Application of time period between the a "hardened of!", normally of minury to the fruit.

d. Includes greenhouse grown 9. Occasional injury to chryscured. Therefore, it is des of these plants (prehloom 10. Do not apply more than 11. "I've the higher rates in pl. times on the periods of intense mit. 12. Oo not harvest treated be 13. Oo not tank mix PLICIR/SOW and summer spray foldar and fruit injury will Apply a minimum of 24 or 15. Use 2.2 lbs. in 150 gals. in 225 gals. of water peapply more than 9 lbs. P within 21 days of harvest.

Do not apply under conditions plantings that might be dama use or consumption.

CONDITIONS OF SALE: 1 ch this material conforms to the fill for use as directed hereon, or representative to make, any guarantee or representation, et 2 Critical and unforeseeable