ORTHO Vegetable & Garden Fungicide EPA Reg. No. 239-2438 Submitted: April 5, 1985

COPY A


PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

WARNING: Causes eye and skin irritation. This product may produce temporary allergic side effects characterized by redness of the eys, mild bronchial irritation and redness or rash on exposed skin areas. Do not get in eyes, on skin or on clothing. Wear long sleeve shirt, long pants, and rubber gloves while mixing and applying this product. Wash skin and hands thoroughly after using. Remove and wash contaminated clothing before reuse. Keep children and pets off treated area until dry.

COPY B

ENVIRONMENTAL HAZARDS: This product is toxic to fish and other aquatic organisms. Do not apply directly to water or dispose of wastes in water. Do not apply when weather conditions favor drift of spray. Do not contaminate water by cleaning of equipment or disposal of wastes.



CONFIDENTIAL

PROPERTY OF
CHEVRON CHEMICAL COMPANY
UNTIL ACCEPTED BY EPA

COPY A

PRECAUTIONARY STATEMENTS TATABLE TO HUMANS & DOMESTIC ANIMALS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

WARNING: Course eye and skin irritation. Skin contact may cause allergic reaction. Do not get in eyes, on skin or on clothing. Wear subner gloves when handling. Wosh skin and hands thoroughly after being. Remove and wash contaminated clothing before reuse. Kenn children and nets all treated area until dry. STATEMENT OF "RACTICAL TREATMENT: In case of eye contact, immediately flux eyes with fresh water for 15 minutes and see a doctor. In case of skin contact, wash skin with soap and water. Get medical attention. I irritation persists. Note to Physicians Emergency. Information. call. (415), 233, 3737.

ENTROMMENTAL HAZARDS. This product is toxic to tash. Keep.

NYRONMENTAL HAZARDS. This product is toxic to wh Keep out of lokes, streams or ponds. Do not contaminate water by leaning of equipment or disposal of water.

DIRECTIONS FOR IJJE COPY B

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

READ ENTIRE LABEL, USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

ORTHO Vegetable & Garden Fungicide mixes readily with water and may be used in an ORTHO SPRAY ETTE, ORTHO Lawn & Garden Sprayer, tank type or power sprayer. Shake bottle well before using. Mix appropriate amount with water and use of once.

Apply at listed rate per crop for disease to be controlled [] in 1 gallon of water to cover on area of 200 sq. ft. (10° \times 20.) planted with listed crop. I TBS: I. Tablespoonlu , I. tsp. I. tea spoonful. Repeat at 7 to 14 day intervals. Use the 7 day schedule during severe disease conditions or wet whather unless otherwise stated. Note: Avoid spraying plants during extremely hat and sunny weather. Do not apply to wilted plants.

Do not graze treated areas or feed treated plant parts to livestock

VEGETABLES - Numbers in () indicate days required between last application and harvest

Beans (14) (Snap Beans) Rust [1 TBS] Begin in early bloom stage or when disease first threatens. Repeat at weekly intervals are as necessary.

Broccoli (0), Brussels Sprouts (0), Cabbage (0), Cauliflower (0): Alternaria Leafspot (Southeast only) [1½ tsp.], Downy Mildew [1½ tsp.] Begin after transplants are set in the field a when conditions favor disease development Repeat it 7 to 0 day intervals. Ring Spot on Brussels Sprouts (California only) [2 tsp.] Begin at time of early sprout development or when conditions favor disease development. Repeat at 7 to 10 day intervals.

Carrots (0) Early and Late Blight [2 tsp.] Begin when disease threatens and repeal at 7 to 10 day intervals or as required to maintain control.

Celery (7): Early and Late Blight [1½ tsp.], Basal Stalk Rot [1½ tsp.]. Begin when plants are set in the field. Repeat on a 4 to 7 day interval.

Corn (14) (Sweet): Helminthosporium Leaf Blight [2 Isp.) Begin when conditions favor disease development. Repeat at a 4 to 7 day interval or as needed. Do not apply to sweet corn for processing.

S ORTHO

128 Gallons
Diluted Spray

Vegetable & Garden Fungicide

Contains DACONIL 2787*

For the Control of Diseases on Vegetables, Roses and Flowers
(a. Gold on label)

CONTROLS BIGHT ON TOMATOES

Active Ingredient 8y Wt.

**Chlorotholonil (retrachloroisophthalonitrile) 29.6%
Inert Ingredients 70.4%

*DACONIL 2787 is a registered trademark of Diamond Shamrock Corp.

**Covered under U.S. Patent No. 3,290,353 and No. 3,331,735

Keep out of reach of children

WARNING

See side panel for additional precautionary statements.

NET CONTENTS 1 QT.

Cucumbers (0), Melons [2 tsp.], Gummy Ster Mildew [2 tsp.], Leaf E Cucumbers? [2 tsp.], States) on Cucumbers Rot (Rhizoctonia) [1 TE in the first true leaf st disease Repeat appliquently if required.

Potatoes (0) Early and [11/2 tsp.] Begin whe threatens Repeat at 7

Tomatoes (0) Anthrac tsp.), Gray Leafspot Rhizoctonia Fruit Rot [peat at 7 to 10 day conditions Alternaria [1 TBS.]--Make 3 appl 5 to 6 weeks before at

ROSES, FLOWERS

Roses: Blackspot [1 ts] young leaves first app the growing season Sp

Chrysanthemums: Ascocisp J. Apply at first siggrowing season

Geraniums: Botrytis Blig ing wet weather or sev

Iris: Botrytis Blight [155] tions when leaves are a Zinnias. Powdery Milde at the first sign of dise

STORAGE AND DISPO STORAGE. Keep pestici centrate or dilute into I tion of feed and foodst in a locked storage are DISPOSAL PRODUCT – by securely wrapping o paper and discard in bottle. Rinse thoroughly

NOTICE Buyer assume in accordance with dire

Chevron Chemical (

Ortho Consumer Products San Francisco CA 94120 7 Product 1000 Made Form 9067 G EPA Reg. No 239 2438

© © EPA Est 239 CA 1, 239 NJ