

FRONT PANEL

ORTHO

St. Augustine & Centipede
Weed and Feed

For use on St. Augustine
and Centipede lawns only

JUN 10 1990
234-2500

Controls: Lippia, Florida Pusley, Pigweed, Plantain and Certain Other
Broadleaf Weeds

Prevents: Sedge, Crabgrass, Spotted Spurge and Paspalums from Becoming
Established

Provides: Fast Acting and Long Lasting Fertilizer for a Green Lawn

Weeds and Feeds 3,000 Sq. Ft.

Money Back Guarantee

See Side Panel

Ingredient Statement as a Herbicide

Active Ingredient	By Wt.
*Atrazine	1.15%
Inert Ingredients98.85%

*Equivalent to 1.10% (2-chloro-4-ethylamino-6-
isopropylamino-S-triazine) and 0.05% related
compounds

Equivalent to 23.0 lbs. Atrazine per ton

12 pt. Keep out of reach of children.

18 pt. CAUTION

See back panel for additional precautionary statements.

NET WT. 12 LBS.

CONFIDENTIAL

PROPERTY OF

ORTHO CHEMICAL COMPANY

USE ACCEPTED BY EPA

LEFT
SIDE PANEL

ORTHO St. Augustine & Centipede
Weed and Feed

Covers 3,000 Sq. Ft.

RIGHT
SIDE PANEL

ORTHO St. Augustine & Centipede
Weed and Feed

Covers 3,000 Sq. Ft.

ORTHO'S MONEY BACK GUARANTEE

If you are not satisfied with the results after using this product,
send us this guarantee with your sales receipt. Your refund check
will be mailed to you promptly.

CHEVRON CHEMICAL CO., ORTHO CONSUMER PRODUCTS DIV., P.O. Box 3744,
San Francisco 94120-7144

BACK PANEL

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

12 pt. CAUTION

Avoid contact with eyes.

STATEMENT OF PRACTICAL TREATMENT

In case of eye contact, immediately wash eyes with fresh water for
15 minutes. If irritation continues, see a doctor.

Note to Physicians: Emergency Information - call (415) 233-3737.

GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen (N)	25%
1.1% Ammoniacal Nitrogen	
20.1% Urea Nitrogen	
3.8% Water Insoluble Nitrogen	
Available Phosphoric Acid (P ₂ O ₅)	3%
Soluble Potash (K ₂ O)	7%

Iron (Fe) 0.9%
Primary Nutrients from Urea, Urea Formaldehyde, Ammonium
Phosphate, and Muriate of Potash.

Trace Nutrient from Iron Sulfate

Chlorine, not more than 5.8%
Potential Acidity 900 lbs. Calcium Carbonate Equivalent per ton

Ingredient Statement as a Herbicide

Active Ingredient By Wt.

*Atrazine 1.15%

Inert Ingredients 98.85%

Equivalent to 1.10% (2-chloro-4-ethylamino-6-isopropylamino-S-triazine) and 0.05% related compounds.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS.

ORTHO St. Augustine & Centipede Weed and Feed rejuvenates starved and weedy lawns by:

1. Providing a balanced fertilizer that will give you a green lawn.
2. Controlling unsightly broadleaf weeds such as Plantain, Pennywort, Chickweed, Creeping Charlie, Evening Primrose, Florida Betony, Black Medic, Mouseear Chickweed, Dandelion, Oxalis, Carolina Geranium, Florida Pusley, Lippia, Henbit, Rabbit Tobacco, Pigweed, Toadflax, Crowfoot and Cudweed that compete with your grass for water and nutrients.
3. Preventing Sedge, Crabgrass, Spotted Spurge and Paspalums from becoming established in your lawn.

WHEN TO APPLY

1. To control the existing broadleaf weeds listed under number 2, above and fertilize grass, apply anytime the weeds are present and actively growing (usually early Spring is best time for application). Repeat in two months if needed. Length of time required to control established weeds is dependent upon their stage of growth. Mature weeds may take as long as 4 or 5 weeks to completely die.

2. To prevent the growth of Sedge, Crabgrass, Spotted Spurge and Paspalums and fertilize grass, apply in January, February or March, before these weeds germinate. This product will not control these weeds if they are already present in the lawn at the time of application.

In newly sprigged St. Augustine grass lawns, apply soon after planting to reduce competition from weeds.

HOW TO APPLY

The contents of this bag will cover 3,000 sq. ft. (100' x 30') when applied according to directions. If soil is dry, water lawn thoroughly the day before treatment. Treat entire lawn evenly. Do not exceed the recommended rate. Immediately after application, water lawn to wash material off the grass blades and into the soil. To avoid drift when using a WHIRLYBIRD Spreader or ORTHO Broadcast Spreader, apply when air is calm.

IMPORTANT NOTICE

ORTHO St. Augustine & Centipede Weed and Feed is for use on St. Augustine and Centipede lawns only. Do not use on Bluegrass, Fescue, Bent, Bahia, Rye, Bermuda or Dichondra lawns. For newly seeded Centipede, wait until the new grass has been mowed at least once before applying ORTHO St. Augustine & Centipede Weed and Feed. Delay sprig planting one month and seeding 6 months after application.

Avoid application under the branch spread or root zone area of trees, shrubs and ornamentals to prevent injury to plants. This should be particularly observed in compact, slowly drained soils especially in the Dallas/Fort Worth area. The root zone area extends from the trunk out to a point under the branch tips. Avoid application on garden plants, flowers and shrubs. Read and follow directions carefully.

FAST, EASY APPLICATION

(Illustrations)

ORTHO Broadcast Spreader*

Use setting No. 5. Swath width 8 ft. Apply 12 lbs to cover 3,000 sq. ft. (100' x 30'). Cover lawn in one direction only. Do not use criss-cross pattern. Due to wide swath, use only in open areas, avoiding trees and shrubs. To prevent overapplication and possible injury, spreader passes should be made 8 feet apart.

ORTHO WHIRLYBIRD Spreader*

Use setting No. 3. Apply 2 WHIRLYBIRDS full (1 lb.) to 250 sq. ft. (25' x 10'). Start cranking the WHIRLYBIRD handle before depressing the thumb lever. To ensure complete coverage, apply the second WHIRLYBIRD

full at right angles to the first. For application in confined areas and along border plantings, control of swath width can be obtained by holding the WHIRLYBIRD close to the ground or by cranking the handle in the opposite direction when using.

ORTHO Drop Spreader*

Use setting No. 5. Cover lawn in one direction only. Do not use criss-cross pattern. To ensure complete coverage, overlap wheel tracks. When making turns, shut off spreader to avoid overapplication.

*Variables inherent in equipment, as well as accuracy of machine adjustment, may cause application rates to deviate. Therefore, Chevron makes no warranty as to uniformity of coverage obtained from settings listed.

PESTICIDE-FERTILIZER MIXTURE containing 1.15% Atrazine.

rpt STORAGE AND DISPOSAL

Avoid contamination of foodstuffs, fish ponds, irrigation ditches or water used for irrigation. Do not store near feed, foodstuffs, seeds, fertilizer, insecticide or fungicides. Keep pesticide in original container. Keep bag closed when not in use.

Store in a dry place. Do not reuse empty container. Wrap container and put in trash collection.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

Chevron Chemical Company
Ortho Consumer Products Division
San Francisco CA 94120-7144
Richmond CA 94804-0036
Form 10184-A Product 5473 Made in U.S.A.
EPA Reg. No. 239-NEW EPA Est. 239-IA-1