

1003

FRONT PANEL
Makes Up To
48 Gallons Spray

ORTHO
Liquid SEVIN® Spray

Kills: Many kinds of Beetles, Bugs and Caterpillars, such as Japanese Beetles, Plant Bugs, Tent Caterpillars, Gypsy Moth, Bagworm, Armyworms and other Leaf Eating Insects

On: Ornamentals, Roses, Flowers, Shrubs, Shade Trees, Fruit Trees, Nut Trees, Vegetables and Lawns

Japanese Beetle, Stink Bug, Tent Caterpillar Active Ingredient By Wt.

Carbaryl
(1-Naphthyl N-methyl
carbamate)..... 22%
Inert Ingredients..... 78%

ACCEPTED
CONTENTS

.2 39 2500

Keep out of reach of children 10 p.c.
CAUTION - 14 p.c.
See back panel for additional precautionary statements.

CONFIDENTIAL

PROPERTY OF

CHEVRON CHEMICAL COMPANY

UNTIL ACCEPTED BY EPA

NET CONTENTS 1 QT.

031 11 11

CONFIDENTIAL

PROPERTY OF

BACK PANEL

ORTHO
Liquid SEVIN® Spray

CHEVRON CHEMICAL COMPANY

UNTIL ACCEPTED BY EPA
PRECAUTIONARY STATEMENTS 10 pt.

HAZARDS TO HUMANS & DOMESTIC ANIMALS 10 pt.

- 10 pt. ← **CAUTION:** Harmful if swallowed or absorbed through the skin. Avoid contact with eyes, skin or clothing. Keep children and animals off treated lawns until these areas are dry.
- 10 pt. ← **STATEMENT OF PRACTICAL TREATMENT:** If swallowed, give a large amount of water to drink, make person vomit, and call a doctor. In case of eye contact, immediately wash eyes with fresh water for at least 15 minutes. See a doctor if irritation persists. For skin, wash with plenty of soap and water. **Note to Physicians:** Emergency Information - call (415) 233-3737. Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.
- 10 pt. ← **ENVIRONMENTAL HAZARDS:** Do not apply directly to water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

DIRECTIONS: ORTHO Liquid SEVIN Spray mixes readily with water and may be used in all sprayers. Shake container well before using. Use 4 tsp. per gallon water or 4 fl. oz. in 6 gallons water, unless otherwise noted.

For best results, apply full coverage spray to all exposed parts of plants, especially upper and lower leaf surfaces. Start spraying when insects or their damage appear. Repeat at 7-10 day intervals or as needed, unless otherwise noted. Numbers in () indicate days required between last application and harvest.

ROSES, ORNAMENTALS, FLOWERS, SHRUBS, SHADE TREES: Japanese Beetles, Gypsy Moth, Certain Aphids, Bagworm, Leaf Beetles, Lace Bugs, Cankerworms, Pine Tip Moth, Webworms, Rose Chater, Flea Beetles, Plant Bugs, Certain Leafminers, Rose Bug, Grasshoppers, Tussock Moth, Periodical Cicada (17 year locust), Scale (Crawlers), Tent Caterpillars, Blister Beetles, Leafrollers & Exposed Sawflies and Thrips. **Note:** Do not use on Boston Ivy or Virginia Creeper.

Combination Spray with Fungicide on Roses: ORTHO FUNGINEX[®] Rose Disease Control may be mixed with this product when listed insect(s) and disease(s) are found on roses. Follow directions and precautions on both labels.

FRUIT TREES, BERRIES AND NUTS: Apples (1), Pears (1), Apricot (3), Peaches (1), Cherries (1), Plum (1), Raspberries (5), Strawberries (1), Citrus (5), Raspberries (7), Pecans (9), Walnuts (9): Apple Maggot, Plum Curculion, Apple Aphid, Leafrollers, Lesser Peach Tree

Borer, Oriental Fruit Moth, Codling Moth, Peach Twig Borer, Scale (Crawlers), Pear Psylla, Certain Leafminers, Mealy Plum Aoud, Fall Webworm, Caterpillars, Cherry Fruitworm, Citrus Citworm, Cherry Maggot, Grape Leafhopper, Leafhoppers, Strawberry Weevil, and Japanese Beetles; Make one application between bud break and bloom, a second application at petal fall and then at 3 to 4 week intervals for 3 to 4 months as necessary. **Apple Maggot** - For best control, apply at 10 day intervals. **Rose Apple Aphid** - Apply before leaves are curled. **Pear Psylla** - Apply when eggs hatch or nymphs are present. **Lesser Peach Tree Borer** - Spray limbs and tree trunks thoroughly, weekly during moth flight. **Scale Insects** - Apply when crawlers are present. **Note:** To avoid foliage injury on McIntosh apples and possible fruit thinning on all varieties, delay use until at least 35 days after bloom.

VEGETABLES: Asparagus (1); Beans (1); Cabbage (3); Broccoli (3); Cauliflower (3); Radish (3); Turnips/Greens (14); Roots (3); Kale (14); Carrots (1); Corn (1); Cucumbers (1); Melons (1); Squash (1); Lettuce/Head (3) Leaf (14); Spinach (14); Swiss Chard (14); Peas (1); Potato (1); Tomato (1); Eggplant (1); Pepper (1); Asparagus Beetle, Blister Beetle, Mexican Bean Beetle, Cucumber Beetle, Flea Beetle, Japanese Beetle, Leafhoppers, Thrips, Cutworm, Armyworm, Corn Earworm, European Corn Borer, Stink Bugs, Caterpillars, Plant Bugs, Imported Cabbageworm, Spittlebug, Pickleworm, Melonworm, Squash Bug, Colorado Potato Beetle, Pea Weevil, Lace Bugs, Tomato Fruitworm, Tomato Hornworm.

LAWN INSECTS: Fleas, Ticks, Ants, Chiggers, Sod Webworm (Lawn Moth), Armyworm, Cutworm, Grasshopper, Earwig - Use 8 fl. oz. (1 cup) in 12 gallons water and cover 600 square feet of established lawn. For best results, use an ORTHO Lawn Sprayer. Mow lawn first and apply after rain or watering. Do not water for at least two days after application. Repeat in 2 or 3 weeks or as necessary.

STORAGE AND DISPOSAL 10 pt.

STORAGE: Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Avoid contamination of food and foodstuffs. Store in a cool, dry place, preferably in a locked storage area. Do not store diluted spray. **DISPOSAL:** **PRODUCT** - Partially filled bottle may be disposed of by securely wrapping original container in several layers of newspaper and discard by trash. **CONTAINER** - Do not reuse empty bottle. Place in original container before discarding to trash.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with the label.

FUNGICIDE (2) Reg. U.S. Pat. & TM. Off. 1974. EPA Reg. No. 105-105-0101 for Trifluralin Fungicide.

SEVIN (2) Reg. U.S. Pat. & TM. Off. 1974. EPA Reg. No. 105-105-0101 for Carbaryl Insecticide.

Chevron Chemical Company
New York, New York, Dept. 100
Sales Dept., P.O. Box 127, New York, New York 10010
Dept. 100, P.O. Box 127, New York, New York 10010
Tel. 212-777-1000
EPA Reg. No. 105-105-0101

© 1974 CHEVRON CHEMICAL COMPANY
All rights reserved. No part of this publication may be reproduced without permission in writing from Chevron Chemical Company.

© 1974 CHEVRON

PROPERTY OF

CHEVRON CHEMICAL COMPANY

MADE ACCURATELY BY EPA

FOR DOCUMENT AVAILABILITY

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

d. Add to your Directions For Use section the statement "Do not rotate to any crop not listed on this label or other carbaryl labels."

3. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 Enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely yours,

Jay S. Ellenberger
Product Manager (12)
Insecticide-Rodenticide Branch
Registration Division (TS-767)

Enclosure