Page 17 15

Please read instructions	on reverse before completing form.		⊬ ···· · For	m Approved.	OMB No. 207	0-0060.	Approval expires 05-31-98
\$EPA	Environmental Protection Washington, DC 204	on Agei	1,,4 (*) et 1 CY		Registrat Amendm Other		OPP Identifier Number
	Application	on for F	esticide ·	- Section			
1. Company/Product Nur 239-2513	nber	-=	2. EPA Prode Denni:	ıct Manager s Edwards, J	r.	1	posed Classification
4. Company/Product (Na 0KTH0-KLC	me) R Soil Insect & Termite Killer		PM#	19			None Restricted
SOLARIS O P. O. Box San Ramor	Applicant (Include ZIP Code) Froup of Monsanto Company 5006, 1, CA 94583-0906 this is a new address	Saci	(b)(i), my pi to:	No	ilar or identic		FIFRA Section 3(c)(3) mposition and labeling
		000	<u></u>				
Amendment - Exp	response to Agency letter dated		_ Age	ency letter dat a Too" Applica	ls in response ted NOTIFIC		<u> </u>
Notification - Exp	lain below.	-	Oth	er - Explain be	alow.307 4	įąci:	
every rec	er Spray reflecting these changes. puirement of PR Notice 95-2.	· .	0-995-01. 1 11	is is to cer	tify that th	is notif	ication meets
1. Material This Product				· · · · · · · · · · · · · · · · · · ·		<u>.</u>	1
Child-Resistant Packagin Yes* No * Certification must be submitted 3. Location of Net Conte	Yes No If "Yes" Unit Packaging wgt. No. per container.	If "Yes Packag	e wgt (No. per container	2. Type of C	Metal Plastic Glass Paper Other (S	The section of the se
Label	च्छित पुराविकार्यक्ष हात्रेज पूर्वकर विकास है है। असे Container अस्त्रेली के अस्त्रेल कर				On Label	. : '	panying product
6. Manner in Which Labo	el is Affixed to Product Lithon Paper Stenc	graph r glued ciled	· · · · · · [Other			Frague (CAT) (CAT)
	'		ion - IV	* .	1		12
1. Contact Point (Comp	lete items directly below for identificati	on of indivi	dual to be co	ntacted, if nec	essary, to pro	cess this	application.)
Name F. X. Kai	uenski	Title Re	gistration S	Specialist		Celephon	No (ligotude Area Code)
	Certific tatements I have made on this form and at any knowingly false or misleading sta able law.	d all attach					6: Dats Application Received (Stamped)
2. Signature 3. Title							CAUCITY CAUCITY
trances	X. Lamerske	Re	gistration S	Specialist		<u>-</u>	
4. Typed Name	X. KAMIEUSKI	5. Date	9/1	6/96			inger de de la companya de la compa



ITF CODE IS 910801

STORAGE AND DISPOSAL STORAGE. Store in a secure, cool dry place. Keep from freezing, DISPOSAL: Do not reuse empty container. Rinse thoroughly, wrap in newspaper and put in trash.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

WARNING: Causes substantial but temporary eye injury. Harmful if swallowed or if on skin, Do not get in eyes or on clothing. Wear goggles or safety glasses. Avoid contact with skin, Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

People and pets may re-enter treated areas after sprayed area has dried.

area has died.

FIRST AID: If IN EYES: Hold eyelids open and flush with a steady, gentle stream of water. Get medical attention, IF SWALLOWED, if the person is alert and able to swallow, give small sips of milk or water. Call a physician or poison control center for further treatment advice. IF ON SKIN, Wash with plenty of soap and water. Get medical attention if intration persists. Note to Physicians: Emergency Information call 1-800-484-2333. Chloropyrifos is a cholmesterase inhibitor. Atropine is antidottal, 2-PAM is artidottal and may be used in conjunction with atropine, but should not be used alone.

ENVIRONMENTAL, HAZARDS. This product is took to fish and whickle, Do not apply directly to water or contaminate water when disposing of ecouprient washwater or disposal of waste. This product is highly took to bees exposed to direct treatment area.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.



Medical Information call 1-800-454-2333 Product Information call 1-800-225-2883

Manufactured for The SOLARIS Group of Monsanto Company P.O. Box 5008 San Ramon CA 94583-0808 Form 9-108-01 / Product 6021 EPA Reg. No. 239-2635 EPA Est. 239-14-3', 58996-MO-14 Superscript is first letter of lot number

Made in USA

[ICON] Outdoor Use Only

ORTHO® ORTHO-KLOR® Soil Insect & Termite Killer [ABN: HomeDefense™ Ortho-Klor® Insect & Termite Killer] [9/10/96]; ABN: ORTHO Borer & Leaf Miner Spray Lawn and Nuisance Pests Makes up to 548 Gallons of Spray [1/2 gal size only] Makes up to 274 Gallons of Spray [1 qt. size only] For Termites, Treats 30 Linear Feet of Trench [1/2 gal size only]

Contains: DURSBAN® Soil Insecticide

CONTROLS: Ants, Subterranean Termites, Powder Post Beetles, Crickets, Chinch Bugs, Wood Borers, Carpenter Ants, Grubs, Cutworms, Fleas, Armyworms, Ticks, Mosquitoes, Grasshoppers, European Crane Fly Larvae, Fire Ants, Sod Webworms, Spiders, Box Elder Bugs, Scorpions, and many other soil insects and home invading pests.

[9/16/96]: Concentrate

FOR USE: Around foundations, outside buildings, lawns, woodpiles, stored lumber and fence posts

For Use in Tank Sprayers, Hose-End Applicators, Or Sprinkling Cans Controls All Major Lawn Pests

Kills Lawn Insects

Kills/Controls Ticks that Transmit Lyme Disease

Helps Keep Your Lawn Healthy - Controls Insect Pests that Can Cause Unsightly Brown Spots And Dead Grass

Kills Insects that Damage Ornamentals, Shade Trees, and Fruit Trees

Kills Borers and Leafminers

[9/16/96]: Kills Pine Beetles, Bronze Birch & Peachtree Borers, and many other insects.

Kills Insects: Peachtree Borer, Dogwood Borer, Lilac Borer, Leafminers,

239-2513.WP4 0-995-01 EPA Accepted March 2, 1995

Notifications of 5/28/96, 9/10/96 & 9/16/96; Amendment of 7/17/96

Needleminers, Leaf Beetles, Twig Borer, Pine Beetles, Elm Bark Beetles

Ideal For Use On Ornamentals, Shade Trees, and Fruit Trees [9/16/96]. Use on Shrubs & Trees
Can be Applied as a Foliar Spray

Can be Applied as a Trunk Spray

Can be Applied with Ortho Hose-End Applicators

Active Ingredient	.
Chlorpyrifos* [0,0-diethyl 0-(3,5,6-Trichloro-2-	·
pyridyl)phosphorothioate]	12.6%
Inert Ingredients	87.4%
* This product contains aromatic petroleum solvent	

KEEP OUT OF REACH OF CHILDREN WARNING

See back panel booklet for additional precautionary statements.

NET CONTENTS 64 FL OZ (1/2 GAL) 1.89 L NET CONTENTS 32 FL OZ (1 QT) 946 mL 19/16/96); NET CONTENTS 16 FL OZ (1 PT) 473 mL For prevention and control of ants, subterranean termites and certain wood destroying insects. This product can be used for treatment of localized areas of wooden structures, firewood piles and stored lumber to control wood infesting borers, carpenter ants and carpenter bees. In addition, controls many common pests in lawns and other outdoor areas around the home. [7/17/96]: Specially formulated for residential use.

DO NOT USE INSIDE DWELLINGS

Peel open for use directions and further information

WARNING: Causes eye irritation. Harmful if swallowed. Do not get in eyes, on skin or clothing. Avoid breathing vapors and spray mist. SEE ADDITIONAL PRECAUTIONARY STATEMENTS AND DIRECTIONS INSIDE BOOKLET.

PULL HERE TO OPEN LABEL. PRESS TO RESEAL

Medical Information call 1-800-454-2333
 Product Information call 1-800-225-2883

ORTHO®, ORTHO-KLOR®, DIAL 'n SPRAY®, HomeDefense™ Trademarks of Monsanto Company
DURSBAN® Trademark of DowElanco

©Monsanto Company 1996

Manufactured For
The Solaris Group of
Monsanto Company
P.O. Box 5008
San Ramon CA 94583-0808
Form 07 741-02 [9/16/96]: 0-995-01
Product 5522
EPA Reg. No. 239-2513
EPA Est. 239-IA-3¹, 58996-MO-1^A
Superscript is first letter of lot number

Made in USA

239-2513,WP4 0-995-01 EPA Accepted March 2, 1995

Notifications of 5/28/96, 9/10/96 & 9/16/96; Amendment of 7/17/96

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

tsp = teaspoon Tbs = Tablespoon 4 fl oz = 1/2 cup 2 Tbs equal ∰ 1 fl. oz.

SHAKE WELL BEFORE USING

FOR OUTDOOR HOME & GARDEN USE ONLY

[9/16/96]. An easy to apply product to control borers and leaf miners that attack ornamental trees, fruit trees and shade trees. For use in tank sprayers and hose-end applicators.

ORTHO-KLOR Soil Insect & Termite Killer is an easy to apply product to control lawn insects, home invading pests, fire ants, wood destroying insects including exposed termites, and borers that attack ornamental plants, fruit trees, and shade trees. When used as directed, it helps lawns and gardens stay vigorous, healthy, and green by controlling bugs that cause problems.

[9/16/96]: BORERS & LEAFMINERS ON TREES & SHRUBS

[9/16/96]: Application timing is important. Consult your County Cooperative Extension Agent for proper time to treat.

Pests	Trees Or Shrubs	Product to Measure When To Apply
Reatles (Cottonwood Leaf, Elm Leaf, Flea, Willow Leaf)	Cottonwood, Elm, Willow	4 Lsp./gal. of water. Apply when pests appear.
Weevils (Black Vine, Yellow Poplar, Pine)	Rhedodendron, Poplar, Pine	Thoroughly wet upper and lower leaf surfaces: Reapply at 7 to 10 day intervals if the needed.

239-2513.WP4 0-995-01 EPA Accepted March 2, 1995

Notifications of 5/28/96, 9/10/96 & 9/16/96; Amendment of 7/17/96

Borers (Ash, Dogwood, Lesser Peachtree, Lilac, Oak, Rhododendron, Bronze Birch, Flatheaded Appletree, Locust, Red Oak)	Ash, Lilac, Oak, Dogwood, Rhododendron, Birch, Locust	1-1/2 fl. oz. (3 Tbs.)/gal. of water: Spray trunks and lower limbs of trees and shrubs when adults begin to emerge:
Lezfminers	Elm, Birch, Locust	1-1/2 floz. (3 Tbs.)/ gal. of water: Spray when leafminers appear in early and Mid- May:
Needleminers	Pine, Spruce	1-1/2 fl. oz. (3 Tbs:)/gal. of water.
Beetles (Ambrosia, Black Turpentine, European Elm Bark, Japanese, June, Southern Pine, Spruce)	Elm, Pine Spruce	10-2/3 fl oz (1- 1/4 cups)/gal of water Apply with tank sprayer only. To prevent infestation, spray the tree trunk in early Spring. To treat infested trees, spray the tree trunk before adult beetles begin to emerge. Avoid contact with foliage to prevent leaf damage.

[9/16/96]: SHRUBS

Borers & Leafminers

Cottonwood, Elm, Willow Cottonwood [9/16/96] Leaf Beetles, Elm Leaf Beetles, Floa Beetles, Willow Leaf Beetles; Rhododendron, Poplar, Pine Blackvine Weevils, Yellow Poplar Weevils, Pine Weevils: Use 4 tsp in 1 gal of water. Apply when pests appear. Thoroughly wet upper and lower leaf surfaces. [7/17/96]. ReapplyRetreat at 7 to 10 day intervals if needed.

239-2513.WP4 0-995-01 EPA Accepted March 2, 1995

Notifications of 5(28/96, 9/10/96 & 9/16/96; Amendment of 7/17/96

Ash, Lilac, Oak, Dogwood, Rhododendron, Birch, Locust—Ash Borers, Dogwood Borers, Lesser-Peachtree Borers, Lilac Borers, Oak Borers, Rhododendron Borers, Bronze Birch-Borers, Flatheaded Appletree Borers, Locust Borers, Red Oak Borers—Spray [9/16/96]: trunks and lower limbs of trees and shrubs when adults begin to emerge; Elm, Birch, Locust—Leafminers—Spray in early and mid May when leafminers appear; Pinc, Spruce—Needleminers: Use 1-1/2 fl oz (3 Tbs) in 1 gal of water.

Elm, Pine, Spruce - Ambrosia Beetles, Black Turpentine Beetles, European Elm-Bark Beetles, Japanese Beetles, June Beetles, Southern Pine Beetles, Spruce Beetles: Use 10 2/3 fl oz (1 1/4 cups) in 1 gal of water. Apply only with a tank sprayer. To prevent infestation, spray [9/16/96], the tree trunk in early Spring. To treat infested trees, spray [9/16/96], the tree trunk before adult beetles emerge.

[9/16/96]: Application timing is important. Consult your County Cooperative Extension Agent for proper time to treat.

Pests	Fruit Trees	Product to Measure When to Apply
Dogwood Borer	Apples Wait 1 month after last application before harvesting fruit.	2 fl. oz. (4 Tbs.)/gal. of water. Soak tree trunks and the soil around the base of trees in the spring. Apply up to 8 times per season. Wait 3 weeks between treatments.
Borers (Lesser Peachtree, Great Peachtree, American Plum)	Sweet Cherries Wait 1 week after application before harvesting fruit.	2 fl. oz. (4 Tbs.)/gal. of water. Spray trunks and lower limbs. Avoid contact with foliage to prevent leaf damage. Apply up to 3 times per season. Apply second application 2 weeks after the first. Make third application after harvest.

Pests	Fruit Trees	Product to Measure When to Apply
Borers (American Plum, Lesser Peachtree, Pacific	Sour Cherries	2 fl. oz. (4 Tbs.)/gal. of water.
Flatheaded, Peach Twig, Peachtree, Shothole)	Wait 2 weeks after application before harvesting fruit.	Spray tree trunks and lower limbs: Apply up to 8 times per season.
Peachtree Borers	Nectarines, Peaches Wait 2 weeks after application before harvesting fruit.	4 fl. oz. (8 Tbs.)/gal. of water. Spray bark from the ground up to the lower branches. Do not spray fruit. Make only 1 application per year.

[9/16/96]: FRUIT TREES

Apples Dogwood Borer: Use 2 fl oz (4 Tbs) in 1 gal of water. Starting in the Spring, spray 9/16/961 soak tree trunks and soil around the base of trees. Apply up to 8 times per season. Wait 3 weeks before [7/17/96]: reapplying retreating. Wait 1 month after last spraying before harvesting fruit.

Sweet Cherries - Lesser Peachtree Borers, Great Peachtree Borers, American Plum Borers: Use 2 fl oz (4 Tbs) in 1-gal of water. Spray trunks and lower limbs. To prevent leaf-damage, avoid foliage contact. Apply up to 3 times per season. Wait 2 weeks before [7/17/96]: reapplying retreating. Wait 1 week after spraying [9/16/96]: application before harvesting fruit. Make 3rd application after harvest.

Sour Cherries — American Plum Borers, Lesser Peachtree Borers, Pacific Flathcaded Borers, Peachtree Borers, Shothole Borers. Use 2 fl oz (4 Tbs) in 1-gal of water. Spray [9/16/96]: tree trunks and lower limbs. Apply up to 3 times per season. Wait 2 weeks after spraying [9/16/96]: application before harvesting fruit. Consult your Extension Service Specialist for the best time to treat.

Nectarines, Peaches Peachtree Borers: Use 4 fl-oz (8 Tbs) in 1 gal of water.

Application timing is important. Consult your Extension Service Specialist for proper time to treat. Spray bark from the ground up to the lower branches. Do not spray fruit. Make only 1 application per year. Wait 2 weeks after spraying

239-2513.WP4

[9/16/96]: application before harvesting fruit.

LAWN PESTS: Armyworm, Lawn Moth Larvae (Sod Webworms), Dichondra Flea Beetles, Chinchbugs, Cutworms, European Crane Fly (Larvae), Fiery Skipper Larvae (California and Hawaii only), Tropical Grass Webworms (Hawaii only), Grasshoppers, Crickets, Chiggers, Clover Mites, Earwigs, Fleas-For use in an ORTHO DIAL 'n SPRAY® Hose-End Sprayer: Add 3-1/2 fl oz undiluted product to sprayer jar, set dial on sprayer at 2 tsp/gallon and spray until jar is empty to treat 1,000 sq ft. For use with an ORTHO Lawn Sprayer: Add 1-3/4 fl oz product to sprayer jar, fill jar with water to the 15 gallon mark and spray evenly over 500 sq ft.

Mow and water 1 to 2 days before treatment. Heavy watering should be withheld for 2 to 3 days following treatment. Spray when pests first appear. Treat entire lawn. Do not attempt to control lawn pests by spot spraying. Repeat treatments may be necessary at 4 to 6 week intervals. For European Crane Fly control, consult your Agricultural Extension Service for proper timing of application for your area.

WHITE GRUBS IN LAWNS: (Larvae of European Chafer and Japanese Beetle) - For use in an ORTHO DIAL 'n SPRAY® Hose-End Sprayer: Add 13 fl oz undiluted product to sprayer jar, set dial on sprayer at 1 oz per gallon and spray until jar is empty to treat 1,000 sq ft. Water lawn thoroughly after application. For use with an ORTHO Lawn Sprayer: Add 6-1/2 fl oz product to sprayer jar. Fill jar with water to the 15 gallon mark and spray evenly over 500 sq ft. Treat entire lawn. Do not attempt to control lawn pests by spot spraying. Water lawn thoroughly to wash the insecticide into the soil. Treat lawn when grubs are young and actively feeding near the soil surface, usually during late July and August. Consult your Agricultural Extension Service Specialist for proper time to treat in your area. Mosquitoes (outdoors): Mosquitoes coming to rest on areas treated for the control of lawn and home invading pests will be killed for varying periods of time after treatment, depending on exposure of treated areas to weathering conditions. Note: Do not use on vegetables or food crops. Do not use on azaleas, camellias, poinsettias, rose bushes or variegated ivy because of possible injury to the plants. [Termite Trench Treatment: the following information in italics will not be printed on our consumer product labels at the time of next printing. We do plan to retain the registered usel

FOR TERMITES (Outdoors): TRENCH TREATMENT: NOTE: This product is not intended for control of termites which do not require contact with the soil for moisture, such as: drywood, dampwood or powderpost termites. Consult the County or State Agricultural Extension Service for termite identification. Remove all wood debris and wood forms. Determine the extent of termite infestation. For prevention and light infestation, treat soil adjacent to both sides of infested foundations of poured concrete, pipes or piers by digging a trench 6 inches wide

and 4 inches deep (or down to the top of the footing whichever is less). For brick and block foundations dig the trench about 12 inches deep. If footing is more than 12 inches deep, make holes with a crowbar, pipe or a rod about one foot apart that extend from the trench bottom to the top of the footing. Make holes closer in hard-packed clay soils. NEVER DIG BELOW THE TOP OF THE FOOTING. Avoid contamination of public and private water supplies by following these precautions. Do not allow a hose or any type of faucet extension to reach into the termite chemical solution while filling or mixing this material with water. Use anti-backflow or anti-siphonage equipment on all filling equipment. These devices can be purchased at hardware or plumbing supply stores. Use of anti-backflow equipment will help insure that the user of this toxic chemical will not contaminate domestic water supplies. Refer to FHA (Federal Housing Administration), state and local specifications for safe distances of treatment areas from wells. Soil in the vicinity of wells, cisterns or ponds should not be treated. Do not treat structures that contain cisterns or wells. Note: When conditions will not permit trenching, surface application adjacent to interior foundation walls may be made but the treated strip should not exceed 18" in width. Apply solution with a sprinkling can. Mix 5-1/3 fl. oz. per gallon of water and treat 2-1/2 linear feet of trench or foundation (one half gallon of ORTHO-KLOR Soil Insect & Termite Killer will make 12 gallons of diluted solution and treat 30 linear feet). Mix solution into soil as it is replaced in the trench. Cover treated soil with a layer of untreated soil. Be sure to treat any accessible voids in hollow blocks in the foundations at same rate as above. Be careful to have adequate ventilation when treating areas under structure. This product is not intended for severe infestations or treatment under slabs, in basements and buildings with insufficient crawl space, etc., contact a licensed pest control operator or termite control service.

Retreatment for subterranean termites should only be made when there is evidence of reinfestation subsequent to the initial treatment, or there has been a disruption of the chemical barrier in the soil due to construction, excavation, landscaping, etc. Reapplication should be made as a spot treatment to those areas.

Annual retreatment of the entire premises must be avoided.

FENCE POSTS-TERMITE PREVENTION (EXCEPT RHODE ISLAND) [5/28/96]: Apply solution with a sprinkling can. Mix 5-1/3 fl oz per gal of water. Apply one gallon of diluted product per post hole. Pour one half gallon of solution into hole and add the remaining 1/2 gallon to the soil when filling the hole so that all the fill soil is treated to form a protective barrier.

NOTE: Do not treat structures that contain cisterns or wells. Do not treat soil that is water saturated or frozen. Consult state and local specifications for recommended distances of treatment from well.

TREATMENT FOR WOOD DESTROYING PESTS (Outdoors): (Powder Post Beetles, Furniture Beetles, Deathwatch Beetles, Round-headed House Borers, Carpenter Ants, Carpenter Bees). Treat localized areas of infested wood. Mix 2½ Tbs per

239-2513.WP4 0-995-01

quart of water. Apply evenly on wood surfaces with a paint brush. Allow to dry and repeat to ensure penetration.

For wood piles and stored lumber apply 5-1/3 fl oz per gal of water. Use ORTHO hose-end sprayers or sprinkling can to deliver a coarse drenching spray. Do not use in structures occupied by animals to be used for food purposes or which produce products for human consumption.

HOME INVADING PESTS (OUTSIDE DWELLINGS): Ants, Bees, Beetles (1), Box Elder Bugs, Carpenter Ants (1), Carpenter Bees (1), Centipedes, Clover Mites, Cockroaches, Crickets, Earwigs, Elm Leaf Beetles, Fleas, Flies, Hornets, House Flies, Millipedes, Mosquitoes, Scorpions (2), Silverfish, Sowbugs, Spiders, Springtails, Ticks, Wasps, Yellowjackets.

NOTE: Number in parenthesis refer to Specific Directions. (1) For added information, see section on Treatment For Wood Destroying Pests. (2) For Scorpions, remove accumulations of lumber, firewood, and other materials which serve as harborage sites and treat beneath and adjacent to infested areas. FOR PEST CONTROL ON OUTSIDE SURFACES AND AROUND BUILDINGS: Spray outside surfaces of buildings and surrounding areas, including porches, window frames, doorways, foundations, eaves, patios, garages, garbage cans, and other areas where pests congregate or can gain entrance. Repeat treatment as needed. Do not treat pets with this product. DO NOT USE IN DWELLINGS.

Treat a band of soil 6 feet wide around buildings, also the building foundation to a height of 2 to 3 feet Spray 10 gal mixture per 1,000 sq ft.

AMOUNT TO USE FOR HOME INVADING PESTS (OUTDOORS): For use in Hose-end sprayer, tank-type sprayer or sprinkling can: Apply 2 tsp in one gallon of water to cover 100 sq ft. For use in an ORTHO DIAL 'n SPRAY: Add 3-1/2 fl oz. undiluted product to sprayer jar, set dial on sprayer at 2 tsp /gal and spray until jar is empty to treat 1,000 sq ft.

IMPORTED FIRE ANT MOUNDS: Mix 2 Tbs per gal of water to each mound, using a sprinkler can. Thoroughly wet mound and treat a 4 foot diameter circle around the mound. For best results, apply in cool weather or in early morning or late afternoon. Treat new mounds as they appear.

STORAGE AND DISPOSAL

STORAGE: Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Avoid contamination of feed, foodstuffs and domestic water supplies. Store in a cool, dry place, preferably in a locked storage area. Do not store diluted spray. Store above freezing. Handle concentrate in a ventilated area. Keep container closed. DISPOSAL: PRODUCT-Partially filled bottle may be disposed of by securely wrapping original container in several layers of newspaper and discard in trash. CONTAINER-Do not reuse empty bottle. Rinse thoroughly before discarding in trash.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

WARNING: Causes eye irritation. Harmful if swallowed. Do not get in eyes, on skin or clothing. Avoid breathing vapors and spray mist. Handle concentrate in ventilated area. Wash thoroughly after handling and before eating or smoking. Do not wear contaminated clothing or shoes. Keep away from food, feedstuffs and domestic water supplies. [CHILD/ PET ICON] Keep children and pets out of treated areas until these areas are dry.

FIRST AID: IF SWALLOWED: DO NOT make person vomit. Call a doctor immediately. IF ON SKIN: In case of contact, remove contaminated clothing and immediately wash skin with soap and water. IF IN EYES: Flush eyes with plenty of water. Call a physician and get prompt medical attention.

Note to Physicians: Emergency Information call 1-800-454-2333. This product contains an aromatic petroleum solvent and may present an aspiration problem. Chlorpyrifos is a cholinesterase inhibitor. If signs and symptoms of cholinesterase inhibition develop, atropine by injection is antidotal; 2-PAM may also be given in conjunction with atropine, but should never be used alone.

ENVIRONMENTAL HAZARDS: This product is toxic to fish, birds and other wildlife. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

PHYSICAL OR CHEMICAL HAZARDS: Do not use or store near heat or open flame.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

Medical Information call 1-800-454-2333
 Product Information call 1-800-225-2883

ORTHO®,ORTHO-KLOR®,DIAL 'n SPRAY®, HomeDefense™ Trademarks of Monsanto Company
DURSBAN® Trademark of DowElanco

[©]Monsanto Company 1996

Manufactured For
The Solaris Group of
Monsanto Company
P.O. Box 5008
San Ramon CA 94583-0808
Form 07 741-02 [9/16/96]: 0-995-01
Product 5522
EPA Reg. No. 239-2513

239-2513,WP4 0-995-01

EPA Est. 239-IA-3¹, 58996-MO-1^A
Superscript is first letter of lot number
Made in USA

Page 17 15

Please read instructions	on reverse before completing form.		⊬ ···· · For	m Approved.	OMB No. 207	0-0060.	Approval expires 05-31-98
\$EPA	Environmental Protection Washington, DC 204	on Agei	1,,4 (*) et 1 CY		Registrat Amendm Other		OPP Identifier Number
	Application	on for F	esticide ·	- Section			
1. Company/Product Nur 239-2513	nber	-=	2. EPA Prode Denni:	ıct Manager s Edwards, J	r.	1	posed Classification
4. Company/Product (Na 0KTH0-KLC	me) R Soil Insect & Termite Killer		PM#	19			None Restricted
SOLARIS O P. O. Box San Ramor	Applicant (Include ZIP Code) Froup of Monsanto Company 5006, 1, CA 94583-0906 this is a new address	Saci	(b)(i), my pi to:	No	ilar or identic		FIFRA Section 3(c)(3) mposition and labeling
		000	<u></u>				
Amendment - Exp	response to Agency letter dated		_ Age	ency letter dat a Too" Applica	ls in response ted NOTIFIC		<u> </u>
Notification - Exp	lain below.	-	Oth	er - Explain be	alow.307 4	įąci:	
every rec	er Spray reflecting these changes. puirement of PR Notice 95-2.	· .	0-995-01. 1 11	is is to cer	tify that th	is notif	ication meets
1. Material This Product				· · · · · · · · · · · · · · · · · · ·		<u>.</u>	1
Child-Resistant Packagin Yes* No * Certification must be submitted 3. Location of Net Conte	Yes No If "Yes" Unit Packaging wgt. No. per container.	If "Yes Packag	e wgt (No. per container	2. Type of C	Metal Plastic Glass Paper Other (S	The section of the se
Label	च्छित पुराविकार्यक्ष हात्रेज पूर्वकर विकास है है। असे Container अस्त्रेली के अस्त्रेल कर				On Label	. : '	panying product
6. Manner in Which Labo	el is Affixed to Product Lithon Paper Stenc	graph r glued ciled	· · · · · · [Other			Frague (CAT) (CAT)
	'		ion - IV	* .	1		12
1. Contact Point (Comp	lete items directly below for identificati	on of indivi	dual to be co	ntacted, if nec	essary, to pro	cess this	application.)
Name F. X. Kai	uenski	Title Re	gistration S	Specialist		Celephon	No (ligotude Area Code)
	Certific tatements I have made on this form and at any knowingly false or misleading sta able law.	d all attach					6: Dats Application Received (Stamped)
2. Signature 3. Title							CAUCITY CAUCITY
trances	X. Lamerske	Re	gistration S	Specialist		<u>-</u>	
4. Typed Name	X. KAMIEUSKI	5. Date	9/1	6/96			inger de de la companya de la compa



ITF CODE IS 910801

STORAGE AND DISPOSAL STORAGE. Store in a secure, cool dry place. Keep from freezing, DISPOSAL: Do not reuse empty container. Rinse thoroughly, wrap in newspaper and put in trash.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

WARNING: Causes substantial but temporary eye injury. Harmful if swallowed or if on skin, Do not get in eyes or on clothing. Wear goggles or safety glasses. Avoid contact with skin, Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

People and pets may re-enter treated areas after sprayed area has dried.

area has died.

FIRST AID: If IN EYES: Hold eyelids open and flush with a steady, gentle stream of water. Get medical attention, IF SWALLOWED, if the person is alert and able to swallow, give small sips of milk or water. Call a physician or poison control center for further treatment advice. IF ON SKIN, Wash with plenty of soap and water. Get medical attention if intration persists. Note to Physicians: Emergency Information call 1-800-484-2333. Chloropyrifos is a cholmesterase inhibitor. Atropine is antidottal, 2-PAM is artidottal and may be used in conjunction with atropine, but should not be used alone.

ENVIRONMENTAL, HAZARDS. This product is took to fish and whickle, Do not apply directly to water or contaminate water when disposing of ecouprient washwater or disposal of waste. This product is highly took to bees exposed to direct treatment area.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.



Medical Information call 1-800-454-2333 Product Information call 1-800-225-2883

Manufactured for The SOLARIS Group of Monsanto Company P.O. Box 5008 San Ramon CA 94583-0808 Form 9-108-01 / Product 6021 EPA Reg. No. 239-2635 EPA Est. 239-14-3', 58996-MO-14 Superscript is first letter of lot number

Made in USA

[ICON] Outdoor Use Only

ORTHO® ORTHO-KLOR® Soil Insect & Termite Killer [ABN: HomeDefense™ Ortho-Klor® Insect & Termite Killer] [9/10/96]; ABN: ORTHO Borer & Leaf Miner Spray Lawn and Nuisance Pests Makes up to 548 Gallons of Spray [1/2 gal size only] Makes up to 274 Gallons of Spray [1 qt. size only] For Termites, Treats 30 Linear Feet of Trench [1/2 gal size only]

Contains: DURSBAN® Soil Insecticide

CONTROLS: Ants, Subterranean Termites, Powder Post Beetles, Crickets, Chinch Bugs, Wood Borers, Carpenter Ants, Grubs, Cutworms, Fleas, Armyworms, Ticks, Mosquitoes, Grasshoppers, European Crane Fly Larvae, Fire Ants, Sod Webworms, Spiders, Box Elder Bugs, Scorpions, and many other soil insects and home invading pests.

[9/16/96]: Concentrate

FOR USE: Around foundations, outside buildings, lawns, woodpiles, stored lumber and fence posts

For Use in Tank Sprayers, Hose-End Applicators, Or Sprinkling Cans Controls All Major Lawn Pests

Kills Lawn Insects

Kills/Controls Ticks that Transmit Lyme Disease

Helps Keep Your Lawn Healthy - Controls Insect Pests that Can Cause Unsightly Brown Spots And Dead Grass

Kills Insects that Damage Ornamentals, Shade Trees, and Fruit Trees

Kills Borers and Leafminers

[9/16/96]: Kills Pine Beetles, Bronze Birch & Peachtree Borers, and many other insects.

Kills Insects: Peachtree Borer, Dogwood Borer, Lilac Borer, Leafminers,

239-2513.WP4 0-995-01 EPA Accepted March 2, 1995

Notifications of 5/28/96, 9/10/96 & 9/16/96; Amendment of 7/17/96

Needleminers, Leaf Beetles, Twig Borer, Pine Beetles, Elm Bark Beetles

Ideal For Use On Ornamentals, Shade Trees, and Fruit Trees [9/16/96]. Use on Shrubs & Trees
Can be Applied as a Foliar Spray

Can be Applied as a Trunk Spray

Can be Applied with Ortho Hose-End Applicators

Active Ingredient	.
Chlorpyrifos* [0,0-diethyl 0-(3,5,6-Trichloro-2-	·
pyridyl)phosphorothioate]	12.6%
Inert Ingredients	87.4%
* This product contains aromatic petroleum solvent	

KEEP OUT OF REACH OF CHILDREN WARNING

See back panel booklet for additional precautionary statements.

NET CONTENTS 64 FL OZ (1/2 GAL) 1.89 L NET CONTENTS 32 FL OZ (1 QT) 946 mL 19/16/96); NET CONTENTS 16 FL OZ (1 PT) 473 mL For prevention and control of ants, subterranean termites and certain wood destroying insects. This product can be used for treatment of localized areas of wooden structures, firewood piles and stored lumber to control wood infesting borers, carpenter ants and carpenter bees. In addition, controls many common pests in lawns and other outdoor areas around the home. [7/17/96]: Specially formulated for residential use.

DO NOT USE INSIDE DWELLINGS

Peel open for use directions and further information

WARNING: Causes eye irritation. Harmful if swallowed. Do not get in eyes, on skin or clothing. Avoid breathing vapors and spray mist. SEE ADDITIONAL PRECAUTIONARY STATEMENTS AND DIRECTIONS INSIDE BOOKLET.

PULL HERE TO OPEN LABEL. PRESS TO RESEAL

Medical Information call 1-800-454-2333
 Product Information call 1-800-225-2883

ORTHO®,ORTHO-KLOR®,DIAL 'n SPRAY®, HomeDefense™ Trademarks of Monsanto Company
DURSBAN® Trademark of DowElanco

©Monsanto Company 1996

Manufactured For
The Solaris Group of
Monsanto Company
P.O. Box 5008
San Ramon CA 94583-0808
Form 07 741-02 [9/16/96]: 0-995-01
Product 5522
EPA Reg. No. 239-2513
EPA Est. 239-IA-3¹, 58996-MO-1^A
Superscript is first letter of lot number

Made in USA

239-2513,WP4 0-995-01 EPA Accepted March 2, 1995

Notifications of 5/28/96, 9/10/96 & 9/16/96; Amendment of 7/17/96

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

tsp = teaspoon Tbs = Tablespoon 4 fl oz = 1/2 cup 2 Tbs equal ∰ 1 fl. oz.

SHAKE WELL BEFORE USING

FOR OUTDOOR HOME & GARDEN USE ONLY

[9/16/96]: An easy to apply product to control borers and leaf miners that attack ornamental trees, fruit trees and shade trees. For use in tank sprayers and hose-end applicators:

ORTHO-KLOR Soil Insect & Termite Killer is an easy to apply product to control lawn insects, home invading pests, fire ants, wood destroying insects including exposed termites, and borers that attack ornamental plants, fruit trees, and shade trees. When used as directed, it helps lawns and gardens stay vigorous, healthy, and green by controlling bugs that cause problems.

[9/16/96]: BORERS & LEAFMINERS ON TREES & SHRUBS

[9/16/96]: Application timing is important. Consult your County Cooperative Extension Agent for proper time to treat.

Pests	Trees Or Shrubs	Product to Measure When To Apply
Reatles (Cottonwood Leaf, Elm Leaf, Flea, Willow Leaf)	Cottonwood, Elm, Willow	4 Lsp./gal. of water. Apply when pests appear.
Weevils (Black Vine, Yellow Poplar, Pine)	Rhedodendron, Poplar, Pine	Thoroughly wet upper and lower leaf surfaces: Reapply at 7 to 10 day intervals if the needed.

239-2513.WP4 0-995-01

Borers (Ash, Dogwood, Lesser Peachtree, Lilac, Oak, Rhododendron, Bronze Birch, Flatheaded Appletree, Locust, Red Oak)	Ash, Lilac, Oak, Dogwood, Rhododendron, Birch, Locust	1-1/2 fl. oz. (3 Tbs.)/gal. of water: Spray trunks and lower limbs of trees and shrubs when adults begin to emerge:
Lezfminers	Elm, Birch, Locust	1-1/2 floz. (3 Tbs.)/ gal. of water: Spray when leafminers appear in early and Mid- May:
Needleminers	Pine, Spruce	1-1/2 fl. oz. (3 Tbs:)/gal. of water.
Beetles (Ambrosia, Black Turpentine, European Elm Bark, Japanese, June, Southern Pine, Spruce)	Elm, Pine Spruce	10-2/3 fl oz (1- 1/4 cups)/gal of water Apply with tank sprayer only. To prevent infestation, spray the tree trunk in early Spring. To treat infested trees, spray the tree trunk before adult beetles begin to emerge. Avoid contact with foliage to prevent leaf damage.

[9/16/96]: SHRUBS

Borers & Leafminers

Cottonwood, Elm, Willow Cottonwood [9/16/96] Leaf Beetles, Elm Leaf Beetles, Floa Beetles, Willow Leaf Beetles; Rhododendron, Poplar, Pine Blackvine Weevils, Yellow Poplar Weevils, Pine Weevils: Use 4 tsp in 1 gal of water. Apply when pests appear. Thoroughly wet upper and lower leaf surfaces. [7/17/96]. ReapplyRetreat at 7 to 10 day intervals if needed.

239-2513.WP4 0-995-01 EPA Accepted March 2, 1995

Notifications of 5(28/96, 9/10/96 & 9/16/96; Amendment of 7/17/96

Ash, Lilac, Oak, Dogwood, Rhododendron, Birch, Locust—Ash Borers, Dogwood Borers, Lesser-Peachtree Borers, Lilac Borers, Oak Borers, Rhododendron Borers, Bronze Birch-Borers, Flatheaded Appletree Borers, Locust Borers, Red Oak Borers—Spray [9/16/96]: trunks and lower limbs of trees and shrubs when adults begin to emerge; Elm, Birch, Locust—Leafminers—Spray in early and mid May when leafminers appear; Pinc, Spruce—Needleminers: Use 1-1/2 fl oz (3 Tbs) in 1 gal of water.

Elm, Pine, Spruce - Ambrosia Beetles, Black Turpentine Beetles, European Elm-Bark Beetles, Japanese Beetles, June Beetles, Southern Pine Beetles, Spruce Beetles: Use 10 2/3 fl oz (1 1/4 cups) in 1 gal of water. Apply only with a tank sprayer. To prevent infestation, spray [9/16/96], the tree trunk in early Spring. To treat infested trees, spray [9/16/96], the tree trunk before adult beetles emerge.

[9/16/96]: Application timing is important. Consult your County Cooperative Extension Agent for proper time to treat.

Pests	Fruit Trees	Product to Measure When to Apply
Dogwood Borer	Apples Wait 1 month after last application before harvesting fruit.	2 fl. oz. (4 Tbs.)/gal. of water. Soak tree trunks and the soil around the base of trees in the spring. Apply up to 8 times per season. Wait 3 weeks between treatments.
Borers (Lesser Peachtree, Great Peachtree, American Plum)	Sweet Cherries Wait 1 week after application before harvesting fruit.	2 fl. oz. (4 Tbs.)/gal. of water. Spray trunks and lower limbs. Avoid contact with foliage to prevent leaf damage. Apply up to 3 times per season. Apply second application 2 weeks after the first. Make third application after harvest.

Pests	Fruit Trees	Product to Measure When to Apply
Borers (American Plum, Lesser Peachtree, Pacific	Sour Cherries	2 fl. oz. (4 Tbs.)/gal. of water.
Flatheaded, Peach Twig, Peachtree, Shothole)	Wait 2 weeks after application before harvesting fruit.	Spray tree trunks and lower limbs: Apply up to 8 times per season.
Peachtree Borers	Nectarines, Peaches Wait 2 weeks after application before harvesting fruit.	4 fl. oz. (8 Tbs.)/gal. of water. Spray bark from the ground up to the lower branches. Do not spray fruit. Make only 1 application per year.

[9/16/96]: FRUIT TREES

Apples Dogwood Borer: Use 2 fl oz (4 Tbs) in 1 gal of water. Starting in the Spring, spray 9/16/961 soak tree trunks and soil around the base of trees. Apply up to 8 times per season. Wait 3 weeks before [7/17/96]: reapplying retreating. Wait 1 month after last spraying before harvesting fruit.

Sweet Cherries - Lesser Peachtree Borers, Great Peachtree Borers, American Plum Borers: Use 2 fl oz (4 Tbs) in 1-gal of water. Spray trunks and lower limbs. To prevent leaf-damage, avoid foliage contact. Apply up to 3 times per season. Wait 2 weeks before [7/17/96]: reapplying retreating. Wait 1 week after spraying [9/16/96]: application before harvesting fruit. Make 3rd application after harvest.

Sour Cherries — American Plum Borers, Lesser Peachtree Borers, Pacific Flathcaded Borers, Peachtree Borers, Shothole Borers. Use 2 fl oz (4 Tbs) in 1-gal of water. Spray [9/16/96]: tree trunks and lower limbs. Apply up to 3 times per season. Wait 2 weeks after spraying [9/16/96]: application before harvesting fruit. Consult your Extension Service Specialist for the best time to treat.

Nectarines, Peaches Peachtree Borers: Use 4 fl-oz (8 Tbs) in 1 gal of water.

Application timing is important. Consult your Extension Service Specialist for proper time to treat. Spray bark from the ground up to the lower branches. Do not spray fruit. Make only 1 application per year. Wait 2 weeks after spraying

239-2513.WP4

[9/16/96]: application before harvesting fruit.

LAWN PESTS: Armyworm, Lawn Moth Larvae (Sod Webworms), Dichondra Flea Beetles, Chinchbugs, Cutworms, European Crane Fly (Larvae), Fiery Skipper Larvae (California and Hawaii only), Tropical Grass Webworms (Hawaii only), Grasshoppers, Crickets, Chiggers, Clover Mites, Earwigs, Fleas-For use in an ORTHO DIAL 'n SPRAY® Hose-End Sprayer: Add 3-1/2 fl oz undiluted product to sprayer jar, set dial on sprayer at 2 tsp/gallon and spray until jar is empty to treat 1,000 sq ft. For use with an ORTHO Lawn Sprayer: Add 1-3/4 fl oz product to sprayer jar, fill jar with water to the 15 gallon mark and spray evenly over 500 sq ft.

Mow and water 1 to 2 days before treatment. Heavy watering should be withheld for 2 to 3 days following treatment. Spray when pests first appear. Treat entire lawn. Do not attempt to control lawn pests by spot spraying. Repeat treatments may be necessary at 4 to 6 week intervals. For European Crane Fly control, consult your Agricultural Extension Service for proper timing of application for your area.

WHITE GRUBS IN LAWNS: (Larvae of European Chafer and Japanese Beetle) - For use in an ORTHO DIAL 'n SPRAY® Hose-End Sprayer: Add 13 fl oz undiluted product to sprayer jar, set dial on sprayer at 1 oz per gallon and spray until jar is empty to treat 1,000 sq ft. Water lawn thoroughly after application. For use with an ORTHO Lawn Sprayer: Add 6-1/2 fl oz product to sprayer jar. Fill jar with water to the 15 gallon mark and spray evenly over 500 sq ft. Treat entire lawn. Do not attempt to control lawn pests by spot spraying. Water lawn thoroughly to wash the insecticide into the soil. Treat lawn when grubs are young and actively feeding near the soil surface, usually during late July and August. Consult your Agricultural Extension Service Specialist for proper time to treat in your area. Mosquitoes (outdoors): Mosquitoes coming to rest on areas treated for the control of lawn and home invading pests will be killed for varying periods of time after treatment, depending on exposure of treated areas to weathering conditions. Note: Do not use on vegetables or food crops. Do not use on azaleas, camellias, poinsettias, rose bushes or variegated ivy because of possible injury to the plants. [Termite Trench Treatment: the following information in italics will not be printed on our consumer product labels at the time of next printing. We do plan to retain the registered usel

FOR TERMITES (Outdoors): TRENCH TREATMENT: NOTE: This product is not intended for control of termites which do not require contact with the soil for moisture, such as: drywood, dampwood or powderpost termites. Consult the County or State Agricultural Extension Service for termite identification. Remove all wood debris and wood forms. Determine the extent of termite infestation. For prevention and light infestation, treat soil adjacent to both sides of infested foundations of poured concrete, pipes or piers by digging a trench 6 inches wide

and 4 inches deep (or down to the top of the footing whichever is less). For brick and block foundations dig the trench about 12 inches deep. If footing is more than 12 inches deep, make holes with a crowbar, pipe or a rod about one foot apart that extend from the trench bottom to the top of the footing. Make holes closer in hard-packed clay soils. NEVER DIG BELOW THE TOP OF THE FOOTING. Avoid contamination of public and private water supplies by following these precautions. Do not allow a hose or any type of faucet extension to reach into the termite chemical solution while filling or mixing this material with water. Use anti-backflow or anti-siphonage equipment on all filling equipment. These devices can be purchased at hardware or plumbing supply stores. Use of anti-backflow equipment will help insure that the user of this toxic chemical will not contaminate domestic water supplies. Refer to FHA (Federal Housing Administration), state and local specifications for safe distances of treatment areas from wells. Soil in the vicinity of wells, cisterns or ponds should not be treated. Do not treat structures that contain cisterns or wells. Note: When conditions will not permit trenching, surface application adjacent to interior foundation walls may be made but the treated strip should not exceed 18" in width. Apply solution with a sprinkling can. Mix 5-1/3 fl. oz. per gallon of water and treat 2-1/2 linear feet of trench or foundation (one half gallon of ORTHO-KLOR Soil Insect & Termite Killer will make 12 gallons of diluted solution and treat 30 linear feet). Mix solution into soil as it is replaced in the trench. Cover treated soil with a layer of untreated soil. Be sure to treat any accessible voids in hollow blocks in the foundations at same rate as above. Be careful to have adequate ventilation when treating areas under structure. This product is not intended for severe infestations or treatment under slabs, in basements and buildings with insufficient crawl space, etc., contact a licensed pest control operator or termite control service.

Retreatment for subterranean termites should only be made when there is evidence of reinfestation subsequent to the initial treatment, or there has been a disruption of the chemical barrier in the soil due to construction, excavation, landscaping, etc. Reapplication should be made as a spot treatment to those areas.

Annual retreatment of the entire premises must be avoided.

FENCE POSTS-TERMITE PREVENTION (EXCEPT RHODE ISLAND) [5/28/96]: Apply solution with a sprinkling can. Mix 5-1/3 fl oz per gal of water. Apply one gallon of diluted product per post hole. Pour one half gallon of solution into hole and add the remaining 1/2 gallon to the soil when filling the hole so that all the fill soil is treated to form a protective barrier.

NOTE: Do not treat structures that contain cisterns or wells. Do not treat soil that is water saturated or frozen. Consult state and local specifications for recommended distances of treatment from well.

TREATMENT FOR WOOD DESTROYING PESTS (Outdoors): (Powder Post Beetles, Furniture Beetles, Deathwatch Beetles, Round-headed House Borers, Carpenter Ants, Carpenter Bees). Treat localized areas of infested wood. Mix 2½ Tbs per

239-2513.WP4 0-995-01

quart of water. Apply evenly on wood surfaces with a paint brush. Allow to dry and repeat to ensure penetration.

For wood piles and stored lumber apply 5-1/3 fl oz per gal of water. Use ORTHO hose-end sprayers or sprinkling can to deliver a coarse drenching spray. Do not use in structures occupied by animals to be used for food purposes or which produce products for human consumption.

HOME INVADING PESTS (OUTSIDE DWELLINGS): Ants, Bees, Beetles (1), Box Elder Bugs, Carpenter Ants (1), Carpenter Bees (1), Centipedes, Clover Mites, Cockroaches, Crickets, Earwigs, Elm Leaf Beetles, Fleas, Flies, Hornets, House Flies, Millipedes, Mosquitoes, Scorpions (2), Silverfish, Sowbugs, Spiders, Springtails, Ticks, Wasps, Yellowjackets.

NOTE: Number in parenthesis refer to Specific Directions. (1) For added information, see section on Treatment For Wood Destroying Pests. (2) For Scorpions, remove accumulations of lumber, firewood, and other materials which serve as harborage sites and treat beneath and adjacent to infested areas. FOR PEST CONTROL ON OUTSIDE SURFACES AND AROUND BUILDINGS: Spray outside surfaces of buildings and surrounding areas, including porches, window frames, doorways, foundations, eaves, patios, garages, garbage cans, and other areas where pests congregate or can gain entrance. Repeat treatment as needed. Do not treat pets with this product. DO NOT USE IN DWELLINGS.

Treat a band of soil 6 feet wide around buildings, also the building foundation to a height of 2 to 3 feet Spray 10 gal mixture per 1,000 sq ft.

AMOUNT TO USE FOR HOME INVADING PESTS (OUTDOORS): For use in Hose-end sprayer, tank-type sprayer or sprinkling can: Apply 2 tsp in one gallon of water to cover 100 sq ft. For use in an ORTHO DIAL 'n SPRAY: Add 3-1/2 fl oz. undiluted product to sprayer jar, set dial on sprayer at 2 tsp /gal and spray until jar is empty to treat 1,000 sq ft.

IMPORTED FIRE ANT MOUNDS: Mix 2 Tbs per gal of water to each mound, using a sprinkler can. Thoroughly wet mound and treat a 4 foot diameter circle around the mound. For best results, apply in cool weather or in early morning or late afternoon. Treat new mounds as they appear.

STORAGE AND DISPOSAL

STORAGE: Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Avoid contamination of feed, foodstuffs and domestic water supplies. Store in a cool, dry place, preferably in a locked storage area. Do not store diluted spray. Store above freezing. Handle concentrate in a ventilated area. Keep container closed. DISPOSAL: PRODUCT-Partially filled bottle may be disposed of by securely wrapping original container in several layers of newspaper and discard in trash. CONTAINER-Do not reuse empty bottle. Rinse thoroughly before discarding in trash.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

WARNING: Causes eye irritation. Harmful if swallowed. Do not get in eyes, on skin or clothing. Avoid breathing vapors and spray mist. Handle concentrate in ventilated area. Wash thoroughly after handling and before eating or smoking. Do not wear contaminated clothing or shoes. Keep away from food, feedstuffs and domestic water supplies. [CHILD/ PET ICON] Keep children and pets out of treated areas until these areas are dry.

FIRST AID: IF SWALLOWED: DO NOT make person vomit. Call a doctor immediately. IF ON SKIN: In case of contact, remove contaminated clothing and immediately wash skin with soap and water. IF IN EYES: Flush eyes with plenty of water. Call a physician and get prompt medical attention.

Note to Physicians: Emergency Information call 1-800-454-2333. This product contains an aromatic petroleum solvent and may present an aspiration problem. Chlorpyrifos is a cholinesterase inhibitor. If signs and symptoms of cholinesterase inhibition develop, atropine by injection is antidotal; 2-PAM may also be given in conjunction with atropine, but should never be used alone.

ENVIRONMENTAL HAZARDS: This product is toxic to fish, birds and other wildlife. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

PHYSICAL OR CHEMICAL HAZARDS: Do not use or store near heat or open flame.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

Medical Information call 1-800-454-2333
 Product Information call 1-800-225-2883

ORTHO®,ORTHO-KLOR®,DIAL 'n SPRAY®, HomeDefense™ Trademarks of Monsanto Company
DURSBAN® Trademark of DowElanco

[©]Monsanto Company 1996

Manufactured For
The Solaris Group of
Monsanto Company
P.O. Box 5008
San Ramon CA 94583-0808
Form 07 741-02 [9/16/96]: 0-995-01
Product 5522
EPA Reg. No. 239-2513

239-2513,WP4 0-995-01

EPA Est. 239-IA-3¹, 58996-MO-1^A
Superscript is first letter of lot number
Made in USA