## [ICON] OUTDOOR USE ONLY

## ORTHO MULTIPURPOSE BORER AND INSECT SPRAY, EPA Reg. No. 239-2635

## Subset 1. ABN: ORTHO Borer & Leaf Miner Spray

Contains: Dursban Insecticide

Controls: Ants, Subterranean Termites, Powder Post Beetles, Crickets, Chinch Bugs, Wood Borers, Carpenter Ants, Grubs, Cutworms, Fleas, Armyworms, Ticks, Mosquitoes, Grasshoppers, European Crane Fly Larvae, Fire Ants, Sod Webworms, Spiders, Box Elder Bugs, Scorpions, And Many Other Soil Insects And Home Invading Pests.

For Use Around Foundations, Outside Buildings, On Lawns, Under Shrubs, On Woodpiles, Stored Lumber, Fence Posts, and Other Wooden Structures

For Use In Tank Sprayers, Hose-End Applicators, Or Sprinkling Cans Controls All Major Lawn Pests Concentrate

New, Water Based Formula

[ICON]: Water Based Formula/Water Based

Kills Lawn Insects

New Easy To Read Label

Low Odor

Low Odor Formula

Armyworms, Chiggers, Kills/Controls Home Invading Pests: Ants, Crickets, Cutworms, Earwigs, Fleas, Grasshoppers, Grubs, Fire Ants, Sod Webworms (Lawn Moth Larvae), Scorpions, Spiders, Ticks, and Many Other

Kills/Controls Ticks That Transmit Lyme Disease

Treats Up To 4,000 sq. ft (Qt. size)

Easy To Use Formula

Easy To Use Clear Formula

Easy To Handle Formula

New, Low Odor Formula

Helps Keep Your Lawn Pest/Insect Free

Controls Insect Pests That Can Cause Unsightly Brown Spots And Dead

Kills Insects That Damage Ornamentals, Shade Trees, And Fruit Trees

Kills Borers And Leafminers

Kills Insects: Peachtree Borer, Dogwood Borer, Lilac Borer, Leafminers,

Kills Pine Beetles, Bronze Birch & Peach Tree Borers and many other

Ideal For Use On Ornamenta As C Shane Trees And Fruit Trees

Use on shrubs and trees

Multi-Purpose

Contains DURSBAN

AUG 19 1998

Under the Federal Insections, Fungicide, and Redenticide Act, as amended for the pesticine

registered under 39 2635 BPA Reg. No. 239 2635

Amendment of 4/13/98; Natifications of 8/29/96, 8/30/96 and 9/10/96

239-2635.A4c

EPA Accepted 8/21/96

Jef 17

Effective Long Lasting Control Proven Effective

Can Be Applied As A Foliar Spray Can Be Applied As A Trunk Spray

Can Be Applied With Ortho Hose-End Applicators

DURSBAN Trademark of DowElanco

Makes Up To 96 Gals Of Diluted Spray

WARNING: KEEP OUT OF REACH OF CHILDREN

See back panel booklet for additional precautionary

statements

Active Ingredient

Net: 16 fl oz (1 pt) 473 ml.

32 fl oz (1 qt) 946 mL

#### FOR OUTDOOR HOME & GARDEN USE ONLY

ORTHO Multipurpose Borer and Insect Spray is an easy to apply, low odor, water based product to control lawn insects, home invading pests, fire ants, wood destroying insects including exposed termites, borers and leafminers that attack ornamental shrubs, fruit trees, and shade trees. For use in tank sprayers and hose-end sprayers.

PULL HERE TO OPEN LABEL. PRESS TO RESEAL. [For Peel/Reseal Labels only]

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

SHAKE WELL BEFORE USING.


tsp = teaspoon
Tbs = Tablespoon
2 Tbs = 1 fl oz
4 fl oz = 1/2 cup

WHEN TO APPLY

Treat when insects or their damage first appear. Repeat as needed.

WHERE & HOW TO APPLY

FOR OUTDOOR USE ONLY. Refer to the table for specific insects and tips on how to control. Use a hose end or tank sprayer to treat lawns, home invading pests, wood destroying insects, ornamental plants, fruit trees and shade trees. Use a sprinkling can to treat fire ant mounds.



239-2635\_A4c

EPA Accepted 8/21/96

Pests	Product To Measure	Area To Treat	Treatment Tips
LAWN PESTS: Note: Treat the entire lawn. Spot treatment may be less effective.			
Armyworm, Chiggers, Chinch Bugs, Clover Mites, Crickets, Cutworms, Dichondra Flea Beetles, Earwigs, European Crane Fly (Larvae), Fiery Skipper Larvae (California and Hawaii only), Fleas, Grasshoppers, Lawn Moth Larvae (Sod Webworms), Ticks, Tropical Grass Webworms (Hawaii only),	Tank Sprayer: 2 tsp. into 3 gals. of water and spray.  ORTHO Lawn Pest & Fertilizer Sprayer: 1-3/4 fl. oz. into container. Fill to the 15 gal. mark with water and spray.  DIAL 'n SPRAY Hose-End Applicator: 1-3/4 fl. oz. into container. DO NOT ADD WATER. Set dial at 2 tsp and spray.	100 sq.ft. (10'x 10')  500 sq.ft. (10'x 50')  500 sq.ft. (10' x 50')	Mow lawn and water thoroughly 1 to 2 days before treatment. Do not water for 2 to 3 days after application. Repeat treatment in 4 to 6 weeks if necessary. For European Crane Fly Larvae control, consult your Agricultural Extension Service for the best time to treat.
White Grubs: (Larvae of European Chafer and Japanese Beetle)	Tank Sprayer: 4 tsp. into 1- 1/2 gals. of water and spray.  ORTHO Lawn Pest & Fertilizer Sprayer: 6-1/2 fl. oz.(13 Tbs.) into container. Fill to the 15 gal. mark with water and spray.	50 sq. ft. (5'x 10') 500 sq.ft. (10'x 50')	Water lawn thoroughly after treatment. This will move the insecticide into the soil where the grubs feed. Treat when grubs are small and feeding near the soil surface. Consult your Agricultural Extension Service for the best time

239-2635.A4c

EPA Accepted 8/21/96

	DIAL 'n SPRAY Hose-End Applicator: 6-1/2 fl. oz.(13 Tbs) into container. DO NOT ADD WATER. Set dial at 1 oz. and spray.	500 sq.ft. (10"x 50")	to treat.
HOME INVADING PESTS OUTDOORS			
Ants, Crickets, Earwigs, Fleas, Millipedes, Sowbugs (Pillbugs), Spiders (including Black Widow Spiders), Scorpions, Ticks (including American Dog, Brown Dog, Cattle Fever, Deer, Gulf Coast, and Lone Star)	Tank Sprayer: 2 tsp. into 1 gal. of water and spray.  ORTHO Lawn Pest & Fertilizer Sprayer: 5 fl. oz. into container. Fill to the 15 gal. mark with water and spray.  DIAL 'n SPRAY Hose-End Applicator: 2-1/2 fl. oz. into container. DO NOT ADD WATER. Set dial	100 sq.ft. (10'x 10')  1500 sq.ft. (5'x 300')  750 sq. ft. (5'x 150')	To help prevent pests from entering buildings, treat a band of soil 5 feet wide around the building. Also treat the foundation up to a height of 2 to 3 feet. Not for use indoors or on pets.
·	at 2 tsp and spray.	130 )	
FIRE ANT MOUNDS			
Fire Ants	Sprinkling Can: 2 Tbs. (1 fl. oz.) into 1 gal. of water	1 mound (4 ft diameter circle)	Apply 1 gal. to each mound. Apply when ants are active. Treat new mounds as they appear.
			* * * * * * * * * * * * * * * * * * *

60f 17

TREATMENT FOR WOOD DESTROYING PESTS OUTDOORS			
Powderpost Beetles, False Powderpost Beetles, Furniture Beetles, Deathwatch Beetles, Carpenter Ants, Carpenter Bees, Apple Twig Borer, Longhorned Beetle	5-1/3 fl. oz. into 1 gal. of water	Treat localized areas of infested wood	Apply as a coarse drenching spray evenly on wood surfaces. Allow to dry and repeat treatment.
FENCE POSTS - TERMITE PREVENTION:	5-1/3 fl. oz. into 1 gal. of water	Apply one gallon of diluted product per post hole.	Pour one half of solution into post hole. Add the rest as you fill the hole with soil so that the fence post is surrounded by a protective barrier. Do not treat soil that is water saturated or frozen. Consult state and local specifications for recommended distances of treatment from wells.

239-2635.A4c

EPA Accepted 8/21/96

#### BORERS & LEAFMINERS ON TREES & SHRUBS

Application timing is important. Consult your County Cooperative Extension Agent for proper time to treat.

Pests	Trees Or Shrubs	Product to Measure When To Apply
Beetles (Cottonwood Leaf, Elm Leaf, Flea, Willow Leaf)	Cottonwood, Elm, Willow	4 tsp./gal. of water. Apply when pests appear. Thoroughly wet upper and lower leaf surfaces. Reapply at 7 to 10 day intervals if needed.
Weevils (Black Vine, Yellow Poplar, Pine)	Rhododendron, Poplar, Pine	
Borers (Ash, Dogwood, Lesser Peachtree, Lilac, Oak, Rhododendron, Bronze Birch, Flatheaded Appletree, Locust, Red Oak)	Ash, Lilac, Oak, Dogwood, Rhododendron, Birch, Locust	1-1/2 fl. oz. (3 Tbs.)/ gal. of water. Spray trunks and lower limbs of trees and shrubs when adults begin to emerge.
Leafminers	Elm, Birch, Locust	1-1/2 fl. oz. (3 Tbs.)/ gal. of water. Spray when leafminers appear in early and Mid- May.
Needleminers	Pine, Spruce	1-1/2 fl. oz. (3 Tbs.)/ gal. of water.
Beetles (Ambrosia, Black Turpentine, European Elm Bark, Japanese, June, Southern Pine, Spruce)	Elm, Pine Spruce	10-2/3 fl. oz. (1-1/4 cups)/gal. of water Apply with tank sprayer only. To prevent infestation, spray the tree trunk in early Spring. To treat infested trees, spray the tree trunk before adult beetles begin to emerge. Avoid contact with foliage to prevent leaf damage.

239-2635.A4c EPA Accepted 8/21/96

Application timing is important. Consult your County Cooperative Extension Agent for proper time to treat.

Pests	Fruit Trees	Product to Measure When to Apply
Dogwood Borer	Apples Wait 1 month after last application before harvesting fruit.	2 fl. oz. (4 Tbs.)/gal. of water. Soak tree trunks and the soil around the base of trees in the spring. Apply up to 8 times per season. Wait 3 weeks between treatments.
Borers (Lesser Peachtree, Great Peachtree, American Plum)	Sweet Cherries  Wait 1 week after application before harvesting fruit.	2 fl. oz. (4 Tbs.)/gal. of water. Spray trunks and lower limbs. Avoid contact with foliage to prevent leaf damage. Apply up to 3 times per season. Apply second application 2 weeks after the first. Make third application after harvest.
Borers (American Plum, Lesser Peachtree, Pacific Flatheaded, Peach Twig, Peachtree, Shothole)	Sour Cherries  Wait 2 weeks after application before harvesting fruit.	2 fl. oz. (4 Tbs.)/gal. of water. Spray tree trunks and lower limbs. Apply up to 8 times per season.
Peachtree Borers	Nectarines, Peaches  Wait 2 weeks after application before harvesting fruit.	4 fl. oz. (8 Tbs.)/gal. of water. Spray bark from the ground up to the lower branches. Do not spray fruit. Make only 1 application per year.

239-2635.A4c EPA Accepted 8/21/96

STORAGE AND DISPOSAL

STORAGE: Store in a secure, cool dry place. Keep from freezing.
DISPOSAL: Do not reuse empty container. Rinse thoroughly, wrap in newspaper and put in trash.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

WARNING: Causes substantial but temporary eye injury. Harmful if swallowed. Do not get in eyes or on clothing. Wear goggles or safety glasses. Avoid contact with skin. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

[Reentry Icon]: People and pets may re-enter treated areas after sprayed area has dried.

FIRST AID: IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists. IF SWALLOWED: Immediately telephone a poison control center, treatment center or a physician for advice. DO NOT make person vomit unless directed to do so by medical personnel. If medical advice cannot be obtained, then immediately take person and product container to an emergency treatment center or hospital. IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists. Note to Physicians: Emergency Information call 1-800-225-2883. Chlorpyrifos is a cholinesterase inhibitor. Atropine is antidotal. 2-PAM is antidotal and may be used in conjunction with atropine, but should not be used alone.

ENVIRONMENTAL HAZARDS: This product is toxic to fish and wildlife. Do not apply directly to water. Do not pour product or rinse water down any drain. This product is highly toxic to bees exposed to direct treatment or residues on blooming plants. Do not apply this product if bees are visiting the treatment area.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

Questions, Comments or Medical Information Call 1-800-225-2883 www.ortho.com?

©Trademark of Monsanto Company ©Monsanto Company 1996 Manufactured for The SOLARIS Group of Monsanto Company P.O. Box 5008 San Ramon CA 94583-0808

EPA Reg. No. 239-2635 EPA Est. 239-IA-3<sup>I</sup>, 58996-MO-001<sup>A</sup> Form Superscript is first letter of lot number

239-2635.A4c EPA Accepted 8/21/96 Amendment of 4/13/98; Notifications of 8/29/96, 8/30/96 and 9/10/96

## Subset 2. ABN: HomeDefense® ORTHO-KLOR Insect & Termite Killer1

[ICON] Outdoor Use Only FOR OUTDOOR, HOMEOWNER USE ONLY [ICON]

Some and the second of the sec

New, Water Based Formula Water Based Formula/Water Based [Icon] Lawn and Nuisance Pests Controls Wood-destroying Insects Concentrate Makes up to 548 Gallons of Spray [ 1/2 gal size only] Makes up to 274 Gallons of Spray [1 qt. size only]

For Termite Prevention, Treats 30 Linear Feet of Trench [1/2 gal] Treats 15 Linear Feet of Trench [quart]

Contains: DURSBAN® Soil Insecticide

CONTROLS: Ants, Carpenter Ants, , Powder Post Beetles, Crickets, Chinch Bugs, Wood Borers, Carpenter Bees, Grubs, Cutworms, Fleas, Armyworms, Ticks, Mosquitoes, Grasshoppers, European Crane Fly Larvae, Fire Ants, Sod Webworms, Spiders, Boxelder Bugs, Scorpions, and many other soil and home invading pests.

Plus Prevents Termite Infestation For Outdoor, Homeowner Post Construction Use Only

(FOR) USE: Around foundations, outside buildings, lawns, woodpiles , stored lumber and fence posts (For) Use in Tank Sprayers, Hose-End Sprayers, or Sprinkling Cans Controls All Major Lawn Pests Kills Lawn Insects Kills/Controls Ticks that Transmit Lyme Disease Helps Keep Your Lawn Healthy - Controls Insect Pests that Can Cause Unsightly Brown Spots And Dead Grass Kills/Controls Carpenter Ants

## Active Ingredient

Chlorpyrifos	.10.7%
Other Ingredients	.89.3%

## KEEP OUT OF REACH OF CHILDREN

WARNING See back panel booklet for additional precautionary statements.

NET	64	FL (	OZ (1	/2 GAL)	1.89	L
NET	32	FL C	)Z (1	QT)	946	mL
NET	16	FL (	DZ (I	PT)	473	mL

239-2635.A4c EPA Accepted \$/21/96

#### PEEL HERE

## Open Resealable Label for Directions & Precautions

#### PRESS TO RESEAL

Easy to apply with

•The Ortho® Dial `n Spray® [or] other hose-end sprayer (picture)

AND CONTROL OF THE PROPERTY OF

- •tank sprayer (picture)
- •sprinkling can (picture)

Alternative Sales Copy #1

Illustrate lawn & select lawn pests such as White Grubs, Chinch Bugs, Lawn Moths & Fleas.

Illustrate a Termite


Illustrate damaged wood infested by Carpenter Ants.

Illustrate a house foundation & select pests such as Spiders, Boxelder Bugs & Ticks.

See inside label for many more uses.

Alternative Sales Copy #2

WHERE TO USE	INSECTS/PESTS	
<ul> <li>Lawns</li> <li>Outdoor Garbage Cans</li> <li>Window Frames &amp; Doorways</li> <li>Porches &amp; Patios</li> <li>Eaves &amp; Foundations</li> <li>Fire Ant Mounds</li> <li>Wood Piles &amp; Stored Lumber</li> <li>Trench Around Building</li> </ul>	Ants, Boxelder Bugs, Carpenter Ants, Carpenter Bees, Centipedes, Chiggers, Chinch Bugs, Clover Mites, Cockroaches, Crickets, Cutworms, Earwigs, Fleas, Lawn Moths, Millipedes, Mosquitoes, Scorpions,	
Foundation [Treats 15 linear feet of Trench]	Spiders (including Black Widow and Brown Recluse Spiders), Subterranean Termites/ Termites (workers & winged), Ticks (including ticks which transmit Lyme Disease), White Grubs	
See inside label for	many more uses.	




120f17

For prevention of subterranean termites. Also controls borers, carpenter ants and carpenter bees found outdoors in localized areas of wooden structures, firewood piles and stored lumber. In addition controls many common pests in lawns and other outdoor areas around the home.

Specially formulated for residential use.

ORTHO-KLOR Soil Insect & Termite Killer<sub>1</sub> is an easy to apply product to control lawn insects, home-invading pests, fire ants & wood-destroying insects. When used as directed, it helps lawns stay vigorous, healthy, and green by controlling bugs that cause problems.

DO NOT USE INSIDE DWELLINGS



239-2635.A4c

EPA Accepted 8/21/96



DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

tsp = teaspoon

Tbs = Tablespoon = 3 teaspoons

Tbs = Tablespoon = 3 teaspoons

Tbs = Tablespoon = 3 teaspoons

SHAKE WELL [BEFORE USING]

#### LAWN PESTS

•Pests that live & feed underground:

European Crane Fly Larvae, Hyperodes (Turfgrass Weevils), Mole Crickets & White Grubs (Including larvae of European Chafer & Japanese Beetle)

•Pests that live & feed above ground:

Including Armyworms, Chiggers, Chinch Bugs, Clover Mites, Crickets, Cutworms, Dichondra Flea Beetles, Earwigs, Fiery Skipper Larvae (California and Hawaii only), Fleas, Grasshoppers, Lawn Moths (Sod Webworms), Ticks & Tropical Grass Webworms (Hawaii only)

## Mixing Instructions for Lawn Pests

Hyperodes, Mole Crickets & White Grubs:
Amount to Use: 2 Tbs (1 fl oz) per gallon of water for each 75 sq ft

. When using the Ortho® Dial 'n Spray®: Set dial to 1 oz. Add 13 fl oz (26 Tbs) to the sprayer jar. DO NOT add water. Spray evenly over 1000 sq ft until jar is empty.

<u>Pests that live & feed above ground and European Crane Fly Larvae:</u>

Amount to Use: 2 tsp per gallon of water for each 100 sq ft
.. When using the Ortho® Dial 'n Spray®: Set dial to 2 tsp.
Add 3½ fl oz (7 Tbs) to sprayer jar. DO NOT add water. Spray
evenly over 1000 sq ft until jar is empty.

## How to Apply

Measure lawn. Calculate square feet by multiplying length times width. [walking man icon]. Spray evenly over measured area.
Treat entire lawn. Spot treatment may be less effective.

#### When to Mow & Water

- •Apply to lawn that was mowed and watered within the last 24 hours.
- •Above Ground Pests: Do not water for 2 to 3 days after application.
- •Underground Pests: Water lawn thoroughly after application to

239-2635\_A4c EPA Accepted 8/21/96 Amendment of 4/13/98; Notifications of 8/29/96, 8/30/96 and 9/10/95

140f17

move the insecticide into the soil where these pests live and feed.

is a transport of the company of the contract of the contract

## When to Apply

•Wait 4 to 6 weeks between applications.

•European Crane Fly Larvae & White Grubs: Treat when pests are small and actively feeding near the soil surface. Consult your County Cooperative Extension Agent for the best time to treat.

#### FIRE ANT MOUNDS

#### Mixing Instructions for Fire Ant Mounds

Use a Sprinkling Can: Add 2 Tbs (1 fl oz) to 1 gallon of water for each mound.

## How to Apply

Apply evenly over and around each mound in a 4 ft diameter circle. Apply gently to avoid disturbing ants and causing them to move and form new mo

#### When to Apply

For best results apply in early morning or late afternoon when fire ants most active. Treat new mounds as they appear.

#### HOME-INVADING PESTS - OUTDOORS

including Ants, Bees, Beetles, Boxelder Bugs, Centipedes, Clover Mites, Cockroaches, Crickets, Earwigs, , Fleas, Flies, Hornets, House Flies, Millipedes, Scorpions, Silverfish, Sowbugs, Spiders (including Black Wid-Brown Recluse Spiders), Springtails, Ticks, Wasps & Yellow Jackets

## Mixing Instructions for Home-Invading Pests

Amount to Use: 2 tsp per gallon of water
When using the Ortho® Dial 'n Spray®: Set dial to 2 tsp. For 3
gallons of spray add 1 fl oz (2 Tbs) to sprayer jar. \* DO NOT add
water. Spray.

Note: After spraying, unused product can be poured back into its original container.

#### •How to Apply

•To keep insects from entering the house, spray a 3 foot(ft) band of soil around the house next to the foundation. Also spray the foundation of the house to a height of 2 feet(ft). '3.'gallons of spray treats 60 linear feet.

•Also spray walks, "garbage cans & garbage areas, "window, frames & doorways, "porches & patios, "under shrubs and 'frees but avoid spraying foliage. DO NOT treat pets with this product. DO NOT

239-2635.A4c EPA Accepted 8/21/96 Amendment of 4/13/98; Notifications of 8/29/96, 8/30/96 and 9/10/95

use indoors.

- Carrier - 1 - The second of the second of

#### MOSQUITOES - OUTDOORS

Mosquitoes coming to rest on areas treated for the control of lawn and home-invading pests will be killed for varying periods of time after application, depending on exposure of treated areas to weathering conditions.

#### CARPENTER ANTS

Plus Carpenter Bees & Wood Boring Beetles (including Powder Post Beetles) -OUTDOORS

Mixing Instructions for Carpenter Ants, Carpenter Bees & Wood Boring Beetles

## Amount to Use:

- . Wood Treatment: 5 1/3 fl oz (10 2/3 Tbs) per gallon of water
- . Perimeter Treatment: 2 tsp per gallon of water

# How to Apply Wood Treatment:

- Fences & Outdoor Wood Surfaces of Buildings. Apply evenly on wood surfaces with a paint brush. Allow to dry and repeat to ensure the insecticide soaks into the wood.
- Wood Piles & Stored Lumber: Use a hose-end or tank sprayer, or sprinkling can to deliver a coarse drenching spray. Treated wood can be burned or used for building one month after treatment.

Perimeter Treatment:

To keep insects from entering the house, use a hose-end or tank sprayer to spray a 3 foot (ft) band of soil around the house next to the foundation. Also spray the foundation of the house to a height of 2 feet. 3 gallons of spray treats 60 linear feet.

#### SUBTERRANEAN TERMITES - OUTDOORS

Prevents subterranean termite infestations. For outdoor, homeowner post construction use only. DO NOT use under slabs, in crawl spaces or indoors. If termites are found within the house or structure contact a licensed pest control operator for treatment.

Wixing Instructions for Termites
Use a Sprinkling Can: Add 5 1/3 fl oz (10 2/3 Tbs) to 1 gallon of water for each 2 1/2 linear feet\*.

\* 1 quart of HomeDefense® Ortho-Klor® will make 6 gallons of diluted solution and treat / cover 15 linear feet.

239-2635\_A4c EPA Accepted 8/21/96 Amendm

A CONTRACT OF MEMORY AND A SECOND OF A SEC

## How to Apply

#### Trench Treatment

- Remove all wood debris and wood forms.
- Poured Concrete Foundations: Dig a trench next to the foundation 6 inches wide and 4 inches deep or down to the top of the footing, whichever is less.
- Brick & Block Foundations:
- Dig a trench 6 inches wide and 12 inches deep.
- To avoid shifting your house foundation never dig below the top of the footing.
- Mix diluted product into soil as soil is replaced in the trench.

Fence Posts - (EXCEPT RHODE ISLAND) Using/use a sprinkling can. Pour 3 gallon of diluted product into fence post hole and add another ½ gallon to the soil when filling the hole so that all the fill soil is treated to form a protective barrier.

When to Apply

Reapplication for subterranean termites should only be made when there has been a disruption of the chemical barrier in the soil due to construction, excavation, landscaping, etc. Reapplication should be made as a spot treatment to those areas. Annual reapplication of the entire premises must be avoided.

#### IMPORTANT

Do not treat structures that contain cisterns or wells. Soil in the vicinity of wells, cisterns or ponds should not be treated. For those situations, contact your County Cooperative Extension Agent or consult a commercial pest control service.

- DO NOT treat soil that is water saturated or frozen or in any conditions where run-off or movement from the treatment area(site) is likely to occur.
- DO NOT apply while it is raining.

239-2635.A4c EPA Accepted 8/21/96



170/17

• DO NOT use on pets.

A SECTION OF THE PROPERTY OF T

• DO NOT use on edible or ornamental plants. Spray drift onto these plants should be rinsed off immediately with water.

・ そうことは大神を大手を

#### STORAGE AND DISPOSAL

STORAGE: Store in a secure, cool dry place. Keep from freezing. DISPOSAL: Do not reuse empty container. Rinse thoroughly, wrap in newspaper and put in trash.

#### PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

WARNING: Causes substantial but temporary eye injury. Harmful if swallowed. Do not get in eyes or on clothing. Wear goggles or safety glasses. Avoid contact with skin. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

[8/30/96]: [Reentry Icon]: People and pets may re-enter treated areas after sprayed area has dried.

FIRST AID: IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists. IF SWALLOWED: Immediately telephone a poison control center, treatment center or a physician for advice. DO NOT make person vomit unless directed to do so by medical personnel. If medical advice cannot be obtained, then immediately take person and product container to an emergency treatment center or hospital. IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists. Note to Physicians:

Emergency Information call 1-800-225-2883. Chlorpyrifos is a cholinesterase inhibitor. Atropine is antidotal. 2-PAM is antidotal and may be used in conjunction with atropine, but should not be used alone.

ENVIRONMENTAL HAZARDS: This product is toxic to fish and wildlife. Do not apply directly to water. Do not pour product or rinse water down any drain.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

Trademark of Monsanto Company

Monsanto Company 1998

Manufactured for

The SOLARIS Group of

Monsanto Company

P.O. Box 5008

San Ramon CA 94583-0808

EPA Reg. No. 239-2635

EPA Est. 239-IA-3<sup>I</sup>, 58996-MO-001<sup>A</sup>

Superscript is first letter of lot number

