

In case of an emergency endangering
life or property involving this
product, call collect, day or night,
Area code 201 - 835-3100.

Antidote: Atropine is an antidote. Ask your physician about obtaining a supply of 1/100 grain (0.6 milligram) tablets for emergency use. If symptoms include blurred vision, stomach cramps or tightness in chest, don't wait for a physician but take two tablets at once. Do not take atropine unless symptoms of poisoning have occurred. Anyone who has been sick enough to have taken atropine must be seen by a physician as soon as possible.

FIRST AID

If swallowed, induce vomiting by sticking finger down throat or by giving soap or strong salty water to drink. Repeat until vomited clear. Never give anything by mouth to an unconscious person.
If inhaled, remove to fresh air. If not breathing, give artificial respiration, preferably mouth to mouth. If breathing is difficult, give oxygen.
In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes while removing contaminated clothing and shoes. Wash clothing before re-use.

CALL A PHYSICIAN AT ONCE IN ALL CASES OF SUSPECTED POISONING.

NOTE TO PHYSICIANS: Warning symptoms include weakness, headache, tightness of chest, blurred vision, nonreactive pinpoint pupils, salivation, sweating, nausea, vomiting, diarrhea and abdominal cramps. Give atropine intramuscularly or intravenously, depending on severity of poisoning, 2 to 4 milligrams (3 to 6 tablets, 1/100 grain each) every 15 minutes until fully atropinized as shown by dilated pupils, dry flushed skin and tachycardia. Twenty to thirty milligrams or more may be required during the first 24 hours. Never give opiates or phenothiazine tranquilizers. Clear chest by postural drainage. Artificial respiration or oxygen administration may be necessary. Observe patient continuously for at least 48 hours. Allow no further exposure to any cholinesterase inhibitor until cholinesterase regeneration has taken place as determined by blood tests. Pralidoxime Hydrochloride (2-PAM PROTOPAM chloride) may be effective as an antidote to atropine. Use according to label directions.

INDICATIONS FOR USE

Do not use this product for any uses other than those specified on this label. BNYH is an effective soil and systemic insecticide for early season insect control which has been combined with fertilizer for efficient and uniform application. Application should be made at planting time. Application at rates higher than those shown on label may result in illegal crop residues.

WHEAT: For control of grasshoppers, aphids and Hessian fly, apply at the rate of 100 lbs. (1.0 B) actual liberate per acre in the seed furrow at planting time. Do not feed or graze foliage within 14 days of treatment. For control of grasshopper control, see label directions and instructions on label.

WARRANTY

American Cyanamid Company warrants that this product, when used in accordance with the directions on the label, conforms to the description of the product on the label and is generally fit for the use thereon described, except insofar as it may be limited by the directions on the label. Any damages arising from the use of this product shall be limited to a refund of the purchase price and shall not include consequential damages or any loss of profits or value of crops. American Cyanamid Company does not warrant that this product will be effective against any pest or disease other than those specified on the label. BNYH is a registered trademark of American Cyanamid Company. BNYH assumes the risk of any loss or damage resulting from the use of this product under abnormal conditions of use or in violation of the directions on the label. American Cyanamid Company is not liable for any such loss or damage.

© 1974

AMERICAN CYANAMID COMPANY
NEW YORK, N.Y. 10017
PHILADELPHIA, PA. 19104

14