

EPA Reg. No. 241-350

1-829
1-11-25
241-350

Under the Federal Insecticide, Fungicide, and Plant Disease Act as amended, for the pesticide registered under EPA Reg. No. 241-350

FOR USE IN SUCCULENT PEAS, DRY EDIBLE PEAS, LENTILS, AND CHICKPEAS

PURSUIT DG can be used on succulent peas, dry edible peas (including chickpeas) and lentils in Washington (Adams, Asotin, Columbia, Garfield, Spokane, Walla Walla and Whitman counties), Idaho (Benewah, Clearwater, Idaho, Kootenai, Latah, Lewis, and Nez Perce counties), Oregon (Union, Umatilla and Wallowa counties) and Montana only.

THIS LABEL EXPIRES DECEMBER 31, 1996

OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.

GENERAL INFORMATION

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT DG application to peas and lentils. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans. Plant peas or lentils at least 1/2 inch deep to reduce risk of crop injury.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

*Registered Trademark of American Cyanamid Company

(see reverse side)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT DG in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

Spray equipment used for PURSUIT DG applications must be drained and thoroughly cleaned with water before being used to apply other products.

DO NOT make more than one application of PURSUIT DG per year.

Allow at least 30 days between application and harvest of succulent peas.

Allow at least 60 days between application and harvest of dry edible peas, chickpeas, and lentils.

When PURSUIT DG is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

BEST AVAILABLE COPY

Apply PURSUIT DG herbicide at a broadcast rate of 1 soluble bag (7.2 ounces per 6 2/3 acres) for preplant incorporated and preemergence applications. At this broadcast rate, one soluble bag of PURSUIT DG will treat 6 2/3 acres of edible beans.

APPLICATION INSTRUCTIONS

Preplant Incorporated Applications. Apply PURSUIT DG within 1 week before planting. **DO NOT** incorporate deeper than 3 inches. Refer to the PURSUIT DG leaflet label for incorporation instructions.

Preemergence Applications. Apply PURSUIT DG after planting, but prior to crop emergence.

PURSUIT DG may be tank-mixed with Sencor DF¹ or Lexone DF² to assist in the control of lambsquarters or mayweed-chamomile (dogfennel). Refer to the Sencor or Lexone label for proper application rates and restrictions.

WEEDS CONTROLLED

PURSUIT DG applied preplant incorporated or preemergence will control:

Buckwheat, wild	Nightshade, Eastern black
Kochia	Nightshade, hairy
Lambsquarters, common*	Pigweed, redroot
Mustard, wild	Shepherdspurse
Nightshade, black	Thistle, Russian

*preplant incorporated applications only

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label:

- Anytime

Chickpeas	Peanuts
Lentils	Soybeans
Peas	IMI-Corn
Lima beans	
- Four months after PURSUIT DG application:

Barley	Rye
Edible beans (dry, snap)	Wheat

Tobacco

- Eighteen months after PURSUIT DG application:

Alfalfa	Popcorn
Cotton	Safflower
Lettuce	Sorghum
Oats	Sweet Corn
- Twenty-six months after PURSUIT DG application:
All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE.
- Forty months after PURSUIT DG application:
Sugar beets

In the event of a crop loss, peas, lentils or chickpeas may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

****Several seed companies have tested a wide range of in-breds for sensitivity to imazethapyr soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with imazethapyr the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into field treated previously with imazethapyr shall be assumed by the user.**

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.

¹Sencor is a trademark of the Parent Company of Farbenfabriken Bayer GmbH, Leverkusen.
²Lexone is a trademark of E. I. duPont de Nemours and Company, Inc.

3-29

EPA Reg. No. 241-350

**FOR USE IN GREAT NORTHERN, RED KIDNEY AND PINTO
TYPE DRY BEANS**

FOR DISTRIBUTION AND USE ONLY IN NEW MEXICO

THIS LABEL EXPIRES APRIL 30, 1996

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN
THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

GENERAL INFORMATION

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT DG application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application. **DO NOT** apply PURSUIT DG postemergence after crop has begun to flower or crop injury may result.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

*Dash is a trademark of BASF Corporation
*Registered Trademark of American Cyanamid Company

(see reverse side)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT DG in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

Spray equipment used for PURSUIT DG application must be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for post applications.

DO NOT use crop oils, petroleum oils, liquid fertilizers or Dash¹.

DO NOT make more than one application of PURSUIT DG per year. **DO NOT** apply PURSUIT DG after June 30. Allow at least 60 days between application and harvest.

BEST AVAILABLE COPY

241-350

tive label restrictions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

APPLICATION RATE AND TIMING

Preemergence Applications. Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre immediately after, or up to 1 day after planting. PURSUIT DG may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide. **DO NOT** apply PURSUIT DG on sand or loamy sand soils.

Postemergence Applications. Apply PURSUIT DG at 1.08 ounces per acre. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture.

DO NOT APPLY PURSUIT DG POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRUE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.

ECO-PAK CONVERSION TABLE

Use Rate (oz/acre)	1 ECO-PAK Soluble Bag Treats
1.08	6-2/3 acres

WEEDS CONTROLLED

PURSUIT DG applied at the broadcast rate of 1.08 ounces per acre preemergence or early postemergence will control:

Mustard, wild	Nightshade, hairy
Nightshade, black	Pigweed, redroot
Nightshade, Eastern black	

Postemergence applications of 1.08 ounces per acre must be to weeds less than 2 inches tall for best results.

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label:

- Anytime
 - Soybeans
 - Peanuts
 - Edible bean types listed on this label (Refer to heading on page 1).
 - IMI-Corn

PE-XXXXX 12/4/94 SD

Peas
Rye
Wheat

- Nine and one-half months after PURSUIT DG application:
 - Field corn
 - Field corn grown for seed*
 - Tobacco
- Eighteen months after PURSUIT DG application:
 - Alfalfa
 - Cotton
 - Lettuce
 - Oats
 - Popcorn
 - Safflower
 - Sorghum
 - Sweet Corn
- Twenty-six months after PURSUIT DG application:
 - All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE.
- Forty months after PURSUIT DG application:
 - Sugar beets
 - Red table beets

In the event of a crop loss Great Northern, red kidney or pinto beans may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

* Several seed companies have tested a wide range of in-breds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.



BEST AVAILABLE COPY

EPA Reg. No. 241-350

**FOR USE IN NAVY, GREAT NORTHERN, RED KIDNEY,
BLACK TURTLE, CRANBERRY, AND PINTO TYPE DRY BEANS,
AND DRY EDIBLE PEAS ONLY**

**FOR DISTRIBUTION AND USE IN NORTH DAKOTA
AND NORTHERN MINNESOTA (see USE AREA below)**

THIS LABEL EXPIRES APRIL 30, 1996

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN
THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

GENERAL INFORMATION

PURSUIT DG application methods and use rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT DG only to the legume vegetables specifically listed in this label. **DO NOT** apply PURSUIT DG in geographical regions outside those specifically defined in this label for each type of legume vegetable.

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT DG application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application.

DO NOT apply PURSUIT DG postemergence after crop has begun to flower or crop injury may result.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Under some conditions (such as heavy texture soil, high organic matter, low pH or low rainfall) PURSUIT DG may cause injury to subsequent planted crops. Sugar beets are particularly sensitive to PURSUIT DG residues in soil. PURSUIT DG is not recommended in crop rotations that include sugarbeets. See the **ROTATIONAL CROP** section of this supplemental label for rotations to other sensitive crops.

Dash is a trademark of BASF Corporation
*Registered Trademark of American Cyanamid Company

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

USE AREA

PURSUIT DG may be used in North Dakota and in Minnesota north of highway #210.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT DG in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, potatoes, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

Spray equipment used for PURSUIT DG application must be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for postemergence applications. **DO NOT** use crop oils, petroleum oils, liquid fertilizers or Dash¹.

DO NOT make more than one application of PURSUIT DG per year.

(see reverse side)

BEST AVAILABLE COPY

241350

Pinto varieties UI-111 and Olathe are more sensitive to PURSUIT DG than other varieties.

DO NOT apply PURSUIT DG after July 10.

DO NOT feed treated forage, hay, or straw to livestock.

Allow at least 60 days between application and harvest.

Postemergence Applications. Apply PURSUIT DG at 0.72 ounces per acre (one ECO-PAY soluble bag treats 10 acres). A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture.

DO NOT APPLY PURSUIT DG POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRIFOLIATE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.

WEEDS CONTROLLED

-) PURSUIT DG applied at the broadcast rate of 0.72 ounces per acre will control:
- wild mustard
 - black nightshade*
 - eastern black nightshade*
- *suppression

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label:

1. Anytime
 - Soybeans
 - IMI-Corn
 - Edible bean and pea types listed on this label (Refer to heading on page 1)
2. Nine and one-half months after PURSUIT DG application:
 - Field corn
 - Wheat
 - Field corn grown for seed**
 - Edible bean and pea types not listed on this label (Refer to heading on page 1)
3. Eighteen months after PURSUIT DG application:
 - Barley
 - Oats
 - Rye
4. Twenty-six months after PURSUIT DG application:
 - All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE.
5. Forty months after PURSUIT DG application:
 - Sugar beets
 - Red table beets

BEST AVAILABLE COPY

****Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.**

) **CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.**

PE-XXXXX 12/5/94 SD

 **CYANAMID**
American Cyanamid Company
Agricultural Products Division
Crop Protection Chemicals Department
Wayne, NJ 07470 ©1994

705-9
EPA Reg. No. 241-350

FOR USE IN GREAT NORTHERN AND RED KIDNEY TYPE DRY BEANS

**FOR DISTRIBUTION AND USE ONLY IN COLORADO, KANSAS, NEBRASKA,
WYOMING, TEXAS AND WISCONSIN**

THIS LABEL EXPIRES APRIL 30, 1996

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN
THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

GENERAL INFORMATION

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT DG application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application.

DO NOT apply PURSUIT DG postemergence after crop has begun to flower or crop injury may result.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

¹Dash is a trademark of BASF Corporation

*Registered Trademark of American Cyanamid Company

(see reverse side)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT DG in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

Spray equipment used for PURSUIT DG application must be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for post applications.

DO NOT use crop oils, petroleum oils, liquid fertilizers or Dash¹.

DO NOT make more than one application of PURSUIT DG per year. **DO NOT** apply PURSUIT DG after June 30. Allow at least 60 days between application and harvest.

BEST AVAILABLE COPY

241-350

ive label restrictions and precautions. NO label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

APPLICATION RATE AND TIMING

Preemergence Applications. Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre immediately after, or up to 1 day after planting. PURSUIT DG may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide. **DO NOT** apply PURSUIT DG on sand or loamy sand soils.

Postemergence Applications. Apply PURSUIT DG at 1.08 ounces per acre. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture.

DO NOT APPLY PURSUIT DG POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRUE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.

ECO-PAK CONVERSION TABLE

<u>Use Rate (oz/acre)</u>	<u>1 ECO-PAK Soluble Bag Treats</u>
1.08	6-2/3 acres

WEEDS CONTROLLED

PURSUIT DG applied at the broadcast rate of 1.08 ounces per acre preemergence or early postemergence will control:

Mustard, wild	Nightshade, hairy
Nightshade, black	Pigweed, redroot
Nightshade, Eastern black	

Postemergence applications of 1.08 ounces per acre must be to weeds less than 2 inches tall for best results.

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label:

1. Anytime
 - Soybeans
 - Peanuts
 - Edible bean types listed on this label (Refer to heading on page 1).
 - IMI-Corn

Rye
Wheat

3. Nine and one-half months after PURSUIT DG application:
 - Field corn
 - Field corn grown for seed*
 - Tobacco
4. Eighteen months after PURSUIT DG application:
 - Alfalfa
 - Cotton
 - Lettuce
 - Oats
 - Popcorn
 - Safflower
 - Sorghum
 - Sweet Corn
5. Twenty-six months after PURSUIT DG application:
 - All crops not listed in this ROTATIONAL CROP GUIDELINE.
6. Forty months after PURSUIT DG application:
 - Sugar beets
 - Red table beets

DO NOT feed treated forage, hay, or straw to livestock.

* Several seed companies have tested a wide range of in-breds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.

BEST AVAILABLE COPY



EPA Reg. No. 241-350

9-879

FOR USE IN NAVY, GREAT NORTHERN, RED KIDNEY, BLACK TURTLE, CRANBERRY, PINTO, AND SMALL WHITE TYPE DRY BEANS, LIMA BEANS, DRY EDIBLE PEAS, ENGLISH PEAS, LENTILS, AND CHICKPEAS

FOR DISTRIBUTION AND USE IN MINNESOTA (south of highway #210) AND SOUTH DAKOTA

THIS LABEL EXPIRES APRIL 30, 1996

U.S. Federal Insecticide, Fungicide, and Rodenticide Act
Registered under
EPA Reg. No. 241-350

OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.

GENERAL INFORMATION

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT DG application to legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application.

DO NOT apply PURSUIT DG postemergence after crop has begun to flower or crop injury may result.

Some crop varieties may be inherently less tolerant to PURSUIT DG applications and crop injury could result. Contact the seed company for information and recommendations about the use of PURSUIT DG postemergence.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

¹Dash is a Trademark of BASF Corporation.

²Registered Trademark of American Cyanamid Company

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT DG in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, potatoes, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

Spray equipment used for PURSUIT DG applications must be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for post applications. **DO NOT** use crop oils, petroleum oils, liquid fertilizers or Dash¹

DO NOT apply to Domino variety black turtle beans.

DO NOT make more than one application of PURSUIT DG per year.

Allow at least 30 days between application and harvest of lima beans and english peas. Allow at least 60 days between application and harvest of dry beans, dry edible peas, lentils and chickpeas.

When PURSUIT DG is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

(see reverse side)

BEST AVAILABLE COPY

at the broadcast rate of 1.08 ounces per acre within 1 week before planting. DO NOT incorporate deeper than 3 inches. Refer to the PURSUIT DG leaflet label for incorporation instructions.

DO NOT apply preplant incorporated to pinto beans.

Preemergence Applications. Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre after planting, but prior to crop emergence. If weeds are present at the time of application, a burndown herbicide such as Roundup¹ or Gramoxone Extra² should be tank-mixed with PURSUIT DG to control emerged weeds.

DO NOT apply preemergence to pinto beans.

Postemergence Applications. FOR USE IN ENGLISH PEAS, NAVY, PINTO, GREAT NORTHERN AND RED KIDNEY TYPE DRY BEANS ONLY. Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture. DO NOT APPLY PURSUIT DG POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRIFOLIATE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.

ECO-PAK CONVERSION TABLE

Use Rate (oz/acre)	1 ECO-PAK Soluble Bag Treats
1.08	6-2/3 acres

WEEDS CONTROLLED

PURSUIT DG applied at the broadcast rate of 1.08 ounces per acre preplant incorporated, preemergence or postemergence will control:

Mustard, wild	Nightshade, hairy
Nightshade, black	Pigweed, redroot
Nightshade, Eastern black	

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label:

- Anytime
 - Chickpeas
 - Lentils
 - Dry edible peas
 - Peanuts
 - Edible beans (dry and lima listed on this label)
 - Soybeans
 - IMI-Corn

¹Roundup is a trademark of Monsanto Agricultural Products Company
²Gramoxone Extra is a trademark of ICI Americas, Inc.

- Wheat
- Nine and one-half months after PURSUIT DG application:
 - Field corn
 - Field corn grown for seed*
 - Tobacco
 - Eighteen months after PURSUIT DG application:

Alfalfa	Oats	Sorghum
Cotton	Popcorn	Sweet Corn
Lettuce	Safflower	
 - Twenty-six months after PURSUIT DG application:
 - All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE.
 - Forty months after PURSUIT DG application:

Sugar beets	Red table beets
-------------	-----------------

In the event of a crop loss, edible beans, dry edible peas, lentils or chickpeas may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

*Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.



CYANAMID

American Cyanamid Company
 Agricultural Products Division
 Crop Protection Chemicals Department
 Wayne, NJ 07470 ©1994

BEST AVAILABLE COPY

EPA Reg. No. 241-350

FOR USE IN NAVY, GREAT NORTHERN, RED KIDNEY, BLACK TURTLE, CRANBERRY, SMALL WHITE TYPE DRY BEANS, LIMA BEANS, SOUTHERN AND ENGLISH PEAS, AND WHITE LUPINS

FOR DISTRIBUTION AND USE IN WISCONSIN ONLY

THIS LABEL EXPIRES APRIL 30, 1996

OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

GENERAL INFORMATION

PURSUIT DG application methods and use rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT DG only to the legume vegetables specifically listed in this label. **DO NOT** apply PURSUIT DG in geographical regions outside those specifically defined in this label for each type of legume vegetable.

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT DG application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application.

DO NOT apply PURSUIT DG postemergence after crop has begun to flower or crop injury may result.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.


REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended:

- 1. Two months after PURSUIT DG application:

Snap beans



PE XXXXX 12/4/94 SD



BEST AVAILABLE COPY

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended for the pesticide
registered under
EPA Reg. No. 241-350

EPA Reg. No. 241-350

12 7 29

FOR USE IN NAVY, GREAT NORTHERN, RED KIDNEY, BLACK TURTLE, CRANBERRY, AND SMALL WHITE TYPE DRY BEANS, LIMA BEANS, AND SOUTHERN AND ENGLISH PEAS ONLY

REFER TO THE MAP ON PAGE 6 FOR GEOGRAPHICAL USE AREA

READ LABEL CAREFULLY - USE INSTRUCTIONS VARY FOR EACH GROUP

OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN
THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.

GENERAL INFORMATION

PURSUIT DG application methods and use rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT DG only to the legume vegetables specifically listed in this label. DO NOT apply PURSUIT DG in geographical regions outside those specifically defined in this label for each type of legume vegetable.

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT DG application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. DO NOT apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application.

DO NOT apply PURSUIT DG postemergence after crop has begun to flower or crop injury may result.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

¹Dash is a trademark of BASF Corporation.

²Registered Trademark of American Cyanamid Company

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT DG in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

Spray equipment used for PURSUIT DG application must be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for postemergence applications. DO NOT use crop oils, petroleum oils, liquid fertilizers or Dash¹.

BEST AVAILABLE COPY

rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

USE RATE

0.72, 1.08 OR 1.44 OUNCES PER ACRE

Depending on soil type and edible bean or pea type, apply PURSUIT DG at a broadcast rate according to the following table.

Use Rate (oz/acre)	LECO-PAK Soluble Bag Treats
0.72	10 acres
1.08	6-2/3 acres
1.44	5 acres

LIMA BEANS

DO NOT apply PURSUIT DG after June 30.

Allow at least 30 days between application and harvest.

Preplant Incorporated Applications. Apply PURSUIT DG at the broadcast rate of 0.72 ounces per acre to sand and loamy sand soils, or 1.08 ounces per acre to all other soils within 1 week before planting. Applied preplant incorporated, PURSUIT DG may be tank-mixed with a registered grass herbicide.

Preemergence Applications. Apply PURSUIT DG at the broadcast rate of 0.72 ounces per acre to sand and loamy sand soils, or 1.08 ounces per acre to all other soils immediately after, or up to 1 day after planting. PURSUIT DG may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide.

NAVY, GREAT NORTHERN, RED KIDNEY, CRANBERRY, BLACK TURTLE, AND SMALL WHITE TYPE DRY BEANS

DO NOT apply to Domino variety black turtle beans.

DO NOT apply PURSUIT DG to sand or loamy sand soils.

DO NOT apply PURSUIT DG after June 20.

Allow at least 60 days between application and harvest.

Preplant Incorporated Applications. Apply PURSUIT DG within 1 week before planting. Applied preplant incorporated, PURSUIT DG may be tank-mixed with a registered grass herbicide. Refer to the following table for use rate.

All other soils with > 2% O.M.

1.08

Preemergence Applications. Apply PURSUIT DG at the rates shown in the preceding table immediately after, or up to 1 day after planting. PURSUIT DG may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide.

ENGLISH PEAS

DO NOT apply PURSUIT DG to sand or loamy sand soils.

Allow at least 30 days between application and harvest.

Preplant Incorporated Applications. Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre up to 1 week before planting. DO NOT apply PURSUIT DG to English peas that have been or will be treated with trifluralin.

Preemergence Applications. Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre immediately after, or up to 1 day after planting. DO NOT apply PURSUIT DG if English peas have been treated with trifluralin.

Postemergence Applications. Apply PURSUIT DG at 1.08 ounces per acre. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture. DO NOT APPLY PURSUIT DG POST-EMERGENCE BEFORE CROP HAS AT LEAST ONE TRUE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT. If English peas have been treated with trifluralin, PURSUIT DG may be applied ONLY at 0.72 ounces per acre for suppression of black nightshade. The use of trifluralin prior to PURSUIT DG application may increase the likelihood and severity of crop injury.

SOUTHERN PEAS

Allow at least 30 days between application and harvest.

Preplant Incorporated Applications. Apply PURSUIT DG at the broadcast rate of 1.44 ounces per acre within 1 week before planting. Applied preplant incorporated, PURSUIT DG may be tank-mixed with a registered grass herbicide.

Preemergence Applications. Apply PURSUIT DG at the broadcast rate of 1.44 ounces per acre immediately after, or up to 1 day after planting. PURSUIT DG may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide.

Early Postemergence Applications. Apply PURSUIT DG at the broadcast rate of 1.44 ounces per acre when peas are at least 3 inches in height. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture

PURSUIT DG applied at the broadcast rate of 1.08 ounces per acre preplant incorporated, preemergence or early postemergence will control:

- | | |
|---------------------------|-------------------|
| Mustard, wild | Nightshade, hairy |
| Nightshade, black | Pigweed, redroot |
| Nightshade, Eastern black | |

WEEDS CONTROLLED

PURSUIT DG applied to low organic matter soils at the broadcast rate of 0.72 ounces per acre preplant incorporated or preemergence will control light to moderate infestations of:

- Nightshade, black
- Nightshade, Eastern black
- Pigweed, redroot

Postemergence applications of 1.08 ounces per acre must be made to weeds less than 2 inches tall for best results.

When applied as directed at the broadcast rate of 1.44 ounces per acre, PURSUIT DG will control or reduce competition from the weeds listed below:

NOTE: C = Control, R = Reduced Competition

(The number under Maximum Leaf Stage indicates the MAXIMUM number of leaves at which weeds should be sprayed postemergence.)

BROADLEAF WEEDS

WEEDS CONTROLLED	SOIL APPLIED	POSTEMERGENCE	
		Maximum Leaf Stage	Size (inches)
Anoda, spurred	C	2	1 - 2
Artichoke, Jerusalem		8	6 - 10
Buffalobur	C**		
Bristly starbur		2	1 - 2
Carpetweed	C		
Cocklebur, common	C**	8	1 - 8
Galinsoga	C		
Jimsonweed	C'	4	1 - 3
Kochia	C	4	1 - 3
Lambsquarters	C'	R	1 - 2
Mallow, Venice	R		
Morningglory,			
entireleaf	R	2	1 - 2
ivyleaf	R	2	1 - 2
pitted	R	2	1 - 2
smallflower	C	4	1 - 3
tall	R	2	1 - 2
Mustard sp.	C	4	1 - 3
Nightshade,			
black	C	4	1 - 3
Eastern black	C	4	1 - 3
hairy	C	4	1 - 3
Pigweed,			
redroot	C	8	1 - 8
smooth	C	8	1 - 8
spiny	C	8	1 - 8
Poinsettia, wild	C		
Puncturevine	C		
Purslane, common	C		
Pusley, Florida	C		
Sida, prickly	C'		

BEST AVAILABLE COPY

BROADLEAF WEEDS (Continued)

WEEDS CONTROLLED	SOIL APPLIED	POSTEMERGENCE	
		Maximum Leaf Stage	Size (inches)
Ragweed,			
common	R	4	1 - 3
giant	R	4	1 - 3
Sage, barnyard		R	1 - 3
Smartweed,			
ladysthumb	C	4	1 - 3
Pennsylvania	C	4	1 - 3
Spurge,			
prostrate	C	4	1 - 3
spotted	C	4	1 - 3
Sunflower, common	C*	4	1 - 3
Velvetleaf	C*	4	1 - 3
Thistle, Canada		R	1 - 3

*When soil applied, common lambsquarters, jimsonweed, prickly sida, velvetleaf and common sunflower are more consistently controlled by preplant incorporated treatments.

**Use soil applications for light to moderate infestations only. Must be preplant incorporated for best results.

GRASS WEEDS*

WEEDS CONTROLLED	SOIL APPLIED	POSTEMERGENCE	
		Maximum Leaf Stage	Size (inches)
Barnyardgrass	R	3	1 - 3
Crabgrass,			
large	R	3	1 - 3
smooth	R	3	1 - 3
Cupgrass, woolly		3**	1 - 3
Foxtail,			
giant	C	6	1 - 6
green	C	3	1 - 3
giant green	C	3	1 - 3
robust purple	C	3	1 - 3
robust white	C	3	1 - 3
yellow	C	3	1 - 3
Goosegrass	R		
Johnsongrass,			
seedling	C	6	1 - 8
rhizome		R	1 - 8
Panicum,			
fall	R		
Texas	R		

*When soil applied to grasses, more consistent control can be obtained from preplant incorporated treatments.

**PURSUIT DG controls emerged woolly cupgrass only.

GRASS WEEDS (Continued)

WEEDS CONTROLLED	SOIL APPLIED	POSTEMERGENCE	
		Maximum Leaf Stage	Size (inches)
Red rice		3	1 - 3
Shattercane	R	6	1 - 8
Signalgrass, broadleaf	R	4	1 - 8
SEDGES			
Nutsedge,			
purple	R	R	1 - 3
yellow	R	R	1 - 3

DO NOT count cotyledon leaves when determining weed stage of growth.

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label:

1. Anytime
 - Soybeans
 - Peanuts
 - IMI-Corn
 - Edible bean and pea types listed on this label (Refer to heading on page 1)
2. Two months after PURSUIT DG application:
 - Snap beans (DE, MD, PA and VA only)
3. Four months after PURSUIT DG application:
 - Barley
 - Edible bean and pea types not listed on this label (Refer to heading on page 1)
 - Rye
 - Wheat
4. Nine and one-half months after PURSUIT DG application:
 - Field corn
 - Field corn grown for seed*
 - Tobacco
5. Eighteen months after PURSUIT DG application:

Alfalfa	Popcorn
Cotton	Safflower
Lettuce	Sorghum
Oats	Sweet Corn
6. Twenty-six months after PURSUIT DG application:
 - All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE.
7. Forty months after PURSUIT DG application:

Sugar beets	Red table beets
-------------	-----------------

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

BEST AVAILABLE COPY

147

ROTATIONAL CROP GUIDELINE (Continued)

In the event of a crop loss, the bean or pea crop may be replanted.


DO NOT feed treated forage, hay, or straw to livestock.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.


USE AREA MAP

PURSUIT DG can be applied to the edible legume vegetables listed only in the states or parts of states shaded in the following map:

Use area includes: South Dakota east of U.S. 81, Nebraska east of U.S. 81, Kansas east of U.S. 81, Maryland and Virginia EXCLUDING THE DELMARVA PENINSULA, and all other states or portions of states shaded in the following map.



PE-17242 Rev. 1 12/04 SD

 **CYANAMID**
American Cyanamid Company
Agricultural Products Division
Crop Protection Chemicals Department
Wayne, NJ 07470 ©1994

EPA Reg. No. 241-350

**FOR PREEMERGENCE USE IN SNAP BEANS GROWN
IN NEW JERSEY ONLY**

THIS LABEL EXPIRES APRIL 30, 1997

OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.

GENERAL INFORMATION

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT DG application to snap beans. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT DG in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

Spray equipment used for PURSUIT DG applications must be drained and thoroughly cleaned with water before being used to apply other products.

DO NOT apply PURSUIT DG after June 20.

Allow at least 30 days between application and harvest.

DO NOT make more than one application of PURSUIT DG per year.

When PURSUIT DG is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

(see reverse side)

BEST AVAILABLE COPY

241-350

broadcast rate of 0.54 ounces per acre (one ECO-PAK soluble bag treats 13-1/3 acres) immediately after, or up to 1 day after planting. PURSUIT DG may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide.

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label:

1. Anytime

Soybeans
Peanuts
Edible beans and peas
IMI-Corn

2. Four months after PURSUIT DG application:

Barley
Rye
Wheat

3. Nine and one-half months after PURSUIT DG application:

Field corn
Field corn grown for seed*
Tobacco

4. Eighteen months after PURSUIT DG application:

Alfalfa	Popcorn
Cotton	Safflower
Lettuce	Sorghum
Oats	Sweet Corn

5. Twenty-six months after PURSUIT DG application:

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE.

6. Forty months after PURSUIT DG application:

Sugar beets Red table beets

In the event of a crop loss, snap beans may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.

PE-XXXXX 12/94 SD

BEST AVAILABLE COPY



CYANAMID

American Cyanamid Company
Agricultural Products Division
Crop Protection Chemicals Department
Wayne, NJ 07470 ©1994

50859

EPA Reg. No. 241-350

FOR POSTEMERGENCE USE IN PINTO TYPE DRY BEANS

**FOR DISTRIBUTION AND USE ONLY IN COLORADO, KANSAS,
WYOMING, NEBRASKA, AND TEXAS**

THIS LABEL EXPIRES APRIL 30, 1996

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN
THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

GENERAL INFORMATION

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT DG application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application.

DO NOT apply PURSUIT DG postemergence after crop has begun to flower or crop injury may result.

Some varieties of pinto beans may be inherently less tolerant to PURSUIT DG applications and crop injury could result. Contact the seed company for information and recommendations about postemergence applications of PURSUIT DG.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible

*Registered Trademark of American Cyanamid Company

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT DG in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, potatoes, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

Spray equipment used for PURSUIT DG application must be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for post applications.

DO NOT make more than one application of PURSUIT DG per year. **DO NOT** apply PURSUIT DG after July 15. Allow at least 60 days between application and harvest.

(see reverse side)

BEST AVAILABLE COPY

241-350

live label instructions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

APPLICATION RATE AND TIMING

Postemergence Applications. Apply PURSUIT DG at 1.08 ounces per acre. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture.

DO NOT use crop oils, petroleum oils, liquid fertilizers or Dash¹.

DO NOT APPLY PURSUIT DG POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRIFOLIATE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.

ECO-PAK CONVERSION TABLE

<u>Use Rate (oz/acre)</u>	<u>1 ECO-PAK Soluble Bag Treats</u>
1.08	6-2/3 acres

WEEDS CONTROLLED

PURSUIT DG applied at the broadcast rate of 1.08 ounces per acre early postemergence will control:

Mustard, wild	Nightshade, hairy
Nightshade, black	Pigweed, redroot
Nightshade, Eastern black	

Postemergence applications of 1.08 ounces per acre must be to weeds less than 2 inches tall for best results.

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label:

- Anytime
 - Soybeans
 - Peanuts
 - Great Northern, red kidney, and pinto type dry beans
 - IMI-Corn

¹Dash is a trademark of BASF Corporation

PE-XXXX 12/594 SD

Rye
Wheat

- Nine and one-half months after PURSUIT DG application:
 - Field corn
 - Field corn grown for seed*
 - Tobacco
- Eighteen months after PURSUIT DG application:
 - Alfalfa
 - Cotton
 - Lettuce
 - Oats
 - Popcorn
 - Safflower
 - Sorghum
 - Sweet Corn
- Twenty-six months after PURSUIT DG application:
 - All crops not listed in this ROTATIONAL CROP GUIDELINE.
- Forty months after PURSUIT DG application:
 - Sugar beets
 - Red table beets

In the event of a crop loss Great Northern or red kidney beans may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

* Several seed companies have tested a wide range of in-breds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.

BEST AVAILABLE COPY



CYANAMID

American Cyanamid Company
Agricultural Products Division
Crop Protection Chemicals Department
Wayne, NJ 07470

©1994

EPA Reg. No. 241-350

FOR USE IN LIMA BEANS ONLY

**FOR DISTRIBUTION AND USE ONLY ON THE DELMARVA PENINSULA OF
MARYLAND, DELAWARE, AND VIRGINIA**

READ LABEL CAREFULLY

THIS LABEL EXPIRES APRIL 30, 1997

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN
THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

GENERAL INFORMATION

PURSUIT DG application methods and use rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT DG only to the legume vegetables specifically listed in this label. **DO NOT** apply PURSUIT DG in geographical regions outside those specifically defined in this label for each type of legume vegetable.

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT DG application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

BEST AVAILABLE COPY

*Registered Trademark of American Cyanamid Company

(see reverse side)

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT DG in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

Spray equipment used for PURSUIT DG application must be drained and thoroughly cleaned with water before being used to apply other products.

241-350

methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

DO NOT apply PURSUIT DG after June 30.

Allow at least 30 days between application and harvest.

Preplant Incorporated Applications. Apply PURSUIT DG at the broadcast rate of 0.72 ounces per acre to sand and loamy sand soils, or 1.08 ounces per acre to all other soils within 1 week before planting. Applied preplant incorporated, PURSUIT DG may be tank-mixed with a registered grass herbicide.

Preemergence Applications. Apply PURSUIT DG at the broadcast rate of 0.72 ounces per acre to sand and loamy sand soils, or 1.08 ounces per acre to all other soils immediately after, or up to 1 day after planting. PURSUIT DG may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide.

ECO-PAK CONVERSION TABLE

<u>Use Rate (oz/acre)</u>	<u>LECO-PAK Soluble Bag Treats</u>
0.72	10 acres
1.08	6-2/3 acres

WEEDS CONTROLLED

PURSUIT DG applied to low organic matter soils at the broadcast rate of 0.72 ounces per acre preplant incorporated or preemergence will control light to moderate infestations of:

Nightshade, black
Nightshade, Eastern black
Pigweed, redroot

PURSUIT DG applied at the broadcast rate of 1.08 ounces per acre preplant incorporated or preemergence will control:

Mustard, wild
Nightshade, black
Nightshade, Eastern black
Nightshade, hairy
Pigweed, redroot

Peanuts
IMI-Corn
Lima beans

2. Two months after PURSUIT DG application:
Snap beans
3. Four months after PURSUIT DG application:
Barley
Other types of edible beans and peas
Rye
Wheat
4. Nine and one-half months after PURSUIT DG application:
Field corn
Field corn grown for seed*
Tobacco
5. Eighteen months after PURSUIT DG application:
Alfalfa
Cotton
Lettuce
Oats
Popcorn
Safflower
Sorghum
Sweet Corn
6. Twenty-six months after PURSUIT DG application:
All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE.
7. Forty months after PURSUIT DG application:
Sugar beets
Red table beets

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

In the event of a crop loss, the bean or pea crop may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.

EPA Reg. No. 241-350

24 8-9

FOR USE IN ENGLISH PEAS ONLY

**FOR DISTRIBUTION AND USE ONLY ON THE DELMARVA PENINSULA OF
MARYLAND, DELAWARE, AND VIRGINIA**

READ LABEL CAREFULLY

THIS LABEL EXPIRES APRIL 30, 1996

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN
THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

GENERAL INFORMATION

PURSUIT DG application methods and use rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT DG only to the legume vegetables specifically listed in this label. **DO NOT** apply PURSUIT DG in geographical regions outside those specifically defined in this label for each type of legume vegetable.

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT DG application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

BEST AVAILABLE COPY

*Registered Trademark of American Cyanamid Company

(see reverse side)

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT DG in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

Spray equipment used for PURSUIT DG application must be drained and thoroughly cleaned with water before being used to apply other products.

RECORDED
241350

with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

DO NOT apply PURSUIT DG to sand or loamy sand soils.

Allow at least 30 days between application and harvest.

Preplant Incorporated Applications. Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre up to 1 week before planting. **DO NOT** apply PURSUIT DG to English peas that have been or will be treated with trifluralin.

Preemergence Applications. Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre immediately after, or up to 1 day after planting. **DO NOT** apply PURSUIT DG if English peas have been treated with trifluralin.

ECO-PAK CONVERSION TABLE

Use Rate (oz/acre)	1 ECO-PAK Soluble Bag Treats
1.08	6-2/3 acres

WEEDS CONTROLLED

PURSUIT DG applied at the broadcast rate of 1.08 ounces per acre preplant incorporated or preemergence will control:

Mustard, wild	Nightshade, hairy
Nightshade, black	Pigweed, redroot
Nightshade, Eastern black	

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label:

- Anytime
 - Soybeans
 - Peanuts
 - IMI-Corn
 - English peas and lima beans
- Two months after PURSUIT DG application:
 - Snap beans

- Wheat
- Nine and one-half months after PURSUIT DG application:
 - Field corn
 - Field corn grown for seed*
 - Tobacco
 - Eighteen months after PURSUIT DG application:
 - Alfalfa
 - Cotton
 - Lettuce
 - Oats
 - Popcorn
 - Safflower
 - Sorghum
 - Sweet Corn
 - Twenty-six months after PURSUIT DG application:
 - All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE.
 - Forty months after PURSUIT DG application:
 - Sugar beets
 - Red table beets

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

In the event of a crop loss, the bean or pea crop may be replanted. **DO NOT** feed treated forage, hay, or straw to livestock.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.

BEST AVAILABLE COPY



CYANAMID

American Cyanamid Company
Agricultural Products Division
Crop Protection Chemicals Department
Wayne, NJ 07470 ©1994

EPA Reg. No. 241-350

**FOR USE IN RED KIDNEY TYPE DRY BEANS, WHITE LUPINS,
LIMA BEANS, AND ENGLISH PEAS ONLY**

FOR DISTRIBUTION AND USE ONLY IN ILLINOIS, IOWA, AND WISCONSIN

READ LABEL CAREFULLY - USE INSTRUCTIONS VARY FOR EACH GROUP

THIS LABEL EXPIRES APRIL 30, 1997

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN
THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

GENERAL INFORMATION

PURSUIT DG application methods and use rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT DG only to the legume vegetables specifically listed in this label. DO NOT apply PURSUIT DG in geographical regions outside those specifically defined in this label for each type of legume vegetable.

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT DG application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. DO NOT apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application. DO NOT apply PURSUIT DG postemergence after crop has begun to flower or crop injury may result.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

*Dash is a trademark of BASF Corporation
*Registered Trademark of American Cyanamid Company

(see reverse side)

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT DG in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

Spray equipment used for PURSUIT DG application must be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for post applications.

DO NOT use crop oils, petroleum oils, liquid fertilizers or Dash!

DO NOT make more than one application of PURSUIT DG per year.

BEST AVAILABLE COPY

241-350

RED KIDNEY TYPE DRY BEANS, WHITE LUPINS AND LIMA BEANS

DO NOT apply PURSUIT DG after June 30.

DO NOT apply PURSUIT DG to white lupins grown on sand or loamy sand soils.

Allow at least 30 days between application and harvest of lima beans.

Allow at least 60 days between application and harvest of red kidney beans or white lupins.

Preplant Incorporated Applications. Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre within 1 week before planting. Applied preplant incorporated, PURSUIT DG may be tank-mixed with a registered grass herbicide.

Preemergence Applications. Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre immediately after, or up to 1 day after planting. PURSUIT DG may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide.

ENGLISH PEAS

Allow at least 30 days between application and harvest.

Preplant Incorporated Applications. Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre up to 1 week before planting. DO NOT apply PURSUIT DG to English peas that have been or will be treated with trifluralin.

Preemergence Applications. Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre immediately after, or up to 1 day after planting. DO NOT apply PURSUIT DG if English peas have been treated with trifluralin.

Postemergence Applications. Apply PURSUIT DG at 1.08 ounces per acre. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture.

DO NOT APPLY PURSUIT DG POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRUE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT. For optimum weed control and crop safety, apply PURSUIT DG prior to 5 nodes before flowering. If English peas have been treated with trifluralin, PURSUIT DG may be applied ONLY at 0.72 ounces per acre for suppression of black nightshade. The use of trifluralin prior to PURSUIT DG application may increase the likelihood and severity of crop injury.

ECO-PAK CONVERSION TABLE

Use Rate (oz/acre)
1.08

LECO-PAK Soluble Bag Treats
6-2/3 acres

PF XXXXX 12/4/94 SD



CYANAMID
American Cyanamid Company
Agricultural Products Division
Crop Protection Chemicals Department
Wayne, NJ 07470 ©1994

Nightshade, black
Nightshade, Eastern black

Postemergence applications of 1.08 ounces per acre must be to weeds less than 2 inches tall for best results.

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label:

1. Anytime
Soybeans
Peanuts
Edible bean and pea types listed on this label (Refer to heading on page 1).
IMI-Corn
2. Four months after PURSUIT DG application:
Barley
Edible bean and pea types not listed on this label (Refer to heading on page 1).
Rye
Wheat
3. Nine and one-half months after PURSUIT DG application:
Field corn
Field corn grown for seed*
Tobacco
4. Eighteen months after PURSUIT DG application:
Alfalfa Oats Sorghum
Cotton Popcorn Sweet Corn
Lettuce Safflower
5. Twenty-six months after PURSUIT DG application:
All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE.
6. Forty months after PURSUIT DG application:
Sugar beets Red table beets

In the event of a crop loss, red kidney beans, white lupins, lima beans, or English peas may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.

BEST AVAILABLE COPY

EPA Reg. No. 241-350

29229

**FOR USE IN NAVY, GREAT NORTHERN, RED KIDNEY, BLACK TURTLE,
COCOA CRANBERRY, AND SMALL WHITE TYPE DRY BEANS ONLY**

**FOR DISTRIBUTION AND USE IN MICHIGAN ONLY
(EXCLUDING UPPER PENINSULA)**

241-350
EPA Reg. No.

THIS LABEL EXPIRES APRIL 30, 1996

OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.

GENERAL INFORMATION

PURSUIT DG application methods and use rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT DG only to the legume vegetables specifically listed in this label. **DO NOT** apply PURSUIT DG in geographical regions outside those specifically defined in this label for each type of legume vegetable.

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT DG application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application.

DO NOT apply PURSUIT DG postemergence after crop has begun to flower or crop injury may result.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

*Dash is a trademark of BASF Corporation
*Registered Trademark of American Cyanamid Company

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT DG in any manner not specifically described in this label

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

Spray equipment used for PURSUIT DG application must be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for postemergence applications. **DO NOT** use crop oils, petroleum oils, liquid fertilizers or Dash¹.

DO NOT make more than one application of PURSUIT DG per year.

Refer to the PURSUIT DG leaflet label for approved combination treatments. When PURSUIT DG is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

(See reverse side)

BEST AVAILABLE COPY

Allow at least 60 days between application and harvest.

Preplant Incorporated Applications. Apply PURSUIT DG within 1 week before planting. Applied preplant incorporated, PURSUIT DG may be tank-mixed with a registered grass herbicide. Refer to the following table for use rate.

Soil Type	Use Rate (oz/acre)
Sands, loamy sands	DO NOT USE
Sandy loams and all other soils with \leq 2% O.M.	0.72
All other soils with $>$ 2% O.M.	1.08

Preemergence Applications. Apply PURSUIT DG at the rates shown in the preceding table immediately after, or up to 1 day after planting. PURSUIT DG may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide.

Postemergence Applications. Apply PURSUIT DG at 0.72 ounces per acre. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture.

DO NOT APPLY PURSUIT DG POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRUE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.

ECO-PAK CONVERSION TABLE

Use Rate (oz/acre)	1 ECO-PAK Soluble Bag Treats
0.72	10 acres
1.08	6-2/3 acres

WEEDS CONTROLLED

PURSUIT DG applied to low organic matter soils at the broadcast rate of 0.72 ounces per acre preplant incorporated, preemergence, or postemergence will control light to moderate infestations of:

Nightshade, black
Nightshade, Eastern black
Pigweed, redroot

PURSUIT DG applied at the broadcast rate of 1.08 ounces per acre preplant incorporated or preemergence will control:

Mustard, wild
Nightshade, black
Nightshade, Eastern black
Nightshade, hairy
Pigweed, redroot

Soybeans

Peanuts

IMI-Corn

Edible bean and pea types listed on this label (Refer to heading on page 1)

- Four months after PURSUIT DG application:
Barley
Edible bean and pea types not listed on this label (Refer to heading on page 1)
Rye
Wheat
- Nine and one-half months after PURSUIT DG application:
Field corn
Field corn grown for seed*
Tobacco
- Eighteen months after PURSUIT DG application:
Alfalfa
Cotton
Lettuce
Oats
Popcorn
Safflower
Sorghum
Sweet Corn
- Twenty-six months after PURSUIT DG application:
All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE.
- Forty months after PURSUIT DG application:
Sugar beets
Red table beets

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.

 **CYANAMID**
American Cyanamid Company
Agricultural Products Division
Crop Protection Chemicals Department
Wayne, NJ 07470 ©1994