

TRIAD

DISINFECTANT-CLEANER

DIRECTIONS FOR EFFICIENT AND ECONOMICAL USE AS A DISINFECTANT • CLEANER • FUNGICIDE • DEODORIZER

GENERAL INDUSTRIAL AND HOUSEHOLD PURPOSES

Dilute 2 FL OZ per gallon of water (1 to 64) for DISINFECTION AND FUNGICIDAL activity against Pathogenic Fungi. NOTE: Pre-clean heavy soil or gross filth deposits with a dilute solution of Triad prior to application at the noted dilution to ensure proper disinfectant action.

FLOORS: Apply with a clean cloth, sponge, mop or sprayer. Rub or provide mechanical agitation to loosen soil. Allow solution to remain in contact with surface for several minutes, then take up with a sponge, mop or vacuum pickup. If floor polish is to be applied, rinse with clear cool water, otherwise rinsing is not needed.

COUNTER TOPS, WALES: Apply with a clean cloth or sponge, rubbing to loosen soil. Wipe off and let dry. No rinsing is needed.

HOSPITALS, CLINICS, NURSING HOMES, PUBLIC AREAS: Dilute 3 FL OZ per gallon of water (3 to 128) for DISINFECTION against PSEUDOMONAS AERUGINOSA and FUNGICIDAL ACTIVITY against PATHOGENIC FUNGI. Follow application directions as noted under General Industrial and Household Purposes.

As a SANITIZER: For all hard surfaces, food processing equipment, food utensils. Dilute 2 FL OZ per 2½ gallons of water (1 to 160). This dilution provides 200 ppm of active quaternary per gallon of water.


A POTABLE WATER RINSE IS REQUIRED BEFORE REUSE OF EQUIPMENT: The product, at this concentration, fulfills the criteria of APPENDIX F of the GRADE "A" PASTEURIZED MILK ORDINANCE 1965 RECOMMENDATION of the U.S. PUBLIC HEALTH SERVICE in water up to 450 ppm of hardness calculated as CaCO₃, when tested by the AOAC GERMICIDE and DETERGENT SANITIZERS OFFICIAL METHOD.

*NOTE: Rinse empty containers thoroughly with water and discard it.

This product authorized for use under USDA Consumer and Marketing Service Inspection Program, for Poultry, Meat, Rabbit and Egg Product Plants. However, the sanitizing directions on appropriate panel must be followed, including a potable water rinse, and all food and packaging equipment must be first removed from the room or carefully protected.

DISINFECTANT CLEANER DEODORIZER FUNGICIDE SANITIZER

- For floors, walls, and hard surfaces in HOSPITALS, NURSING HOMES, HOTELS, INSTITUTIONS, PUBLIC AREAS, INDUSTRIAL AREAS, AND CONVEYANCES
- Excellent for OPERATING ROOMS, LABORATORIES, REST ROOMS, CLINICS, ISOLATION WARDS, NURSERIES and FOOD SERVICE AREAS
- Effective in SCHOOLS, rest rooms, locker rooms, garbage pails, sink tops, cafeterias, corridors, class rooms and offices.
- Excellent cleaner for restaurants, doctor and dentist offices, sewage disposal plants, barber and beauty shops, swimming pool surfaces, bakeries and breweries.
- TRIAD DISINFECTANT-CLEANER passes the AOAC Use Dilution Test against *Staphylococcus Aureus*, *Salmonella Choleraesuis*, at 2 FL OZ per gallon of water. Also effective against *Pseudomonas Aeruginosa* at 3 FL OZ per gallon of water. Fungicidal against pathogenic fungus - *Trichophyton Interdigitale* at 2 FL OZ per gallon of water.



ACTIVE INGREDIENTS	6.2%
Alkyl (60% C14, 30% C16, 5% C12, 5% C18) dimethyl benzyl ammonium chlorides	1.6%
Alkyl (68% C12, 32% C14) dimethyl ethylbenzyl ammonium chlorides	1.6%
Sodium carbonate	3.0%
INFRT INGREDIENTS	93.8%
Includes efficient cleaning detergents, builders, sequestering agents, dye, odorants and water.	

DANGER:

KEEP OUT OF REACH OF CHILDREN.

SEE FIRST AID & PRECAUTIONS ON BACK PANEL

Net Contents Liquid Gallons

DANGER. KEEP OUT OF REACH OF CHILDREN.
Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Protect eyes and skin when handling concentrate. Harmful if吞 swallowed. Avoid contamination of food. As an breathing spray mist. This product is toxic to fish. Do not contaminate water by the cleaning of equipment or discharge of waste.

FIRST AID: In case of contact immediately flush eyes or skin with plenty of water for at least 15 minutes. If in eyes call a physician. Remove and wash contaminated clothing before reuse. If swallowed, drink PROMPTLY a large quantity of milk, egg whites, dilute solution or if these are not available drink large quantities of water. Avoid aspiration. Call a physician immediately.

NOTE TO PHYSICIAN: Product causing damage may contraindicate the use of antacids, laxatives. Measures against circulatory shock, respiratory depression, and convulsions may be needed.

EPA Reg. No. 257-296

EPA Est. 257-ME-1

(f) **FULD-STALFORT, INC.**
HAVRE de GRACE, MARYLAND 21078