



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OCT 14 2010

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

Mr. Mike See
Bayer Cropscience
2 T.W. Alexander Drive, PO Box 12014
Research Triangle Park, NC 27709

Subject: SC 450 Herbicide
EPA Registration Number 264-1061
Submission dated August 11, 2010

Dear Mr. See:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act as amended is acceptable, provided you make the following changes before you release the product for shipment.

1. Add an appropriate EPA Establishment Number to the label
2. Remove "Wash thoroughly with soap and water after handling" from the Precautionary Statements section
3. Revise the heading "GENERAL INFORMATION" to "USE INFORMATION" or "PRODUCT INFORMATION"
4. On page 3, in the section Use Restrictions and Precautions, add the following restriction: "A 25 foot buffer for ground applications, or a 200 foot buffer for aerial applications, must be maintained between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (including grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrub lands), sensitive freshwater habitats (including lakes, rivers, sloughs, ponds, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats."
5. On page 10, remove the statement "The applicator should be familiar with and take into account the information covered in the Aerial Drift Reduction Advisory Information"
6. On page 10, add "you must" before "maintain a 25 foot buffer for ground applications, or a 200 foot buffer for aerial applications"
7. On page 10, revise "such as" to "including" in the buffer statement

Page 2


EPA Registration Number 264-1061

8. On page 10, revise "should" to "must" in the section Wind

Submit one (1) copy of final printed labeling incorporating the above changes before you release the product for shipment. Amended labeling will supercede all previously accepted ones. A stamped copy of labeling is enclosed for your records. If you have any questions, please contact Hope Johnson at 703-305-5410.

NOTE: The Corvus Herbicide label (EPA Registration Number 264-1066) and Capreno Herbicide label (EPA Registration Number 264-1063) referenced in this application must be revised to include a mandatory buffer statement per #4 above in the Restrictions section. Submit a label amendment for these products within 30 days from the date of this letter.

Sincerely,



James A. Tompkins
Product Manager 25
Herbicide Branch
Registration Division (7505P)

ACCEPTED
with COMMENTS
in EPA L. Dated
OCT 14 2010

OCT 14 2010
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No. 264-1061

GROUP	2	HERBICIDE
-------	---	-----------

SC 450 HERBICIDE

For selective control of weeds in corn

ACTIVE INGREDIENT:

THIENCARBAZONE-METHYL* (Methyl 4-[[[(4,5-dihydro-3-methoxy-4-methyl-5-oxo-1H-1,2,4-triazol-1-yl)carbonyl]amino]sulfonyl]-5-methyl-3-thiophenecarboxylate)..... 19.2

OTHER INGREDIENTS:..... 80.8

TOTAL:..... 100.0%

*SC 450 Herbicide is formulated as a suspension concentrate of THIENCARBAZONE-METHYL at a concentration of 1.88 pounds of active ingredient per gallon.

E.P.A. Reg. No. 264-1061

E.P.A. Est. No.

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.
For PRODUCT USE Information Call 1-800-331-2867.

FIRST AID

IF SWALLOWED:	<ul style="list-style-type: none"> • Immediately call a poison control center or doctor for treatment advice. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Have person sip a glass of water if able to swallow. • Do not give anything by mouth to an unconscious person.
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a physician if irritation persists.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

NOTE TO PHYSICIAN: No specific antidote is available. All treatments should be based on observed signs and symptoms of distress in the patient. Overexposure to materials other than this product may have occurred.

PRECAUTIONARY STATEMENTS

CAUTION

HAZARD TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed or absorbed through the skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some of the materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

Applicators and other handlers must wear: Long-sleeved shirt and long pants, chemical-resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride, shoes plus socks and protective eye wear. When mixing/loading or cleaning equipment, wear a chemical resistant apron in addition to the other required PPE. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for

cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering control statement:

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [(40 CFR §170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Users should remove clothing/PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinseate.

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having "high potential for reaching surface water via runoff", according to the pesticide's "mean" soil partition coefficient (Kd) for several days after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and natural surface water features such as ponds, streams and springs will reduce potential loading of Thien carbazon-methyl from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

Ground Water Advisory

This chemical has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.

ENDANGERED SPECIES PROTECTION REQUIREMENTS:

This product may have effects on federally listed threatened or endangered species or their critical habitat in some locations. When using this product, you must follow the measures contained in the Endangered Species Protection Bulletin for the county or parish in which you are applying the pesticide. To determine whether your county or parish has a Bulletin, and to obtain that Bulletin, consult <http://www.epa.gov/espp/>, or call 1-800-447-3813 no more than 6 months before using this product. Applicators must use Bulletins that are in effect in the month in which the pesticide will be applied. New Bulletins will generally be available from the above sources 6 months prior to their effective dates.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read entire label before using this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the same area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticides.

Do not drain or rinse equipment near desirable vegetation.

Avoid spray drift from treated areas. Refer to the Spray Drift Management section of this label for additional information.

Non-target plants may be adversely affected if the pesticide is allowed to drift from areas of application. To prevent damage to crops and other desirable plants, read and follow all directions and precautions on this label before using.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil or water, is coveralls over long-sleeved shirt and long pants, socks and shoes and chemical-resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE

Store unused product in original container only, out of reach of children and animals. NEVER TRANSFER THIS PRODUCT TO ANOTHER CONTAINER FOR STORAGE.

PESTICIDE DISPOSAL

Dispose wastes resulting from the use of this product on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then puncture and dispose of in a sanitary landfill, or by incineration; or, if allowed by State and local authorities, by burning. If burned, stay out of smoke

GENERAL INFORMATION

SC 450 Herbicide is a selective herbicide for control of important broadleaf and grassy weeds infesting field corn, sweet corn and popcorn. SC 450 Herbicide can be used in corn grown for silage, grain or seed. SC 450 Herbicide can be applied for post harvest, preplant (surface-applied or incorporated), preemergence, or postemergence uses.

SC 450 Herbicide is effective in controlling triazine resistant populations of weed species which are listed in the "BROADLEAF AND GRASS WEEDS CONTROLLED BY SC 450 HERBICIDE" tables in this label.

Corn hybrids and certain male pollenators within blended corn varieties vary in their response to SC 450 Herbicide. Not all hybrids or male pollenators within blended corn varieties have been tested for sensitivity to SC 450 Herbicide. You should consult with your seed provider, your local Bayer CropScience representative and/or other knowledgeable agricultural professionals for advice on tolerance of hybrids or varieties containing male pollinator lines before applying SC 450 Herbicide. If the tolerance of a hybrid or variety containing male pollinator lines is not known, you should apply SC 450 Herbicide to a small area to first determine if the hybrid is tolerant prior to spraying large acreages of that hybrid.

SC 450 Herbicide also contains a safener, which greatly reduces or prevents the temporary yellowing or stunting crop response associated with this herbicide chemistry. If symptoms appear, corn quickly outgrows the effect and develops normally.

USE RESTRICTIONS AND PRECAUTIONS

- Plant corn at least 1½ inches deep. Corn seed must be completely covered with soil and furrow firmed.
- Do not apply this product through any type of irrigation system.
- Do not apply more than 2.73 fl oz of SC 450 Herbicide per 365 day period or exceed the maximum labeled rate for any given soil type in a single application.
- Do not apply more than 0.04 pounds per acre of THIENCARBAZONE-METHYL active ingredient from all sources per 365 day period, or exceed the maximum labeled rate for any given soil type in a single application.
- SC 450 Herbicide also contains 1.88 pounds of the safener Cyprosulfamide per gallon. Do not apply more than 0.20 pounds per acre of CYPROSULFAMIDE safener from all sources per 365 day period, or exceed 0.14 pounds per acre of CYPROSULFAMIDE safener in a single application.
- DO NOT apply SC 450 Herbicide postemergence beyond the 2-leaf collar stage (V2) to corn with liquid fertilizers as the primary spray carrier. Only apply postemergence to corn with water as the primary spray carrier plus recommended adjuvants. See spray additives section.
- Apply SC 450 Herbicide spray mixtures within 24 hours of mixing to avoid degradation of this product.
- Avoid spray drift from treated areas. Refer to the Spray Drift Management section of this label for additional information.
- DO NOT apply more than two applications of SC 450 Herbicide to a corn crop in one growing season from 30 days before planting through the V12 growth stage.
- Do not apply more than one application of SC 450 Herbicide during the post harvest application period (from harvest of preceding crop to 30 days prior to planting of next crop).
- Allow at least 14 days between applications of SC 450 Herbicide.
- When weeds are stressed due to drought, heat, lack of fertility, flooding, or prolonged cool temperatures control can be reduced or delayed because weeds are not actively growing. To obtain optimum weed control with an application of SC 450 Herbicide follow label directions and use when weeds are actively growing.

- Weed control may be reduced if the application is made when weeds are dust covered or in the presence of heavy dew, fog, and mist/rain or when weeds are under stress due to drought.
- Tank contamination can cause severe damage to other crops. Careful management of tank cleanout is required.
- DO NOT graze livestock, or harvest corn forage or sweet corn ears within 45 days of application of SC 450 Herbicide.

ROTATIONAL CROP RESTRICTIONS

Rotational crops vary in their crop response to low concentrations of SC 450 Herbicide remaining in the soil. The amount of SC 450 Herbicide that may be present in the soil depends on soil moisture, soil temp, application rate, elapsed time since application and other environmental factors. When SC 450 Herbicide is used in combination with other products, always follow the most restrictive rotational crop requirements.

The active ingredient, thien carbazone-methyl is used in multiple products. When determining rotational crop intervals the total application rate of thien carbazone-methyl from all sources should be included on an equivalent basis to SC 450 Herbicide rates when determining crop rotation intervals from Tables 1 and 2.

The following rotational crops may be planted after applying SC 450 Herbicide:

Table 1. Minimum recropping intervals for various crops following SC 450 Herbicide application at rates totaling greater than 0.93 fl oz of product per acre.

Rotational Interval (elapsed time)	Crop	Minimum precipitation requirement ¹
0 Months ²	Yellow field corn	None
3 Months ²	Wheat	None
9 Months ²	Barley, Cotton, Soybean ³ , Sweet corn ³ , Popcorn ³ , White field corn ³	15 inches of cumulative precipitation from application to planting of rotational crop
17 Months ³	Alfalfa, Green and Dry Beans, Sorghum, Oats, Sunflower, Canola, Potato, Sugar beet and All other crops	30 inches of cumulative precipitation from application to planting of rotational crop

¹The amount of cumulative precipitation required before planting a rotational crop is in addition to the required rotational interval given in months. Furrow or flood irrigation not to be included in total. No more than 7 inches of overhead irrigation included in total.

²Crop varieties planted back at intervals of one year or less should not have known acute sensitivity to ALS-inhibiting and/or SU herbicides.

³When soil pH is 7.5 or above crop plant back should be delayed to the next interval, and to 24 months for crops listed in the 17 month interval above.

Table 2. Minimum recropping intervals for various crops following SC 450 Herbicide application at rates totaling greater than 0.93 fl oz of product per acre or less.

Rotational Interval (elapsed time)	Crop	Minimum precipitation requirement ¹
0 Months ²	Yellow Field Corn	None
2 Months ²	Soybean ³	None
3 Months ²	Wheat	None
9 Months ²	Barley, Cotton, Sorghum, Sweet corn ⁴ , Popcorn ⁴ , White field corn ⁴	15 inches of cumulative precipitation from application to planting of rotational crop
17 Months ⁴	Alfalfa, Green and Dry Beans, Oats, Sunflower, Canola, Potato, Sugar beet and All other crops	30 inches of cumulative precipitation from application to planting of rotational crop

¹The amount of cumulative precipitation required before planting a rotational crop is in addition to the required rotational interval given in months. Furrow or flood irrigation not to be included in total. No more than 7 inches of overhead irrigation included in total.

²Crop varieties planted back at intervals of one year or less should not have known acute sensitivity to ALS-inhibiting and/or SU herbicides.

³When soil pH is 7.5 or above soybean plant back should be delayed to the 9 month interval.

⁴When soil pH is 7.5 or above crop plant back should be delayed to the next interval, and to 24 months for crops listed in the 17 month interval above.

INSECTICIDE INTERACTION INFORMATION

Soil applied insecticide interaction information

When SC 450 Herbicide and certain organophosphate (OP) or carbamate insecticides are applied to corn injury can result. DO NOT USE SC 450 Herbicide in the same season as Lorsban®, Counter® 15G, Counter® 20G, Dyfonate®, or Thimet®.

For all corn hybrids, the following table describes the uses of soil-applied insecticides prior to an application of SC 450 Herbicide:

Soil Applied Insecticide	Use Pattern	Use of SC 450 Herbicide in the Same Season
Aztec [®] , Regent [®] , Tefluthrin (e.g. Force [®])	All	No use precautions
Chlorpyrifos (e.g. Lorsban [®] 15G), Terbufos (e.g. Counter [®] 15G, Counter [®] 20CR), Phorate (e.g. Thimet [®]), Fonophos (e.g. Dyfonate [®])	All	Do Not Use

Foliar Insecticide Interaction Information

Foliar applications of an organophosphate or carbamate insecticide should not be made within 7 days of an application of SC 450 Herbicide or crop injury may result.

MIXING INSTRUCTIONS

Application with water or liquid fertilizer (prior to V-2 stage ONLY for liquid fertilizer) as a carrier: Fill the spray tank 1/4 to 1/2 of the required volume of water or liquid fertilizer prior to the addition of SC 450 Herbicide and begin agitation. Slowly add the proper amount of SC 450 Herbicide, and begin bringing the water or liquid fertilizer to the desired level. Maintain sufficient agitation to ensure a uniform spray mixture during application. If SC 450 Herbicide is applied in a tank mixture with other pesticides, add SC 450 Herbicide to the spray tank first and ensure it is thoroughly dispersed before adding other pesticides. If ammonium sulfate (AMS) is the nitrogen fertilizer source, it is preferred that the AMS go into the tank after the SC 450 Herbicide and before other pesticide tank mix partners. If UAN is the nitrogen fertilizer source, it should be added to the tank after other pesticide tank mix partners. Add surfactants to the tank last. Continue to fill the tank with carrier to the desired volume while agitating. **CONTINUE AGITATION DURING APPLICATION TO ENSURE A UNIFORM SPRAY MIXTURE.**

Re-suspending SC Products in Spray Solution: Like other suspension concentrates (SC's), SC 450 Herbicide will settle if left standing without agitation. If the spray solution is allowed to settle for one hour or more, reagitate the spray solution for a minimum of 10 minutes before application.

TANK CLEANUP PROCEDURE

Sprayer Cleanup: To avoid injury or exposure to non-target crops, thoroughly clean all mixing and spray equipment, including pumps, nozzles, lines and screens with a good quality tank cleaner, on approved rinse pad or on the field site where an approved crop is to be grown.

Cleaning Equipment After SC 450 Herbicide Application

Special attention must be given to cleaning equipment before spraying a crop other than corn. Mix only as much cleaning solution as needed.

1. Flush tank, hoses, boom and nozzles with clean water.
2. Use a pressure washer with a high quality commercial spray tank cleaner in water to clean the inside of the spray tank. Take care to wash all parts of the tank, including the inside top surface. If a pressure washer is not available, completely fill the sprayer with the cleaning solution to ensure contact of the cleaning solution with all internal surfaces of the tank and plumbing. Start agitation in the sprayer and thoroughly recirculate the cleaning solution for at least 15 minutes. All visible deposits must be removed from the spraying system.
3. Flush hoses, spray lines, and nozzles for at least 1 minute with the cleaning solution.
4. Dispose of rinseate from steps 1-3 in an appropriate manner.
5. Repeat steps 2-4.
6. Remove nozzles, screens and strainers and clean separately in the cleaning solution after completing the above procedures.
7. Rinse the complete spraying system with clean water.
8. Cleanup should be conducted on an approved rinse pad or the field site where an approved crop is to be grown.

TANK MIXTURES

SC 450 Herbicide can be applied in tank mixture with many other pesticides registered for use on approved crops. Refer to "Tank Mix Combination" section for application rates and other restrictions.

COMPATIBILITY

If SC 450 Herbicide is to be tank mixed with liquid fertilizers or other pesticides, compatibility should be tested prior to mixing. To test for compatibility, use a small container and mix a small amount (0.5 to 1 qt) of spray, combining all ingredients in the same ratio as the anticipated use. If any indications of physical incompatibility develop, do not use this mixture for spraying. Indications of incompatibility usually will appear within 5-15 minutes after mixing. Read and follow all parts of the label of each tank-mix partner.

APPLICATION PROCEDURES

APPLICATION TIMING

SC 450 Herbicide may be:

- Used in either conventional, conservation tillage, or no-till crop management systems.
- Applied either preplant, preplant incorporated (less than 2" deep), preemergence or early postemergence for use in corn production.
- Applied postemergence to corn.
- Used during the period from harvest of preceding crop to 30 days prior to planting of next crop.

SC 450 Herbicide treatments applied to the soil surface are most effective in controlling weeds when adequate rainfall is received within 14 days after application. If cultivation is necessary because of soil crusting, soil compaction or weed germination before rain occurs, use shallow tillage such as rotary hoe to lightly incorporate SC 450 Herbicide and make certain corn seeds are below the tilled area. If treated soil is moved during tillage practices in such a way that the herbicide barrier is no longer intact, weeds may emerge from areas where treated soil has been removed. Do not incorporate with a drag harrow after planting.

Preplant Surface-Applied: SC 450 Herbicide may be applied up to 21 days before planting corn. SC 450 Herbicide may be applied up to 30 days prior to planting when used in a planned sequential application program such as SC 450 Herbicide followed by Ignite[®] 280 SL, Buctril[®] or other post applied herbicides. Refer to the label of the respective sequential partner for specific use directions. Moving treated soil out of the row or moving untreated soil to the surface during planting may result in reduced weed control.

Preplant Incorporated: SC 450 Herbicide may be applied up to 21 days before planting corn. SC 450 Herbicide may be applied up to 30 days prior to planting when used in a planned sequential application program such as SC 450 Herbicide followed by Ignite[®] 280 SL, Buctril[®] or other post applied herbicides. Refer to the label of the respective sequential partner for specific use directions. Apply to the soil and uniformly incorporate in the top two inches of soil before planting using a finishing disc harrow, field cultivator or similar implement capable of providing uniform two inch incorporation. Do not incorporate SC 450 Herbicide deeper than 2" or weed control may be reduced.

Preplant/Preemergence Burndown: Prior to corn emergence, when weeds are present at the time of treatment, SC 450 Herbicide plus adjuvants (see spray additives section on this label) is recommended for burndown of labeled weeds 6" or less in height. Weed growth ceases within hours after SC 450 Herbicide is applied. Weed death generally occurs within 7 to 14 days after application but is progressively slower as weeds approach the 6-inch height. For preplant and preemergence burndown methylated seed oil (MSO) can be substituted for crop oil concentrate (COC) as an adjuvant. When weeds are greater than 6" in height, faster weed kill is desired, or weeds not controlled by SC 450 Herbicide are present the addition of a burndown herbicide (e.g., Gramoxone[®] Extra, Glyphosate, or 2,4-D) is recommended. Follow directions for use and precautions and restrictions on the label of any tank-mix partner included for the burndown.

Preemergence: Apply SC 450 Herbicide during planting (behind the planter after furrow closure) or after planting, but before weeds or crop emerge. Failure to thoroughly close and firm the seed furrow may allow herbicide to directly contact the seed which can cause injury.

Early Postemergence: SC 450 Herbicide alone and/or in certain tank mixtures may be applied to corn from spiking through the 2-leaf collar growth stage (V2, the first leaf has a rounded tip) at rates as defined in Table 1 under the Specific Uses Directions section of this label.

Postemergence:

Broadcast: SC 450 Herbicide alone and/or in certain tank mixtures may be applied broadcast to corn from spiking through the 6-leaf collar growth stage (V6, the first leaf has a rounded tip) at rates as defined under the Postemergence Rates heading of the Specific Use Directions section of this label. Do not make broadcast postemergence applications beyond the V6 growth stage.

Directed: Applications of SC 450 Herbicide may be applied when corn is greater than V6 and less than V12 stage of growth as a directed postemergence spray. Drop Nozzles must be used for applications of SC 450 Herbicide after the V6 stage of growth. Do not apply to corn that is more mature than V12 (i.e. more than 12 visible leaf collars) stage of growth. Applications of SC 450 Herbicide on corn that is V6 to V12 increases the potential for crop response. The risk may be greatly reduced, but not eliminated, by using drop nozzles properly placed between corn rows to optimize coverage on the weeds and minimize spray contact in the whorl and the leaf axles of the corn stalks. Use drop nozzles and appropriate spacing to direct spray below the corn whorl and upper leaves. The top of the target weed canopy must be sufficiently below the whorl and upper leaves of the crop to permit this application and provide adequate spray coverage. The height differential required between the crop and weed canopy will depend on the specific equipment used.

DO NOT graze livestock, or harvest corn forage or sweet corn ears within 45 days of application of SC 450 Herbicide.

Post-Harvest: SC 450 Herbicide may be used alone or in combination with other registered herbicides as a fall to early spring-applied ground broadcast application for burn down and residual control of certain winter annual broadleaf and other weeds listed on this label.

Apply SC 450 Herbicide to unfrozen ground from harvest of preceding crop to 30 days prior to planting of next crop. Apply SC 450 Herbicide beginning at the first sign of germination of any of the winter annual broadleaf weeds listed on this label. This application will reduce weed cover before spring planting. A fall application of SC 450 Herbicide is not intended to provide weed control throughout the

succeeding cropping period, but is part of a weed management program to remove weed covers that rob residual moisture and nutrients from the soil, slow soil warming, and hamper field preparation for and planting of succeeding crops. Weed control in succeeding crops will require additional applications of residual and/or postemergence herbicides for season-long control.

Apply 0.91 to 2.25 fl oz of SC 450 Herbicide plus adjuvants (see spray additives section on this label) for burndown of labeled weeds 6" or less in height. Weed growth ceases within hours after SC 450 Herbicide is applied. Weed death generally occurs within 7 to 14 days after application but is progressively slower as weeds approach the 6-inch height. For post-harvest applications, methylated seed oil (MSO) can be substituted for crop oil concentrate (COC) as an adjuvant. For optimum control, weeds must be in vegetative stages of growth. The length of residual control will increase with the application rate of SC 450 Herbicide.

If emerged weeds are present and are greater than 6 inches in height or diameter, use 2,4-D or an appropriate alternative postemergence herbicide in the tank mixture.

Control of established common dandelion requires a tank mixture containing at least 1 pint/acre (4-pound/gallon) of 2,4-D, and/or a tank-mixture with AUTUMN™ Herbicide.

Field corn, soybean and other crops can only be planted at the intervals specified in the "Rotational Crop Restrictions" section of this label after a post-harvest application of SC 450 Herbicide.

When using SC 450 Herbicide in tank mix combinations, follow the entire label of the most restrictive tank mix partner label.

SPRAY ADDITIVES

SC 450 Herbicide is a suspension concentrate that requires the use of an external adjuvant and a nitrogen fertilizer source to achieve optimum weed control.

Crop Oil Concentrate

Use Crop Oil concentrate (COC) at 1 gallon per 100 gallons of water (1% v/v), with a minimum of 1.25 pt/A. COC should contain at least 80% crop oil and 10% emulsifier or greater. The use of adjuvants such as non-ionic surfactants or refined vegetable oils will result in unacceptable or erratic weed control.

Ammonium Nitrogen Fertilizer

Use 1.5 qt/A of a high-quality urea ammonium nitrate (UAN) or 1.5 lb/A minimum (or 8.5 lb per 100 gallons) of a spray-grade ammonium sulfate (AMS). Use UAN under conditions of low relative humidity for greater weed control.

For tank mixtures with Ignite® 280 SL Herbicide

SC 450 Herbicide can be tank mixed with Ignite® 280 SL Herbicide. Ignite® 280 SL Herbicide can only be used on corn seed designated as LibertyLink®. Do not use MSO/ESO or COC adjuvants in this mixture, only add AMS at 8.5 lbs/100 gallons (1.5 lb/A minimum). Follow all other directions for adjuvants as listed on the Ignite® 280 SL Herbicide label. Follow the most restrictive directions for use and precautions as defined on both herbicide labels.

For tank mixtures with Glyphosate (including Roundup and Touchdown branded products)

SC 450 Herbicide can be tank mixed with glyphosate for use on glyphosate-tolerant corn. SC 450 Herbicide will enhance broadleaf control, combat glyphosate-resistant weeds and reduce glyphosate induced weed shifts. Do not use MSO/ESO or COC adjuvants in this mixture. Add AMS at 8.5 lbs/100 gallons (1.5 lb/A minimum) and any additional surfactant as directed on the Glyphosate label. Follow the most restrictive directions for use and precautions as defined on both herbicide labels.

GROUND APPLICATION

DO NOT OVERLAP SPRAY PATTERNS BEYOND EQUIPMENT MANUFACTURERS' RECOMMENDATIONS AS EXCESSIVE RATES MAY RESULT IN ADVERSE CROP RESPONSE.

Apply SC 450 Herbicide alone or in tank mixtures by ground equipment in a minimum of 10 gallons of spray mixture per acre. Uniform, thorough spray coverage is important to achieve consistent weed control. Use sprayers that provide accurate and uniform application. For emerged weed control in dense weed populations or under adverse growing conditions, 15 to 20 gallons of water per acre is recommended. Good coverage is essential to achieve optimum control of emerged weeds.

To minimize spray drift to non-target areas, apply this product using nozzles which deliver a coarse or larger spray droplet as defined by ASAE standard S-572 and as shown in nozzle manufacturer's catalogues. Keep the spray boom at the lowest possible spray height above the target surface. Refer to nozzle manufacturer's recommendations for proper nozzle, pressure setting and sprayer speed for optimum product performance and minimal spray drift.

Over applications, uneven application, sprayers not properly calibrated, boom overlapping or improper incorporation may decrease the level of weed control and/or increase the level of adverse crop response.

Maintain constant ground speed while applying product to ensure proper distribution.

MAINTAIN ADEQUATE AGITATION AT ALL TIMES, INCLUDING MOMENTARY STOPS.

BANDED APPLICATION

Banding herbicide application equipment must be carefully calibrated to prevent crop exposure to concentrations of SC 450 Herbicide that exceed the labeled rate for the soil type. It is critical to insure that the calibrated band width equates to actual band width realized in field applications. Bands actually delivered at a width narrower than targeted will concentrate the product and increase the risk for crop response.

EVEN FLAT SPRAY TIP NOZZLES AND A BAND WIDTH OF NO LESS THAN 12" MUST BE USED.

Band Treatment- Apply a broadcast equivalent rate and volume per acre. The following equations may be used to make the required calculations as follows:

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast RATE per acre} = \text{Amount product needed per acre.}$$

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast spray VOLUME per acre} = \text{Amount Band spray VOLUME needed per acre}$$

AERIAL APPLICATION

Calibrate the spray equipment prior to use. Apply SC 450 Herbicide in a minimum of 5 gallons of water per broadcast acre. The use of nozzles and spray pressure that deliver **MEDIUM** spray droplets as indicated in the nozzle manufacturer's catalogs and in accordance with ASAE Standard S-572 are highly recommended for optimum spray coverage and canopy penetration. **DO NOT** use raindrop nozzles. Aerial applications of this product should be made at a maximum height of 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind. Aerial applications of this product should also be made with low drift nozzles at a maximum pressure of 40 psi. Uniform, thorough spray coverage is important to achieve consistent weed control.

See the **Spray Drift Management** section of this label for additional information on proper application of SC 450 Herbicide.

SPECIFIC DIRECTIONS FOR USE IN CORN

Application of SC 450 Herbicide at less than recommended rates for the appropriate soil will only provide suppression of sensitive weeds.

TABLE 1: SC 450 Herbicide USE RATES When Applied Post-harvest, Preplant Surface, Preplant Incorporated, Preemergence, and Early Postemergence (2 leaf-collar stage or prior) in Corn

Application Timing	fl oz of SC 450 Herbicide per Acre ¹	
	Soil Texture	
	Coarse Soils 2.0% O.M. ² or less (Sand, Loamy sand, Sandy loam)	Coarse Soils greater than 2.0% O.M. ² (Sand, Loamy sand, Sandy loam) And Medium Soils (Loam, Silt loam, Silt, Sandy clay loam) And Fine Soils (Silty clay loam, Clay loam, Sandy clay, Silty clay, Clay)
Post-harvest Preplant³ (Surface Applied or Incorporated) Preemergence Early postemergence	1.33	2.25 ⁴

¹If soils are 2.0% or less in O.M. and have a pH of 7.5 or greater, reduce the rate selected from the table by 0.2 fl. oz.

²O.M. = Organic Matter by weight

³SC 450 Herbicide may be applied alone or in recommended tank-mixes up to 21 days prior to planting. SC 450 Herbicide may be applied up to 30 days prior to planting when used in a planned sequential application program such as SC 450 Herbicide followed by Ignite[®] 280 SL Herbicide, Buctril, or other postemergence applied herbicides.

⁴For coarse textured soils with greater than 2.0% O.M., or medium textured soils with 2.0% O.M. or less, and where densities of weeds controlled by SC 450 Herbicide are light to moderate, an appropriate rate down to 1.80 fl oz per acre may be selected.

Postemergence Rates

- Apply SC 450 Herbicide to emerged weeds at 0.91 fl oz/A per application. Always add appropriate adjuvants to the spray tank (see Spray Additives Section of this label).
- Applications of SC 450 Herbicide at rates less than 0.91 fl oz/A postemergence may result in incomplete weed control and reduction in residual activity.
- Broadcast applications of SC 450 Herbicide may be made to corn from spiking through the 6-leaf collar stage of growth (V6, the first leaf has a rounded tip). Do not apply broadcast to corn that is more mature than the V6 stage of growth.
- Directed postemergence applications of SC 450 Herbicide can be made to corn from the 6-leaf collar stage through the 12-leaf collar stage (V12) of growth. Do not apply to corn that is more mature than the V12 stage of growth.
- Follow all precautions and warnings for using ALS-inhibiting or Sulfonyl urea (SU) herbicides on a particular hybrid/variety.
- Best results with SC 450 Herbicide used postemergence are obtained when it is applied to young actively growing weeds. SC 450 Herbicide will affect weeds that are larger than the recommended height, however such applications may result in incomplete weed control.
- Do not exceed 0.91 fl oz/A of SC 450 Herbicide per postemergence application beyond a 2-leaf collar corn stage (V2).
- DO NOT apply more than two postemergence applications of SC 450 Herbicide to a corn crop in one growing season.
- Allow at least 14 days between applications of SC 450 Herbicide.
- DO NOT graze livestock, or harvest corn forage or sweet corn ears within 45 days of application of SC 450 Herbicide.

TANK-MIX COMBINATIONS

SC 450 Herbicide is recommended as the first herbicide applied in an integrated weed control program that includes sequential postemergence herbicide applications.

Tank mix combinations may be used and applied at the same timings as SC 450 Herbicide unless otherwise specified in the tank-mix partner label. Multiple tank mixtures are allowed unless otherwise specified by the respective product labels. Check all tank-mix product labels for proper rates and compatibilities for multiple tank-mixes.

SC 450 MAY BE TANK-MIXED WITH THESE HERBICIDES FOR CONTROL OF CERTAIN BROADLEAF AND GRASS WEEDS IN CORN.

Tank-mixes with SC 450 Herbicide are not limited to the tank-mix partners listed on this label. Refer and follow the label of each tank-mix partner used for precautionary statements, directions for use, geographic and other restrictions.

TANK-MIX PARTNERS

- 2,4-D
- Atrazine
- Autumn™ Herbicide
- Define™ SC
- Balance® FLEXX
- Glyphosate (including Roundup® and Touchdown® branded products)
- Paraquat (including Gramoxone® branded products)
- LAUDIS™
- Ignite® 280 SL Herbicide
- SIMAZINE/PRINCEP

BROADLEAF AND GRASS WEEDS CONTROLLED BY SC 450 HERBICIDE

ANNUAL GRASS WEEDS		
Barnyardgrass Crabgrass, large Foxtail, giant	Foxtail, green Foxtail, yellow Millet, wild proso Oat, wild	Panicum, fall Sandbur, field ¹ Shattercane Signalgrass, broadleaf
ANNUAL BROADLEAF WEEDS		
Amaranth, Palmer ^{1,2} Lambsquarters, common Morningglory, ivyleaf Morningglory, pitted ¹	Pigweed, redroot ² Purselane, common Ragweed, common ² Sesbania, hemp	Sida, prickly ¹ Sunflower, wild Velvetleaf
¹ These weeds will be suppressed/or be reduced in competition. Reduced competition weeds will be stunted in growth and/or be of reduced populations as compared to non-treated areas. Commercially acceptable control may require the application of an appropriate tank mixture or sequential herbicide treatment. ² SC 450 Herbicide may not provide acceptable control of ALS-resistant biotypes.		

RESISTANCE MANAGEMENT

Naturally occurring biotypes of certain weed species with resistance to ALS-inhibiting herbicides are known to exist. Some weed populations may contain plants naturally resistant to SC 450 Herbicide or other herbicides with the same mode of action (ALS/AHAS

enzyme inhibitors). Repeated use of herbicides with the same mode of action, allow resistant weed populations to gain density and spread. To manage the development and spread of resistant weed populations, use herbicides with different modes of action in sequence, tank mixture, rotation, or in conjunction with alternate cultural practices. SC 450 Herbicide can be used effectively as one of the modes of action in a Balance® FLEXX or Ignite® 280 SL Herbicide program or rotation.

SPRAY DRIFT MANAGEMENT

SC 450 Herbicide is not volatile. Spray drift may result in injury to non target crops or vegetation. To avoid spray drift, do not apply when wind speed is greater than 10 MPH or during periods of temperature inversions. Do not apply when circumstances or weather conditions, wind speed or wind direction may cause spray drift to non-target areas, unprotected persons or to food. AVOIDING SPRAY DRIFT AT THE APPLICATION SITE IS THE RESPONSIBILITY OF THE APPLICATOR.

The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops.

1. The distance of the outer most nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.
3. All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers.

Where states have more stringent regulations, they shall be observed. The applicator should be familiar with and take into account the information covered in the Aerial Drift Reduction Advisory Information.

One of the most effective ways to reduce drift potential is apply the largest droplets that provide sufficient coverage and weed control. Applying larger droplets reduces drift potential, but may not prevent spray drift if unfavorable environmental conditions exists (e.g. wind direction toward sensitive areas or wind speed greater than 10 MPH).

To minimize risk of spray drift, select nozzles and pressures that deliver MEDIUM spray droplets as indicated in nozzle manufacturer's catalogs and in accordance with ASAE Standard S-572. Nozzles that deliver COARSE spray droplets may be used to further reduce spray drift potential, however, application volume per acre (GPA) should be increased to maintain coverage and weed control.

Only apply this product when the potential for drift to adjacent non-target areas is minimal (e.g., when wind is 10 MPH or less and is blowing away from sensitive areas).

To avoid potential adverse effects to non-target areas, maintain a **25 foot buffer for ground applications**, or a **200 foot buffer for aerial applications** between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots hedgerows, riparian areas and shrub lands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

SWATH ADJUSTMENT:

When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind, smaller drops, etc.).

WIND:

Drift potential is lowest between wind speeds of 2 - 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. **NOTE:** Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

TEMPERATURE AND HUMIDITY:

When making applications in low relative humidity, adjust equipment to produce larger droplets in order to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry. Avoid spraying during conditions of low humidity and/or high temperatures.

TEMPERATURE INVERSIONS:

Do not make aerial or ground applications into areas of temperature inversions because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience LP. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience LP is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE LP'S ELECTION, THE REPLACEMENT OF PRODUCT.

NET CONTENTS: 2.5 gallons

Aztec, Balance Pro, Buctril, Ignite, and LibertyLink are registered trademarks of Bayer.
Autumn is a trademark of Bayer.
Gramoxone Max, Force are registered trademarks of a Syngenta Group Company.
Roundup is a registered trademark of Monsanto Company.
Counter, and Regent are registered trademarks of BASF.
Lorsban is a registered trademark of DOW.
Dyfonate, or Thimet.

Produced for



Bayer CropScience

Bayer CropScience LP
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)
<http://www.bayercropscienceus.com>

SC 450 Herbicide (PENDING) 08/10/10