

[279-40]

INSTRUCTIONS

Apply to control *Colorado Potato Beetle* and *Rice Beetle* use 20 to 30 pounds per acre. (through coverage) of under surfaces of leaves in particular to effect control.

Apply to control *Rice Beetle* use 10 to 15 pounds per acre.

Apply to control *Colorado Potato Beetle* and *Rice Beetle* use 20 to 30 pounds per acre.

Apply to control *Cucumber Beetle*, *Squash Bug* and *Aphids* use 20 to 30 pounds per acre.

Apply to control *Colorado Potato Beetle* and *Rice Beetle* use 20 to 30 pounds per acre.

Apply to control *Red Wasp* and *Wasp* use 20 to 30 pounds per acre. Apply at 7-day intervals and use as necessary for control. For best results, make first application when bees are active in field and repeat 3 to 7 days later, on this interval, applications at 7-day intervals may be necessary.

Apply to control *Aphid*, *Rice Beetle* and *Wasp* use 20 to 30 pounds per acre. First application when insects appear and repeat 3 days later.

Apply to control *Cucumber Beetle*, *Squash Bug* and *Aphids* use 20 to 30 pounds per acre.

Apply to control *Rice Beetle* and *Rice Beetle* use 20 to 30 pounds per acre.

Apply to control *Rice Beetle*, *Colorado Potato Beetle* and *Rice Beetle* use 20 to 30 pounds per acre.

Apply to control *Cucumber Beetle*, *Squash Bug* and *Aphids* use 20 to 30 pounds per acre.

APPROVED

DATE

SIGNATURE

ACCEPTED

UNDER THE FEDERAL INSECTICIDE, FUNGICIDE AND ROSENICIDE ACT FOR ECONOMIC POISON REGISTERED UNDER NO. SUBJECT TO ATTACHED COMMENTS