

C-O-C-S

PREPARED ADDITIONAL DIRECTIONS FOR USE

PEACHES: Leaf Curl - Use 4 pounds per 100 gallons of water.
Apply at pre-blossom to petal fall.

**DIRECTIONS FOR USE OF C-O-C-S
DO NOT USE IN UNDILUTED FORM.**

THE CONTENTS OF THIS PACKAGE SHOULD NOT BE USED AS A DUST. Use Niagara C-O-C-S for the control of those diseases against which an insoluble copper fungicide may be used. Make the first application when the disease to be controlled is reported to be present in the area. Repeat at 7 to 10 day intervals or as required to maintain effective control. Use in sufficient water to thoroughly wet the plants.

Fill the spray tank 1/2 to 3/4 full of water. With the agitator running slowly sift in the required amount of Niagara C-O-C-S, adding afterwards the required amount of insecticide and lime, if needed. Finish filling the spray tank with water and apply at once.

CARROT: Cercospora Leaf Spot, Macrosporium Leaf Spot—Use 4 pounds per acre.

CELERY: Early Blight (Cercospora), Late Blight (Septoria)—Use 4 pounds per acre.

CUCUMBERS, MELONS: Anthracnose, Angular Leaf Spot, Downy Mildew, Scab—4 pounds per acre.

POTATOES: Early Blight, Late Blight—Use 4 pounds per acre. If late blight infection is severe the dosage rate on large plants should be increased to 7 pounds per acre. Spray plants thoroughly.

PEPPERS, TOMATOES IN COLD FRAMES, GREENHOUSE, ETC.: Damping off—Use 4 pounds in 100 gallons of water or 2 ounces in 3 gallons. The suspension should be kept well agitated. Apply at 4 to 7 day intervals, beginning when the plants break through the ground. Spray the plants lightly but thoroughly. Also apply to the soil surrounding the plants. Do not use on cabbage and related crops for the control of damping-off.

TOMATOES: Early Blight, Late Blight, Septoria Leaf Spot—Use 4 pounds per acre. If late blight infection is severe the dosage rate on large plants should be increased to 7 pounds per acre. Spray plants thoroughly.

Bacterial Spot—Use 5 to 6 pounds per acre. In the seed bed, apply every 5 to 6 days after the first leaves appear. In the field, apply at 5 to 7 day intervals if the disease is known to occur in the area. During periods of rainy weather, use the shorter interval between sprays. Spray plants thoroughly.

TART CHERRY: Cherry Leaf Spot—Use 1 1/2 pounds plus 3 pounds of hydrated spray lime in 100 gallons of water.

GRAPES: Downy Mildew, Powdery Mildew, Black Rot—Use 2 pounds per 100 gallons plus 4 pounds of hydrated lime. Spray thoroughly both upper and lower leaf surfaces.

SUGAR BEETS: Cercospora Leaf Spot—Use 6 pounds per acre in sufficient water to obtain thorough coverage. Begin applications when disease first appears and repeat at 7 to 10 day intervals for as long as disease threatens. May be applied up to day of harvest.

CITRUS including Grapefruit, Lemons, Limes, Oranges, Tangerines, Tangelos: Melanose, Scab, Greasy Spot—Use 6 pounds in 500 gallons of water. Apply in dormant season before Spring growth appears. Repeat 1 to 3 weeks after petal fall. For melanose and scab repeat 4 weeks later. Do not apply after fruit is 1/4 inches in diameter.

ORNAMENTALS:

AZALEA: Bud Blight, Twig Blight—Use 4 pounds per 100 gallons of water.

VINCA: Canker, Dieback—Use 4 pounds per 100 gallons of water.

MAPLE: Tar Leaf Spot—Use 4 pounds plus 2 pounds hydrated lime per 100 gallons of water.

OAK (WHITE): Anthracnose—Use 4 pounds plus 2 pounds hydrated lime per 100 gallons of water.

PACHYSANDRA: Canker, Twig Blight, Leaf Spot—Use 3 pounds plus 3 pounds spray lime per 100 gallons of water.

ACCEPTED
Dec 10, 1968
UNDER THE FEDERAL INSECTICIDE AND FUNGICIDE ACT
REGISTERED UNDER NO. 279-69
TO: AGRICULTURAL COMMISSION

279-69

Niagara
C-O-C-S

Bag No. 5855



CODE 557

3 LBS. NET WT.

C-O-C-S
an insoluble COPPER COMPOUND
FUNGICIDE

USDA REG. NO. 279-69

ACTIVE INGREDIENT:

*Copper expressed as metallic..... 55.0%

INERT INGREDIENTS: 45.0%

*from copper oxychloride sulphate 100.0%

CAUTION
KEEP OUT OF REACH OF CHILDREN.
HARMFUL IF SWALLOWED!

Rinse equipment and DISPOSE OF WASTES by burying in non-crop lands away from water supplies. Container should be disposed of by burying with wastes or by burning. Keep out of smoke.

DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROPS OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS BECAUSE ANIMAL, PLANT OR CROP INJURY, ILLEGAL RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR.

DEALERS SHOULD SELL IN ORIGINAL PACKAGES ONLY

TERMS OF SALE OR USE: On purchase or use of this product buyer and user agree to the following conditions:

WARRANTY: Niagara warrants that this product, as of the time of sale by Niagara, (1) conforms to the ingredient statement on the label, and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS. NIAGARA MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

DAMAGES: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages. Notice of any claim must be reported to Niagara within 10 days of discovery by buyer or user, failing which buyer or user waives any claim for such damage.

EXCUSE FROM NON-CONTROLLABLE CONDITIONS: Because of certain critical conditions created or incurred by buyer or user or over which Niagara has no control, buyer and user, as a condition of purchase or use, assume responsibility for and release Niagara from liability arising out of the handling of use of the product attributable to such causes. No person is authorized to vary or waive any statement contained herein.



NIAGARA CHEMICAL DIVISION
FMC CORPORATION
MIDDLEPORT, N. Y.



MANUFACTURED IN U.S.A. FOR



PLANTS

AYER, MASS.	GREELEY, COLO.	MALAGA, N.J.
NAPOLEON, Q.	FORT MEAL, IA.	SOUTH HAVEN, MICH.
WYOMING, ILL.	JACKSONVILLE, FLA.	AYDEN, N.C.
DOTHAN, ALA.	GREEKVILLE, MISS.	OPELOUSAS, LA.
PINE BLUFF, ARK.	POMPANO BEACH, FLA.	TAMPA, FLA.

019



Printed in U.S.A.