

CODE
114

CALCIUM ARSENATE

NET WEIGHT
50 LBS.

INSECTICIDE - HERBICIDE

DIRECTIONS FOR USE

ORNAMENTAL TURF, HOME LAWNS: Crabgrass - Use 13 lbs. per 1,000 sq. ft. of turf. Apply in sufficient water to obtain uniform coverage of treated area. Application should be made in the late winter or spring prior to the emergence of the crabgrass seedlings. The treated area should be well watered immediately after application. Keep children and animals off treated area until this material has been thoroughly watered into the soil and the grass has completely dried. Apply to well-established lawns only. This material may injure certain varieties of bentgrass, blue grass and fescues. Injury will usually be in the form of a thinning of the grass stand.

Multiple applications during the same year, or single application of the above dosage for two or more consecutive years is not recommended. Treatment should be limited to a single application of the above dosage during the first year treatment is undertaken and in succeeding years, treatments of one-third the initial dosage may be applied should the problem persist. This reduction of the dosage in subsequent treatments is designed to prevent excessive thinning of turf. If additional crabgrass control is desired within one year from the date of a calcium arsenate application, another recommended control measure should be used.

If reseeding of the treated area is desired, such reseeding should not be attempted within 6 months of the time of treatment.

POULTRY HOUSES: For control of the larval (maggot) stage of the house fly and soldier fly in litter droppings under caged chickens and turkeys on wire floors. Use 2.4 lbs. in 4 gallons of water. Agitate well and apply as a coarse spray to 1000 sq. ft. of droppings or manure piles on a preventative schedule. Repeat as manure is added. Use only on droppings where poultry cannot come in contact with the material. Do not spray directly on poultry or feed and water. Repeat on 5-7 day intervals.

CAUTION: Do not apply manure treated with this product to food crops.

TOMATOES: Tomato Fruit Worm, Hornworm, Flea Beetles - Use at the rate of 3 lbs. per acre in sufficient water to provide uniform coverage. Repeat applications as required. Remove residues remaining at harvest by washing, wiping, brushing or other effective means.

U. S. D. A. Reg. No. 279-85

ACTIVE INGREDIENTS:

Tricalcium Arsenate ----- 70.0%

INERT INGREDIENTS: ----- 30.0%

100.0%

Arsenic expressed as metallic ----- 26.0%


Arsenic in Water Soluble Forms expressed as metallic - 0.5%



KEEP OUT OF REACH OF CHILDREN

ANTIDOTE: Give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Then give two tablespoonfuls of epsom salts or milk of magnesia in water, and plenty of milk and water. Have victim lie down and keep quiet. Call a Physician IMMEDIATELY!

WARNING: Poisonous if swallowed. Avoid contact with skin, eyes, or clothing. Wash thoroughly after using. Avoid breathing dust or spray mist.



WARNING - POISON

This package contains ARSENIC or combination thereof for agricultural purposes only. Do not keep or use in a room where food is being manufactured or prepared or where powdered food is stored.

Do not use or store in or around the home.

USAGE CAUTIONS

This product is toxic to wildlife. Keep out of any body of water. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Dispose of wastes by burying in non-crop lands away from water supplies. Container should be disposed of by burying with wastes or by burning. Keep out of smoke.