

CODE

30319

C-O-C-S® COPOZIM SPRAY

NET WEIGHT

30 LBS.

FUNGICIDE - NUTRITIONAL

DIRECTIONS FOR USE

CROPS - Tomatoes, potatoes, celery, cucumbers, squash, melons and peppers.

DISEASES - Early blight, late blight, downy mildew, anthracnose, Alternaria leaf spot, gray leaf spot, and bacterial leaf spot.

INSECTS - None

NUTRITIONAL - Aids in correcting deficiencies of zinc and manganese.

Use 6 pounds to 100 gallons of water. Draw spray tank about 2/3 full of water. While agitator is running pour in required amount of this material. Keep agitator running until tank is sprayed out. If insecticides such as lead arsenate, DDT, BHC,

U. S. D. A. Reg. No. 279-485

ACTIVE INGREDIENTS:

*Copper, expressed as metallic-----33.3%

INERT INGREDIENTS: **-----66.7%

100.0%

*The copper is in the form of copper-oxochloride-sulfate

**Manganese expressed as metallic ---8.0%

Zinc expressed as metallic -----3.0%

CAUTION

KEEP OUT OF REACH OF CHILDREN

HARMFUL IF SWALLOWED!

USAGE CAUTIONS

This product is toxic to fish. Keep out of any body of water. Apply this product only as specified on this label.

Rinse equipment and dispose of wastes by burying in non-crop lands away from water supplies. Bags should be disposed of by burying with wastes or by burning. Keep out of smoke.

Chlordane or Parathion wettable powders are used add to tank last.

Begin applications when diseases are apt to appear and repeat at 7 to 10 day intervals.

The rate of application per acre ranges from 75 to 150 gallons depending on the size of the plants.

