

[279-776] 1 of 3

Code: 30237

KOLODUST[®] MALATHION 50

DIRECTIONS FOR USE

Make thorough, full coverage application when insects or disease first appear and repeat as necessary. May be applied by air or ground. Observe days interval between application and harvest indicated by number in () following the crop.

APPLES (3): Aphids, Mealybug, Codling Moth, Fruittree Leafroller and Powdery Mildew — Use 50 to 80 pounds per acre. Sulphur and Malathion may injure certain varieties of apples, check tolerance prior to use.

BLACKBERRIES (1), BOYSENBERRIES (1), DEWBERRIES (1), LOGANBERRIES (1), RASPBERRIES (1): Powdery Mildew, Aphids and Rose Scale — Use 40 to 60 pounds per acre.

CHERRIES (3): Black Cherry Aphid, Fruittree Leafroller and Brown Rot — Use 50 to 80 pounds per acre. May injure foliage of some varieties.

GRAPES (3): Leafhopper, Drosophila and Powdery Mildew — Use 30 to 50 pounds per acre. Ground application preferred. Emulsion may cause injury to foliage on some varieties.

PECANS (0): Pecan Nut Casebearer — Use 100 pounds per acre. Apply when first brood larvae begin to enter nuts and repeat after 7 days as necessary. Pecan Bud Moth — Apply 10 pounds per acre by airplane. Apply 4 to 5 times, 2 to 3 weeks apart when insects appear.

PEACHES (7), NECTARINES (7): Peach Silver Mite, Aphids, Oriental Fruit Moth, Brown Rot and Powdery Mildew — Use 50 to 80 pounds per acre.

PEARS (1): Pear Psylla, Mealybug, Codling Moth, Fruittree Leafroller and Powdery Mildew — Use 50 to 80 pounds per acre.

PLUMS (3), PRUNES (3): Aphids, Brown Rot and Prune Rust — Use 50 to 80 pounds per acre.

STRAWBERRIES (3): Aphids, Lygus Bugs, Spittlebug, Field Crickets, Thrips, Potato Leafhopper, Whitefly, Strawberry Leafroller and Powdery Mildew — Use 50 pounds per acre.

BEANS (1): Mexican Bean Beetle, Leafhoppers, Aphids, Cucumber Beetle and Powdery Mildew — Use 30 to 40 pounds per acre. Sulphur may injure certain varieties, check sulphur tolerance prior to use.

BROCCOLI (3), CABBAGE (7), KALE (7), TURNIPS (3): Aphids, Imported Cabbageworm and Cabbage Looper — Use 35 to 60 pounds per acre.

BRUSSELS SPROUTS (7), COLLARDS (7): Aphids and Imported Cabbageworm — Use 35 to 60 pounds per acre.

CARROTS (7): Aphids and Leafhoppers — Use 35 to 50 pounds per acre.

CAULIFLOWER (7): Aphids — Use 50 to 60 pounds per acre. Imported Cabbageworm — Use 30 pounds per acre.

FUNGICIDE-INSECTICIDE

EPA REGISTRATION NO. 279-776 AA

ACTIVE INGREDIENTS:	BY WT.
Malathion; 0,0-Dimethyl dithiophosphate of diethyl mercaptosuccinate	4.00%
Sulphur	42.00%
INERT INGREDIENTS:	54.00%
	100.00%

CAUTION

KEEP OUT OF REACH OF CHILDREN


Avoid breathing of dust. May be harmful if swallowed. Avoid contact with skin. Wash thoroughly after using. Avoid contamination of feed and foodstuffs.

NOTE TO PHYSICIAN: Malathion may cause cholinesterase inhibition. Atropine is antidotal.

For Emergency Assistance Call
716-735-3765

Applications of materials containing Sulphur should not be made just prior to or during periods of excessively high temperatures or within 10 days to two weeks of an oil spray.

RECEIVED
279-776
EPA



[279-776] 2 of 3

4 DUST

**Net We
50 Pounds**

COTTON (0): Cotton Aphid, Leafhoppers, Whitefly, Cotton Leafworm, Cotton Leafperforator, Thrips, Lygus Bugs, Fleahoppers, Fall Armyworms, Grasshoppers, Garden Webworms — Use 30 to 60 pounds per acre.

LETTUCE (Head-7, Leaf-14): Aphids and Leafhoppers — Use 30 pounds per acre. Cabbage Looper — Use 35 to 50 pounds per acre.

MUSTARD GREENS (7): Aphids, Imported Cabbageworm, Cabbage Looper and Flea Beetles — Use 35 to 50 pounds per acre.

ONIONS (3): Thrips and Onion Maggots — Use 35 to 50 pounds per acre. Apply when Maggot flies appear and repeat every 4 days.

PEAS (3) (Forage-7): Pea Aphid, Pea Weevil and Powdery Mildew — Use 30 pounds per acre.

PEPPERS (3): Pepper Maggots — Use 35 pounds per acre.

POTATOES (0): Aphids, Leafhoppers, False Chinch Bug and Mealybug — Use 35 to 60 pounds per acre.

TOMATOES (1): Aphids, Tomato Russet Mite and Drosophila — Use 40 to 50 pounds per acre.

ORNAMENTALS: Aphids, Rust and Powdery Mildew. Make thorough full coverage application. Injury may occur on certain ferns and on some species of crassula. Do not use on canaerti juniper.

ENVIRONMENTAL AND DISPOSAL CAUTIONS


This product is toxic to fish. Keep out of lakes, ponds and streams. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from the areas treated. Do not apply where run-off is likely to occur. Apply this product only as specified on the label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your cooperative Agriculture Extension Service.

DISPOSE OF WASTES by burying in non-crop land away from water supplies. Empty containers should be buried with wastes or burned. Keep out of smoke.

ADDITIONAL INFORMATION APPEARING ON BAGS TO WHICH THIS LABEL WILL BE ATTACHED.

[279-776] 3 of 3



Dealers Should Sell in Original Packages Only.

DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROPS OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS BECAUSE ANIMAL, PLANT OR CROP INJURY, ILLEGAL RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR.

TERMS OF SALE OR USE: On purchase or use of this product buyer and user agree to the following conditions:

WARRANTY: FMC warrants that this product, as of the time of sale by FMC, (1) conforms to the ingredient statement on the label, and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS. FMC MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

DAMAGES: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages. Notice of any claim must be reported to FMC within 10 days of discovery by buyer or user, failing which buyer or user waives any claim for such damage.

EXCUSE FROM NON-CONTROLLABLE CONDITIONS: Because of certain critical conditions created or incurred by buyer or user or over which FMC has no control, buyer and user, as a condition of purchase or use, assume responsibility for and release FMC from, liability arising out of the handling or use of the product attributable to such causes. No person is authorized to vary or waive any statement contained herein.

Manufactured in U.S.A. by:
FMC CORPORATION
AGRICULTURAL CHEMICAL DIVISION
Middleport, N.Y. 14105

FMC
Chemicals