

Code: 1085

50 Pounds Net

GRANULAR 10% CHLORDANE

DIRECTIONS FOR USE

Granular 10% Chlordane is designed for application to soil for the control of insects attacking the roots and young plants of certain crops and for control of Ants in Citrus groves.

CORN, MAIZE, SOYBEANS, CORN, TOMATOES and CARROTS to control Wireworms. Apply 25-50 pounds per acre, depending upon soil type, as an overall soil application prior to planting. Sandy soils require lower amounts. Work into top 3 to 6 inches of soil.

CORN to control Corn Rootworm. Apply 20 pounds per acre as an overall soil application prior to planting. Work into top 3 to 6 inches of soil.

IN CITRUS GROVES (Oranges, Grapefruit, Kumquats, Lemons, Lime, Tangerines, Tangelos) to control Ants. Use 50 pounds per acre. Apply to surface of ground when ants appear and repeat as necessary to maintain control. Do not apply after fruit formation.

BEANS, CELERY, CUCURBITS, LETTUCE, PEPPERS and TOMATOES to control Cutworms and Flea Beetles. Use 20 pounds per acre. Apply to surface of soil around base of plants when insects first appear. Do not apply after edible parts start to form. May be injurious to certain varieties of cucurbits.

BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER, CELERY, CORN, PEPPERS, POTATOES, SWEET POTATOES and TURNIPS to control White Grubs, Wireworms and Root Maggots. Use 25-50 pounds per acre depending upon soil type. Sandy soils require lower amount. Apply as an overall soil application prior to planting. Work into top 3 to 6 inches of soil.

ON LAWNS and TURF to control Ants and European Earwig use 10-15 pounds per acre. To control White Grubs, Wireworms, Sod Webworm, and Cutworms use 25 pounds per acre. For 4-5 years control increase dosage to 50-100 pounds per acre. On lawn and turf areas wash down into turf and soil. For non-seeded areas, apply and cultivate into soil to a depth of 4-6 inches immediately after applying. Keep children and domestic pets off lawns and turf until they have been thoroughly watered and completely dry. Birds feeding over treated lawn may be harmed.

STRAWBERRY to control Strawberry Root Weevil, White Grubs and Strawberry Rootworm use 100 pounds per acre. Apply as an overall soil application prior to planting. Work into top 3 inches of soil.

INSECTICIDE

U.S.D.A. REGISTRATION NO. 279-1023

ACTIVE INGREDIENT:	BY WT.
*Technical Chlordane - - - - -	10.00%
INERT INGREDIENTS - - - - -	90.00%
	100.00%

*Equivalent to 6% Octachloro-4,7-Methanotetrahydroindane and 4% related compounds.


CAUTION

KEEP OUT OF REACH OF CHILDREN

Avoid inhalation.

Avoid contamination of feed and foodstuffs.

Japanese Beetle Grub



MANUFACTURED BY


DIAL CORPORATION

NEW YORK, N. Y.

ROSELAND, N. Y.

Toxic to fish and wildlife. Do not contaminate water, by direct application, cleaning or disposal of wastes and containers in area not specified on the label.

Dispose of wastes by burying in non-crop land supplies. Bag should be disposed of by burying or burning. (Keep Out of Smoke).



USAGE CAUTION: Do not allow this material to contact neighboring crop or non-crop areas or use in a time other than in accordance with directions. Injury, excessive residues or other undesirable effects may occur.

Dealers Should Sell in Original Packages Only

50 Pounds Net Weight GRANULAR 10% CHLORDANE

SE
 gned for application to
 ing the roots and young
 of Ants in Citrus groves.
 S to control Wireworms.
 g upon soil type, as an
 ing. Sandy soils require
 hes of soil.
 0 pounds per acre as an
 ng. Work into top 3 to 6
 Kumquats, Lemons, Lime,
 Use 50 pounds per acre.
 s appear and repeat as
 apply after fruit form-

PERS and TOMATOES to con-
 pounds per acre. Apply to
 hen insects first appear.
 to form. May be injurious

AULIFLOWER, CELERY, CORN,
TURNIPS to control White
 25-50 pounds per acre de-
 quire lower amount. Apply
 o planting. Work into top

European Earwig use 10-15
 s. Wireworms, Sod Webworm,
 For 4-5 years control in-
 e. On lawn and turf areas
 i-seeded areas, apply and
 inches immediately after
 pets off lawns and turf
 ared and completely dry.
 be harmed.

t Weevil, White Grubs and
 r acre. Apply as an over-
 g. Work into top 3 inches

INSECTICIDE
 U.S.D.A. REGISTRATION NO. 279-1025


ACTIVE INGREDIENT:	BY WT.
*Technical Chlordane - - - - -	10.00%
INERT INGREDIENTS - - - - -	90.00%
	100.00%

*Equivalent to 6% Octachloro-4,7-Methanotetrahydroindane
 and 4% related compounds.

CAUTION
KEEP OUT OF REACH OF CHILDREN

Avoid inhalation.
 Avoid contamination of feed and foodstuffs.


Japanese Beetle Grub



MANUFACTURED BY U.S.A. BY
 PRAIRIE INDUSTRIES DIVISION
 CHEMICAL CORPORATION
 BROOKLYN, N. Y.

Toxic to fish and wildlife. Do not contaminate any body
 of water, by direct application, cleaning of equipment
 or disposal of wastes and containers nor apply to any
 area not specified on the label.

Dispose of wastes by burying in non-crop lands away from water
 supplies. Bag should be disposed of by burying with wastes or by
 burning. (Keep Out of Smoke).



Handwritten signature or initials: JH-1-23

USAGE CAUTION: Do not allow this material to drift onto
 neighboring crop or non-crop areas or use in a manner or at a
 time other than in accordance with directions, because plant
 injury, excessive residues or other undesirable results may
 occur.

Dealers Should Sell in Original Packages Only.

Cameraman's Note

Poor Copy