CODE 35

THIRAM 4.9 DUST

NET WEIGHT

50 LBS.

EPA Reg No 279-1561

ACTIVE IN	GREDIENT
------------------	----------

*Tetramethylthiuram disulfide

CAUTION

KEEP OUT OF REACH OF CHILDREN.

Consumption of alcoholic beverages increase possibility of harm.

For Emergency Assistance Call 716-735-3765 MAR 2 8 1974

ONE OF THE FEDERAL INSECTICIDE ACTION REGISTER

ED UNDER NOW 72 26 STORES

279-1561

ENVIRONMENTAL AND DISPOSAL CAUTIONS

This product is toxic to fish. Keep out of lakes, ponds and streams. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

JISPOSE OF WASTES by burying in non-crop lands away from water supplies. Container should be disposed of by burying with wastes or by burning. Keep out of sinoke.

DIRECTIONS FOR USE

APPLE SCAB: Apply as a protective fungicide in the pre-bloom, bloom and early cover stages using 40 pounds per acre on mature or nearly mature size trees. Make repeated applications as required to keep the foliage fully protected from the time new growth starts in the spring.

CEDAR APPLE RUST: Apply in pre-bloom, bloom, calyx and first cover stages as is usually determined for apple scab. Apply 40 pounds per acre on mature or nearly mature size trees.

OTHER APPLE DISEASES: For the control of Black Rot on fruit and foliage, Fly Speck, Sooty Blotch, Bitter Rot, Brooks Spot, Apple Blotch, Botryospheria Spot and Black Pox make applications as indicated for scab control since these are effective in controlling the diseases listed. When weather conditions favor the disease, shorten the application in erval and make applications in advance of infection periods.

FEACH BROWN ROT: For blossom blight apply at to 4 day intervals during bloom, timing the applications to keep newly opening blossoms covered. For fruit rot apply at petal fall and two weeks later. Take one or more additional applications beginning one month before harvest repeating at 7 day intervals. Begin earlier if any rot is seen on green fruit. Apply 40 pounds per acre. Do not apply within 7 days of harvest.

PEACH RHIZOPUS ROT: Apply as directed for fruit phase of brown rot.

STRAWBERRIES: For Gray Mold (Botrytis) apply 40 pounds per acre at 10 day intervals beginning at early blossom stage and continuing as necessary. Good coverage of buds, blossoms and developing fruit is required for best results. If applied within 3 days of harvesting remove residue by washing.

ADDITIONAL INFORMATION APPEARING ON BAGS TO WHICH THIS LABEL WILL BE ATTACHED.

LABEL

279-1561

Dealers Should Sell in Original Packages Only.

DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGH-BORING CROPS OR NON-CROP AREAS OR USE IN A MAN-NER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS BECAUSE ANIMAL, PLANT OR CROP INJURY, ILLEGAL RESIDUES OR OTHER UN-DESIRABLE RESULTS MAY OCCUR.

TERMS OF SALE OR USE: On purchase or use of this product buyer and user agree to the following conditions:

WARRANTY: FMC warrants that this product, as of the time of sale by FMC, (1) conforms to the ingredient statement on the label, and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS FMC MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED

DAMAGES: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages. Notice of any claim must be reported to FMC within 10 days of discovery by buyer or user, failing which buyer or user waives any claim for such damage. EXCUSE FROM NON-CONTROLLABLE CONDITIONS: Because of certain — cal conditions created or incurred by buyer or user or over which FMC has no control, buyer and user, as a condition of purchase or use, assume responsibility for and release FMC from, liability arising out of the handling or use of the product attributable to such causes. No person is authorized to vary or waive any statement contained herein.

Manufactured in U.S.A by: FMC CORPORATION AGRICULTURAL CHEMICAL DIVISION Middleport, N.Y. 14105

