

CODE
113

C-O-C-S[®] THIODAN[®] 3 DUST

FUNGICIDE - INSECTICIDE

NET WEIGHT
50 LBS.

DIRECTIONS FOR USE

Apply at first sign of disease or insect activity and repeat as required to maintain control. Dust plants thoroughly, particularly the underside of the leaves.

CUCUMBERS, PUMPKINS, SUMMER SQUASH, AND WINTER SQUASH: Aphids, Cucumber Beetles, Striped Cabbage Flea Beetle, Squash Vine Borer, Bacterial Wilt, Scab, Leaf Blights, Downy Mildew - Use 25 to 33 lbs. per acre. May be applied up to day of harvest. For vine borer apply weekly to flower buds, stems and vines beginning when moths first appear.

GRAPES: Grape Leafhopper, Downy Mildew, Powdery Mildew - Use 35 to 50 lbs. per acre. Do not use on Concord variety as injury is likely to occur. Do not apply within 7 days of harvest.

MELONS: Aphids, Cucumber Beetles, Striped Cabbage Flea Beetle, Cabbage Looper, Omniverous Leafroller - Use 35 to 50 lbs. May be applied up to harvest.

PEPPERS: Flea Beetle, Green Peach Aphid, Bacterial Spot - Use 33 lbs. per acre. Do not apply within 1 day of harvest.

POTATOES: Flea Beetle, Colorado Potato Beetle, Leafhopper, Aphids, Early Blight, Late Blight, Southern Armyworm, Green Stink Bugs, Leaf-footed Plant Bug and other plant bugs, Potato Tuberworm, Whitefly, False Chinch Bug, Potato Psyllid, European Corn Borer, Three-lined Potato Beetle - Use 30 lbs. per acre. Apply as needed up to harvest. Do not plant root crops other than carrots, potatoes, and sweet potatoes as follow-up crops.

E. P. A. Reg. No. 279-1589

ACTIVE INGREDIENTS:

*Endosulfan (Hexachlorohexahydromethano-
2,4,3-benzodioxathiepin oxide) ----- 3.0%
Copper, expressed as metallic ----- 6.0%

INERT INGREDIENTS: ----- 91.0%
100.0%

*Thiodan U. S. Pat No. 2,799,685
The copper is in the form of basic sulfates and chlorides.

CAUTION KEEP OUT OF REACH OF CHILDREN.

Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing of dust. Avoid contact with skin, eyes, or clothing. In case of contact with skin or eyes, flush with plenty of water; for eyes, get medical attention. Wash with soap and water after handling and before eating or smoking; wear clean clothing. Avoid contamination of feed and foodstuffs. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals.

Workers entering treated fields within 24 hours should wear protective clothing.

ANTIDOTE

Endosulfan is a central nervous system stimulant. There is no specific antidote. Barbituric acid derivatives may be used for convulsions. Treat symptomatically.

TOMATOES: Aphids, Blister Beetle, Colorado Potato Beetle, Flea Beetle, Green Stink Bug, Whitefly, Hornworm, Early Blight, Late Blight, Septoria Leaf Spot, Bacterial Spot - Use 30 to 33 lbs. per acre. Do not apply within 1 day of picking.

USAGE CAUTIONS

This product is toxic to fish and wildlife. Keep out of any body of water. Birds feeding on treated areas may be killed. Do not apply when weather conditions favor drift from the areas treated. Do not apply when runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.



This product is toxic to bees and should not be applied when actively visiting the area.

DISPOSE OF WASTES by burying in non-crop lands away from water supplies. Container should be disposed of by burying with wastes or by burning. Keep out of smoke.

Jul 6, 1971

277-1119

ADDITIONAL INFORMATION APPEARING ON BAGS TO WHICH THIS LABEL WILL BE ATTACHED



Dealers Should Sell in Original Packages Only.

DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO FIGHTING FIRE OR TO BE IN NEARBY AREAS OR USE IN A MANNER NOT AS INTENDED HEREIN. IN ACCORDANCE WITH FEDERAL AND STATE LAWS, BECAUSE ANIMAL, PLANT OR HUMAN LIFE IS AT RISK, RESIDUES OR OTHER UNDESIRABLE BY-PRODUCTS MAY OCCUR.

MAY 1964



NIAGARA CHEMICAL DIVISION
FMC CORPORATION
MIDDLEPORT, NEW YORK

