11**79**



NET WEIGHT

FUNGICIDE-INSECTICIDE

DIRECTIONS FOR USE

Apply when insects or disease first appear and repeat as required to maintain effective control. Coverage of both upper and lower leaf surfaces is essential for good control. Observe use limitations. It insect control is required beyond the following use patterns, supplement the control program with other suitable pesticides. Observe days interval between last application and harvest indicated by number in () following crop.

COTTON: Cotton Stink Bugs, Lygus Bugs Bell-worms, Cabbage Looper-Use 35 lbs per acre. Apply at 3 to 5 day intervals depending upon the severity of the infestation and local conditions. Do not apply after bolls begin to open. Do not graze meat or dairy animals in treated fields.

GRAPES (7): L. afhoppers, Leaffolders (Calif.) Powderv Mildew-Use 30 to 35 lbs. per acre. Do not apply to Concord variety of grapes.

PEAS, SEED (0): Pea Aphids, Pea Weevil-Use 35 Ibs per acre. Do not feed treated vines or threshings to dairy or meat animals.

POTATOES (0): Aphids Colorado Potato Beetle, Stink Bugs, Plant Bugs (including Shield Shaped Bugs), Potato Flea Beetle Potato Leathopper Potato Tuberworm, Southern Amyworr, Use 30 to 35 lbs per acre; Furopean Corn Borer, 2 tato Psyllid-Use 35 lbs per acre; False Chinch Bug, Use 35 lbs per acre; Three lined Potato Beetle Use 17 lbs, per acre.

USDA REG. NO. 279 1618

					100.0
INERT INGREDIEN	TS .				477, ()
methano 2.4,3-ber				.l.,	3.0
: Endosultan (Hex.	achlorohes	ahyo	110		
Sultai					50 O
ACTIVE INGREDIE	N18				


Throdan, U.S. Pat. Ao. 2,799,685

Sultur fineness 98 1 thm 325 mesh

CAUTION KEEP OUT OF REACH OF CHILDREN.

Harmful it swallowed, inhaled or absorbed through skin. Avoid breathing of dust. Avoid confact with skin, eyes or clothing. In case of confact with skin or eyes, flush with plenty of water, for eyes, get medical attention. Wash with soap and water after handling and before eating or sacking; wear clean clothing. Avoid contamination of feed and food-stuffs. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals.

Workers entering treated fields within 24 hours should wear protective clothing.



Thiodan is a registered trademark of Canadian Hoechst Limited

USAGE TAUTION:

Applications of spraying or dusting materials containing sultur should not be made just prior to or during periods of excessively high temperatures.

The product is toxic to fish and wildlife. Keep out of any bod, of water. Apply this product only as specified on the label.

DISPOSE OF WASTES by burying in non-crop lands away from water supplies. Container should be disposed or by Eurying with wastes or by burning. Keep out of smoke

BORING CROES OF TON CEOF AREAS OF SOLING MANAGED OR AT A TIME OFFIEL THAN IN ACCOURT OR A LEGISLIC TON A MANAGED DIRECTION BECAUSE AN MALL PLANS OF SOLID FOR MANAGED HAVE OFFIELD OR A LEGISLIC TON A LEGISLIC TON AND A LEGISLIC TON A LEGISLIC TON AND A LEGISLIC TON AND A LEGISLIC TON A LEG

Dealers Should Sell in Original Packages Only

TERMS OF SALE OR USE On purchase ruse fibis or dictable result user archiver the translations of the same and user archiver the translations of the same archiver the same archives are same archives and same archives are same arc

I. AMA also the control of the property of the property of the control of the control of the control of the property of the control of the

EXPLISE FROM NON-CONTROL CAPITE CONTROL NO cause of freedral control of a red freedral control of a red freedral control of a grant freedral control of the freedral control of a red freedral control of the freedral from the freedral from the freedral control of the product attributed per control of the product of water and street control of the product of water and street control of the product of water and street control of the freedral of the freedral control of the freedral cont

- - - -

CODE 1179

() following crop

animals in treated fields

apply to Concord variety of grapes

ings to dairy or meat animals.

DIRECTIONS FOR USE

Apply when insects or disease first appear and re-

peat as required to maintain effective control. Cov-

tial for good control. Observe use limitations if

insect control is required beyond the following use

patterns, supplement the control program with other

suitable pesticides. Observe days interval between

last application and harvest indicated by number in

COTTON: Cotton Stink Bugs, Lygus Bugs Boll-

worms, Cabbage Looper-Use 35 lbs per acre Apply

at 3 to 5 day intervals depending upon the severity

of the infestation and local conditions. Do not apply after bolls begin to open. Do not graze meat or dairy

GRAPES (7): Lafhoppers, Leatfolders (Calif)

Powdery Mildew-Use 30 to 35 lbs per acre. Do not

PEAS, SELD (0): Pea Aphids Pea Weevil-Use 35 This per acre. Do not feed treated vines or thre h

POTATOES (0): Aphids Colorado Potato Beetle

Stink Bugs, Plant Bugs (including Shield Shaped

Burs), Potato Flea Beetle, Potato Leafhopper, Po-

tat) Tuberworm Septhem Armyworm-Use 30 to 35

This per acre; European Corn Borer, Potato Psyllid-


Use 35 lbs per acre; False Chinch Bug-Use 35 lbs

per acre; Three-lined Potato Beetle, Use 17 lbs. per



SULFUR 50

FUNGICIDE-INSECTICIDE



ACTIVE INGREDIENTS: Sulfur

Sulfur fineness 98" thru 325 mesh

CAUTION KEEP OUT OF REACH OF CHILDREN.

skin, eyes or clothing. In case of contact with skin or eyes, flush with plenty of water; for eyes, get handling and before eating or smoking; wear clean clothing. Avoid contamination of feed and foodstuffs. Do not apply or allow to drift to areas oc-

Workers entering treated fields within 24 hours should wear protective clothing.



50 0%

The land of the second with the second se INERT INGREDIENTS: 47.0%

100.0%

*Thiodan, U.S. Pat. Ao. 2,799,685

Harmful it swallowed, inhaled or absorbed through skin. Avoid breathing of dust. Avoid contact with medical attention. Wash with soap and water after cupied by unprotected humans or beneficial animals.

NET WEIGHT LBS. **50**

USAGE CAUTION:

Applications of spraying or dusting materials containing sulfur should not be made just prior to or during periods of excessively high temperatures.

This product is toxic to fish and wildlife. Keep

out of any body of water. Apply this product only

as specified on this label

DISPOSE OF WASTES by burying in non-crop lands away from water supplies. Container should be disposed of by burying with wastes or by burning. Keep out of smoke

DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGH-BORING CROPS OR NON CEOP AREAS OR U.S. IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LAPIT DIRECTIONS BECAUSE ANIMAL, PLANT OF CHOP BILLING HILLEGAL RESIDUES OF OTHER UNDESERVED RESULT MAY OFFITR

Dealers Should Sell in Original Packages Only.

TERMS OF SALE OR USE. On purchase or use of this product torror and over a read to the tollowing a military

WARRANTY Ningara & irrants that this product, is of the fiif sale to Neagara 1 is intrinsic the ingredient databases of the Directions of User EXCLUDEAS SO WALRANDED THE PRODUCT IS GOLD AS BY NIAGARA MAKES NO CHIEF

WARRANTY LAPRESS OR IMPORTS PAMAGES - Burney reads as a second for trace to to correctly one of percent tak aparte berek sekrater, berek elektrick bereit and late are districtly to the property of the service of the serv by topics or a containing was to have be seen as a containing

to be with the open TACUSE FROM NON CONTROL (ABILE) COMMITTONS OF B regions of electromy estimates and the fitting of the artists of the electron for the second control of the con For the second of the which New and because of the Contract of the and a condition of purchase of war are more respected. and replace Niggary from Jacky's and my of it the back is or use of the product attributation of south care is a South care

authorized to vary or waise any statement contains I herein.

- - -

910B