

ACCEPTED
MAY 17 1967
UNDER THE FEDERAL INSECTICIDE ACT
279-1794 SUBJECT

CODE
537

Viagara®

NET CONTENTS
½ Pint
liquid

STIKCOL®

for control of PREHARVEST DROP OF APPLES

USDA Reg. No. 279-1794

ACTIVE INGREDIENT:

*Triethanolamine salt of silvex (2-(2,4,5-Trichlorophenoxy) propionic acid) 5.3%
INERT INGREDIENTS: 94.7%

100.0%

*Equivalent to 3.4% silvex (2-(2,4,5-Trichlorophenoxy) propionic acid) by weight or 7.5 grams per one-half pint.

WARNING

KEEP OUT OF REACH OF CHILDREN

Avoid contact with skin, eyes or clothing. Avoid spray drift to susceptible plants as this product may injure beans, peas, ornamentals, etc. Do not use the same spray equipment for other purposes unless thoroughly cleaned. Do not store near fertilizers, seeds, insecticides or fungicides. Do not contaminate water used for domestic or irrigation purposes.

fmc NIAGARA CHEMICAL DIVISION **fmc**
FMC CORPORATION
MIDDLEPORT, NEW YORK

DIRECTIONS FOR USE

The effectiveness of this spray to prevent apple drop depends upon good spray coverage and having foliage free from mite damage, magnesium deficiency, frost injury and severe scab infection.

McINTOSH: For Immediate Market. Apply about 7 to 8 days before harvest drop normally begins. Use ½ pint in 100 gallons of water (20 ppm). Pick fruit at proper maturity but do not let fruit over-ripen on trees. **Fruit to be Stored.** Apply spray 4 to 5 days before harvest drop normally begins. Use ¼ pint in 100 gallons of water (10 ppm). This should control drop for 2 weeks. Fruit should be picked before chemical has lost its effectiveness.

SUMMER VARIETIES: For varieties such as Early McIntosh, Melton and Wealthy, apply spray 2 weeks before harvest. Use ½ pint in 100 gallons of water. For Williams Early Red, use ¼ pint in 100 gallons of water and apply 2 weeks before harvest.

FALL VARIETIES: For Baldwin, Rome, Jonathan, Delicious, Spy, York and Staymen, apply spray about 2 weeks before harvest. Use ½ pint in 100 gallons of water.

USE IN SPEED SPRAYERS: This material has been used successfully in speed sprayers both in a concentrate or dilute form. For effective spraying of this chemical with a speed sprayer, contact your local agricultural authorities.

PRECAUTIONS

1. Time your sprays and use correct concentrations according to varieties involved and degree of ripeness desired.
2. Fruit should be harvested at proper maturity, but do not allow fruit to become over-ripe on trees before picking.
3. Do not use this material at a higher concentration than is recommended.
4. Do not use more than one application on a given variety.

Dealers Should Sell in Original Packages Only.

USAGE CAUTION: DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROP OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH DIRECTIONS, BECAUSE PLANT INJURY, EXCESSIVE RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR.

NOTICE OF DISCLAIMER: This product is sold without warranty, express or implied, except that it conforms to the label ingredient statement. Seller shall not be responsible for injury, damage or loss of any kind resulting from handling, storage or use of the product, regardless of the negligence or fault of Niagara or Seller. Buyer accepts and uses the product on these conditions.

056

Printed in U. S. A.