

ENVIRONMENTAL AND DANGER SAL CAUTIONS

This product is toxic to fish, birds and other wildlife. Birds and other wildlife in treated areas may be killed. Keep out of lakes, ponds and streams. Do not contaminate water by cleaning of equipment, or disposal of wastes and containers. Do not apply when weather conditions favor drift from the areas treated. Do not apply where run-off is likely to occur. Apply this product only as specified on the label.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

Do not re-use empty 2 gallon containers. Destroy them by perforating or crushing. Bury or discard in a safe place away from water supplies.

Do not store below 30° F. Material if frozen will thaw out without impairment on raising temperature.

Do not use in a manner or at a time other than in accordance with label directions because animal, plant or crop injury, illegal residues or other undesirable results may occur.

For Emergency Assistance Call: 716—735-3765

DIRECTIONS FOR USE

Apply when pests first appear and repeat as required. Add required amount of this material to partially filled spray tank. Continue to fill tank and keep agitator running. Do not allow mixture to stand. Do not combine with emulsifiable liquids unless previous use has proven them to be compatible. BEFORE USING SHAKE OR ROLL CONTAINER. Do not use in undiluted form. Observe days interval between last application and harvest indicated by number in () following the crop.

124W

MANUFACTURED BY STAUFFER CHEMICAL COMPANY

AGRICULTURAL CHEMICAL DIVISION

MIDDLEPORT, N.Y.
FRESNO, CALIF.

YAKIMA, WA
RICHMOND, CALIF.

**CODE:
30537**

**Net Contents
2 Gallons**

TRITHION[®] 4.0 FLOWABLE

INSECTICIDE-MITICIDE

EPA REGISTRATION NO. 279-1807 AA

ACTIVE INGREDIENT:	BY WT.
*S-[[p-Chlorophenyl)thio]methyl] 0,0-diethyl phosphorodithioate	44.00%
INERT INGREDIENTS:	56.00%
	100.00%

Contains 4 pounds per gallon of *Carbophenothion (Trithion)

WARNING

KEEP OUT OF REACH OF CHILDREN

May Be Fatal if Swallowed. Do not get in eyes, on skin or clothing. For commercial application, wear rubber gloves, protective clothing, rubber over-shoes and goggles. In case of contact, wash immediately with soap and water; for eyes flush with water for 15 minutes. Do not contaminate feed and foodstuffs. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash contaminated clothing with soap and hot water before re-use. Do not permit children or domestic animals on treated areas until it is completely dry. Trithion upon use, may cause cholinesterase inhibition. ATROPINE IS ANTIDOTAL. Commercial operators should maintain a supply of Atropine 1/100 grain tablets.

TRITHION is a registered trademark of Stauffer Chemical Company for an insecticide-miticide.

Use Trithion in a thorough manner. Do not use if adjusted in minimum of

ALMONDS: Spider) — Use if almond harvest

APPLES (30) (Apple, Rose Twospotted) not apply material

APRICOTS mer: Mites

CITRUS: Mite Spider) — Use between application California Red, low — Use days of harvest to kumquat summer or fall

FIGS (7): Two

OLIVES (60) pint plus 1 to

PLUMS (30) (Brown, European) than 4 pints

POST-HARVEST: Apple, Apricot, Quince): To Spider Mites

Dosage rate: sufficient water

Net Contents
2 Gallons
FLOWABLE

BY WT.

44.00%

56.00%

100.00%

Trithion)

LDREN

othing. For com-
ng, rubber cover-
y with soap and
contaminate feed
soap and water
th soap and hot
imals on treated
se cholinesterase
rs should main-

Company for an

FRUIT AND NUT CROF

Use Trithion 4.0 Flowable in amounts given below per 100 gallons of water in a thorough coverage spray in normal spray-schedule. If concentrate sprays are used the amount of Trithion 4.0 Flowable per 100 gallons should be adjusted in order to apply the same quantity of Trithion per acre. Apply a minimum of 1 quart per acre.

ALMONDS: Summer Sprays: Mites (Brown, European Red, Twospotted Spider) — Use 1 to 2 quarts in a minimum of 50 gallons of water per acre. If almond hulls are to be used for feed, do not apply within 7 days of harvest.

APPLES (30), CRABAPPLES (30), PEARS (30), QUINCE (30): Summer: Aphids (Apple, Rosy Apple, Wooly Apple), Leafhopper, Mites (Brown, European Red, Twospotted Spider) — Use ½ pint. Apply only after full leaf expansion. Do not apply more than 2½ pints per acre.

APRICOTS (30), CHERRIES (30), PEACHES (30), NECTARINES (30): Summer: Mites (Brown, European Red, Twospotted Spider) — Use ½ pint.

CITRUS: Mites (Citrus Red, Citrus Rust, Sixspotted, Texas Citrus, Yuma Spider) — Use 1 pint. Do not apply within 14 days of harvest. Allow 30 days between applications. Do not use more than 7½ pints per acre. Scales (California Red, Citricola, Cottoncushion, Florida Red, Purple, Brown Soft, Yellow) — Use 1 pint plus 1 to 2 gallons spray oil. Do not apply within 30 days of harvest. Do not use more than 10 pints per acre. DO NOT APPLY to kumquat or citron. Trithion may cause injury to grapefruit if used in summer or fall before the fruit is fully colored.

FIGS (7): Twospotted Spider Mites — Use ½ to 1 pint.

OLIVES (60): Scales (Black, Oleander, Olive, San Jose) — Use ½ to 1 pint plus 1 to 2 gallons of spray oil.

PLUMS (30), PRUNES (30): Summer: Leaf Curl Aphid, Leafhoppers, Mites (Brown, European Red, Twospotted) — Use ½ to 1 pint. Do not apply more than 4 pints per acre.

POST-HARVEST APPLICATION TO DECIDUOUS FRUIT TREES: (Almond, Apple, Apricot, Cherry, Crabapple, Nectarine, Peach, Pear, Plum, Prune, and Quince): To decrease overwintering Population of Aphids, Leafhoppers, Spider Mites — Use ½ to 1 pint.

VEGETABLE, VINE AND FRUIT CROPS

Dosage rates are given in terms of TRITHION 4.0 FLOWABLE per acre in sufficient water to provide good coverage.

BEANS (Suculent) (Snap and Lima) (7): Bean Aphid, Potato Leafhopper, Serpentine Leafminer, Twospotted Spider Mites — Use 1 to 1½ pints. Do not graze or feed crop residue to livestock.

SUGAR BEETS (14): Bean Aphid, Potato Flea Beetle, Serpentine Leafminer, Twospotted Spider Mites — Use 1 pint.

CANTALOUPE (5), WATERMELON (5): Melon Aphid, Potato Flea Beetle, Serpentine Leafminer, Twospotted Spider Mites — Use 1 to 2 pints.

COTTON: Cotton Aphid, Fleahopper, Cotton Leafworm, Cotton Leafperforator, Mites (Cinnabar, Tropical Spider, Twospotted Spider) — Use 1 to 2 pints. Apply 1 quart per acre for control of Cotton Aphids and Cotton Leafperforator. Treated plants should not be grazed or fed to livestock.

CUCUMBER (7), SUMMER SQUASH (7): Melon Aphid, Potato Leafhopper, Serpentine Leafminer, Twospotted Mites — Use 1 to 2 pints.

EGGPLANT (7): Mites (Tropical Spider, Twospotted Spider), Potato Flea Beetle, Potato Leafhopper, Serpentine Leafminer — Use 1 to 2 pints.

ALFALFA, CLOVER: Pea Aphid, Potato Leafhopper, Twospotted Spider Mite, Spotted Alfalfa Aphid — Use 1 to 2 pints per acre. Do not make more than 1 application per season. Do not cut or graze for hay within 28 days after application.

BEETS, GARDEN: Bean Aphid, Potato Flea Beetle, Serpentine Leafminer, Twospotted Spider Mites — Use ½ to 1 pint. Do not apply within 21 days of harvest if tops are to be used for food or feed.

GRAPES (30): Grape Leafhopper, Twospotted Spider Mites — Use 1 pint per acre in sufficient water for coverage. Do not apply to sensitive varieties such as Charbono, Gamay and Muscat.

Dealers Should Sell in Original Packages Only.

WARRANTY: FMC warrants that this product, as of the time of sale by FMC, (1) conforms to the ingredient statement on the label, and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS. FMC MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

EPA EST., 279-CA-2

Printed in U.S.A.