

Code: 8

Net Contents
5 Gallons

SINOX-W

GENERAL INFORMATION

MIXING INSTRUCTIONS: Sinox-W is a water soluble material which dissolves readily with mild agitation

WEEDS CONTROLLED: Sinox-W can be used effectively as a post emergence (selective) treatment for the control of such broadleaved weeds as annual mustard, annual thistles, chickweed, lambsquarters, pigweed, smartweed, ragweed, fan or Fench weed, shepherd's purse, wild lettuce, wild radish, yellow rocket and annual seedling grasses. Generally higher rates of application are needed for seedling grass control than for seedling broadleaved weed control. **Apply Sinox-W when weeds are small, young, succulent and actively growing. Established perennial weeds are not usually controlled.**

APPLICATION PROCEDURES: Apply during warm, sunny weather when crop foliage is dry. Under conditions of cool weather or drought only partial control may be obtained. When temperatures are 70° to 85° F less Sinox-W is needed to give good control. Use enough spray solution and maintain adequate pressure to give complete and uniform coverage of all weed foliage. For ground rig application use 40 to 60 gallons of spray mix per acre and for airplane application 15 gallons per acre. Pressure may vary from 25 to 75 psi depending on equipment used. A flat fan-shaped spray pattern is recommended.

USAGE CAUTIONS: Sinox-W may produce some foliage burn of crop leaves but this effect is usually only temporary. Do not apply when air temperatures are 85° F or above or expected to be during the 24 hour period following application. During periods of high humidity, or when soil moisture is high resulting in high humidity close to the soil surface, make certain that the crop plant has a good coat of wax on the leaves before spraying. Do not spray just before a rain.

DIRECTIONS FOR USE

ALFALFA, and ALSIKE, LADINO and RED CLOVERS: Apply 1/2 to 1 gallon per acre by ground rig or airplane when seedlings have two or more trifoliolate leaves. Do not graze treated areas or feed hay to livestock from treated fields within 40 days of treatment.

CORN (Field, Pop, Sweet): 1 to 1 1/2 gallons per acre while corn is in tight roll stage. Apply by ground.

GARLIC, ONIONS: Apply 1 to 1 1/2 gallons per acre after two leaf stage and before crop is 4 to 6 inches tall. Use air or ground application. **Do not apply Sinox-W on onions which are to be used as green table onions.** The following varieties of onions are not injured by the recommended rates of Sinox-W: White Globe, Yellow Globe, Australian Brown, Italian Red and Sweet Spanish. Any other variety should be tested before field scale applications are made. **NOTE:** When using in inland areas use the lower dosages and do not apply if temperatures exceed 75° F.

SELECTIVE WEED KILLER

EPA REGISTRATION NO. 279-1841 AA

ACTIVE INGREDIENT:
Amr onium salt of dinoseb (2-sec-butyl-4,6-dinitrophenol) 13%
INERT INGREDIENTS: 87%
100%

Contains the equivalent of 1 pound DNOSBP per gallon
Contains Nitrated Phenols.



DANGER - POISON



KEEP OUT OF REACH OF CHILDREN

May be Fatal If Swallowed or Absorbed Through Skin! Poisonous if Swallowed or Absorbed Through Skin! Do not breathe spray mist. Do not get in eyes, on skin or on clothing. In case of contact, remove clothing and wash thoroughly with soap and water; for eyes, flush with water and get medical attention. Keep away from domestic animals and foodstuffs.

FIRST AID

INTERNAL: To induce vomiting give 2 tablespoonful of salt in a glass of warm water. Repeat until vomiting fluid is clear. **CALL A PHYSICIAN IMMEDIATELY!**

EXTERNAL: Wash with rubbing alcohol and follow with a thorough scrubbing with soap and water. Remove contaminated shoes and clothing immediately. If splashed in eyes, immediately flush eyes with plenty of water for 15 minutes and get medical attention.

NOTE TO PHYSICIAN: Active ingredient is metabolic stimulant. Treat symptomatically. **THERE IS NO SPECIFIC ANTIDOTE.**

NOT FOR USE OR STORAGE IN OR AROUND THE HOME

WARNING: FLAMMABLE! KEEP AWAY FROM HEAT AND OPEN FLAME. Protect from temperature below 15° F.

ENVIRONMENTAL AND DISPOSAL CAUTIONS

This product is toxic to fish, wildlife and birds. Birds and other wildlife in treated areas may be killed. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from the areas treated. Do not apply where run-off is likely to occur. Apply only as specified on this label.

Do not re-use empty 5 gallon container. Destroy it by perforating or crushing. Bury in a safe place away from water supplies.

PEAS: 1/2 to 3/4 gallon per acre applied when peas are 2 to 8 inches tall. **Apply before pea flower buds appear.** Allow enough time for plants to harden following cloudy weather or periods of high humidity **before spraying.** Apply with ground rig or airplane. Do not graze treated areas or feed hay to livestock from treated fields within 40 days after treatment.

SMALL GRAINS (BARLEY, OATS, WHEAT) WITH LEGUME COMPANION CROP (ALFALFA and ALSIKE, LADINO and RED CLOVERS): Apply 3/4 to 1 gallon per acre when grain is 3 to 6 inches tall and legume crop plants have two or more trifoliolate leaves. Make sure crop foliage is dry at time of application. Apply from ground or air equipment. Do not graze treated areas or feed hay to livestock from treated fields within 60 days after treatment. **NOTE, ALL CROPS:** Sprayed areas should not be disturbed by cultivation or any other means until weeds become troublesome.

GLADIOLUS: Use 3/4 gallon per acre and apply when leaves are still short and straight. May be applied after plants become tall if a low pressure fan spray is directed at the base of the plants and tops of leaves are not wetted.

DO NOT USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS BECAUSE ANIMAL, PLANT OR CROP INJURY, ILLEGAL RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR.

Dealers Should Sell in Original Packages Only.

TERMS OF SALE OR USE: On purchase or use of this product buyer and user agree to the following conditions:

WARRANTY: FMC warrants that this product, as of the time of sale by FMC, (1) conforms to the ingredient statement on the label, and (2) is reasonably fit for the purposes set forth in the Directions for Use. **EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS. FMC MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.**

DAMAGES: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages. Notice of any claim must be reported to FMC within 10 days of discovery by buyer or user, failing which buyer or user waives any claim for such damage.

EXCUSE FROM NON-CONTROLLABLE CONDITIONS: Because of certain critical conditions created or incurred by buyer or user or over which FMC has no control, buyer and user, as a condition of purchase or use, assume responsibility for and release FMC from liability arising out of the handling or use of the product attributable to such causes. No person is authorized to vary or waive any statement contained herein.


104W

MANUFACTURED IN USA BY

AGRICULTURAL CHEMICAL DIVISION

MIDDLEPORT, N.Y. YAKIMA, WA
FRESNO, CA RICHMOND, CA

FMC



EPA EST. 279-CA-2 Printed in U.S.A.