

Code: 667

50 Pounds Net Weight

TRITHION[®] 3 DUST

DIRECTIONS FOR USE

Apply when pests first appear and repeat as necessary. May be applied with ground or airplane application equipment. Ten-day interval between last application and harvest indicated by number in parentheses following the crop.

ALFALFA: Two-Spotted Mites, Pea Aphid, Spotted Alfalfa Aphid, Potato Leafhopper, Meadow Spittlebug. Use 20-33 pounds per acre. Meadow Spittlebug. Use 33 pounds per acre. Do not apply more than one application per cutting. Do not graze or cut hay within 28 days after application.

ALMOND, APRICOT, CHERRY, NECTARINE, PEACH, PLUM, PRUNE: Aphids, Orchard Mites. Use 40-60 pounds per acre. If almond hulls are used as feed do not apply within 7 days of harvest. Do not apply to apricot, cherry, nectarine, peach, plum, and prune within 30 days of harvest.

APPLE (30), CRABAPPLE (30), PEAR (30), QUINCE (30): Apple Aphid, European Red Mite, Red Apple Aphid, Woolly Apple Aphid, Two-Spotted Mites, Codling Moth. Use 40 pounds per acre.

BEETS, GARDEN: Bean Aphid, Leafhoppers, Serpentine Leafminer, Two-Spotted Mites, Potato Flea Beetle. Use 15-40 pounds per acre. Do not apply within 21 days of harvest. Tops are to be used in feed or forage.

CANTALOUPE (5), WATERMELON (5): Melon Aphid, Serpentine Leafminer, Potato Flea Beetle, Two-Spotted Mites. Use 20-40 pounds per acre.

CITRUS (14): Citrus Red Mite, Citrus Rust Mite, Six-Spotted Mite, Texas Citrus Mite, Yema Mite. Use 35-100 pounds per acre. Allow thirty days between applications. Do not apply to lemons or limes.

CLOVER: Two-Spotted Mite, Pea Aphid, Potato Leafhopper, Meadow Spittlebug. Use 20-33 pounds per acre. Use 32 pounds per acre for the control of the Meadow Spittlebug. Do not apply more than one application per cutting. Do not graze or cut hay within 28 days after application.

CORN (FIELD AND SWEET): Grass Mite, Pacific Mite, Two-Spotted Mite. Use 33 pounds per acre. Do not make more than one application per season. Do not graze or cut corn forage within 21 days after application.

COTTON: Cotton Aphid, Cotton Leaf Perforator, Cotton Fleahopper, Cotton Flea Beetle, Cotton Mite, Tobacco Mite, Two-Spotted Mite. Use 20-30 pounds per acre. Apply 30 pounds per acre for Cotton Leaf Perforator. Do not graze dairy animals or meat animals in treated areas.

CUCUMBER (7), SUMMER SQUASH (7): Mite, Aphid, Potato Leafhopper, Serpentine Leafminer, Two-Spotted Mite. Use 20-30 pounds per acre.

EGGPLANT (7): Mite, Aphid, Potato Leafhopper, Potato Flea Beetle, Triplicorn Mite, Two-Spotted Mite. Use 20-30 pounds per acre.

INSECTICIDE - MITICIDE

EPA REGISTRATION NO. 279-2035

ACTIVE INGREDIENT	BY WT
Carbophenothion Salt, Chlorophenyltin dimethyl 0,0-diethyl phosphorodithioate	3.90%
INERT INGREDIENTS	97.00%
	100.00%

Trithion is a registered trademark of Stauffer Chemical Company, for an insecticide. U.S. Patent No. 2,793,224

CAUTION

KEEP OUT OF REACH OF CHILDREN

Hazardous if Swallowed. Avoid breathing dust. Do not get in eyes, on skin, or on clothing. Wear dust respirator and goggles. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Do not breathe fumes and food stuffs. Wash all contaminated clothing with soap and hot water before reuse. Do not store near food and food products.

Inhalation upon repeated, prolonged or careless use may cause dizziness, headache, and nausea. Keep supply of Atropine 1:1000 for treatment of anticholinergic poisoning.

ACCEPTED

MAR 12 1970

U.S. DEPARTMENT OF AGRICULTURE
 OFFICE OF PLANT INDUSTRY
 WASHINGTON, D.C. 20250

FIGS (7): Two-Spotted Mite. Use 30-40 pounds per acre.

GRAPES (30): Grape Leafhopper, Two-Spotted Mites. Use 20-33 pounds per acre.

ONIONS (GREEN OR DRY BULB): Onion Maggot. Use 33-66 pounds per acre, depending on soil types. Apply at time of seeding in the planting furrow, using a combination of broadcast and sprayer methods, or at application equipment.

ORNAMENTALS: Citrus Red Mite, Citrus Rust Mite, Violets, Fungus Gnats, Destructive Shrubs, Spider Mites, Aphids. Use 15-25 pounds per acre.

PEAS (7): Pea Aphid, Serpentine Leafminer, Two-Spotted Mite. Use 20-40 pounds per acre. Do not graze or feed crop residue to livestock.

PEPPER (7), PIMENTO (7): Serpentine Leafminer, Potato Flea Beetle, Potato Leafhopper, Tropical Mite, Two-Spotted Mite. Use 20-40 pounds per acre.

SNAP BEANS, LIMA BEANS, SOYBEAN (IN POD): Bean Aphid, Mexican Bean Beetle, Serpentine Leafminer, Potato Leafhopper, Two-Spotted Mites. Use 20-40 pounds per acre. On Soybeans do not apply within 7 days of harvest. Do not feed treated forage to livestock. On Beans at the 20 pound per acre rate do not apply within 7 days of harvest. Do not apply more than twice per season. Do not graze or cut vire for hay within 21 days after application. At a rate above 20 pounds per acre, do not apply within 7 days of harvest. Do not feed treated forage to livestock.

SORGHUM: Two-Spotted Mite. Use 33 pounds per acre. Do not make more than one application per season. Do not graze or cut forage within 21 days after application.

SPINACH (21), BEET TOPS (21): Serpentine Leafminer, Two-Spotted Mite, Bean Aphid. Use 15-40 pounds per acre.

STRAWBERRY (3): Two-Spotted Mites, Strawberry Aphid. Use 20-40 pounds per acre.

SUGAR BEETS (14): Bean Aphid, Serpentine Leafminer, Potato Flea Beetle, Two-Spotted Mite. Use 20-30 pounds per acre.

TOMATOES (7): Serpentine Leafminer, Potato Psyllid, Potato Flea Beetle, Potato Leafhopper, Tobacco Mite, Tropical Mite, Two-Spotted Mites. Use 20-40 pounds per acre.

USAGE CAUTIONS


This product is toxic to fish and wildlife. Keep out of lakes, ponds, and streams. Birds fed treated forage may be killed. Do not apply when weather conditions favor drift from areas treated. Do not apply to any area not specified on the label.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

Do not use (WASH) treated forage for crop animals from water supply. Empty troughs and buckets with water or beered. Keep out of sinks.

Printed in U.S.A.

ADDITIONAL INFORMATION APPEARING ON BAGS TO WHICH THIS LABEL WILL BE ATTACHED.



Dealers Should Sell in Original Packages Only.

DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROPS OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS BECAUSE ANIMAL, PLANT OR CROP INJURY, ILLEGAL RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR.

TERMS OF SALE OR USE - On purchase or use of this product buyer and user agree to the following conditions:

WARRANTY - Niagara warrants that this product, as of the time of sale by Niagara, is conforming to the ingredient statement on the label and is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS. NIAGARA MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

DAMAGES - Dealer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to repair or replacement of the product. Damages shall not include consequential damages. Notice of any claim must be reported to Niagara within 90 days of purchase by buyer or user, failing which buyer or user waives and releases Niagara from liability for such damage.

EXCUSE FROM NON-CONTROLLABLE CONDITIONS - Because of certain critical conditions created or induced by buyer or user or over which Niagara has no control, buyer and user, as a condition of purchase, release Niagara's responsibility for and release Niagara from liability arising out of the handling or use of the product attributable to such causes. No person is authorized to vary or waive any statement contained herein.

MANUFACTURED IN U.S.A. BY



NIAGARA CHEMICAL DIVISION
FMC CORPORATION
MIDDLEPORT, NEW YORK

