

CODE 744

2 1/2 LBS. NET WT.

TEDION[®] 50

WETTABLE POWDER

MITICIDE

EPA Reg. No. 279-2127

ACTIVE INGREDIENT:

*Tetradifon (4-chlorophenyl 2,4,5-trichlorophenyl sulfone)	50.0%
INERT INGREDIENTS:	50.0%
	100.0%

*Tedion, U. S. Pat. No. 2,812,281

CAUTION

KEEP OUT OF REACH OF CHILDREN

This product is toxic to shrimp, crayfish, crabs and other aquatic invertebrates. Keep out of lakes, ponds, streams, marshes and tidal estuaries. Apply this product only as specified on this label.

Rinse equipment and DISPOSE OF WASTES by burying in non-crop lands away from water supplies. Container should be disposed of by burying with wastes or by burning. Keep out of smoke.

DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROPS OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS BECAUSE ANIMAL, PLANT OR CROP INJURY, ILLEGAL RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR.

DEALERS SHOULD SELL IN ORIGINAL PACKAGES ONLY


TERMS OF SALE OR USE: On purchase or use of this product buyer and user agree to the following conditions:

WARRANTY: Niagara warrants that this product, as of the time of sale by Niagara, (1) conforms to the label statement, (2) the label and (3) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED, THE PRODUCT IS SOLD AS IS. NIAGARA MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

DAMAGES: Buyer's remedy is exclusive and limited to replacement of the product if it is found to be damaged. Damages for the use of the product are limited to the amount of the purchase price. Niagara Chemical Division, Niagara, New York, is not liable for any damages, including consequential damages, arising from the use of this product.

EXCUSE FROM NON-CONTROLLABLE CONDITIONS: Because of certain critical conditions beyond the control of Niagara Chemical Division, the use of this product may be limited. Niagara Chemical Division is not liable for any damages, including consequential damages, arising from the use of this product attributable to such conditions. Niagara Chemical Division's liability is limited to the amount of the purchase price of the product.

TEDION[®] Registered trademark of N. V. Phillips Roxane



DIRECTIONS FOR USE

DO NOT USE IN UNDILUTED FORM

FRUITS

Kill of adult mites is not rapid and several days may elapse before full effect of this material is noted. However, Tedion 50 Wettable Powder has long residual activity and should remain effective for several weeks. Observe days interval between last application and harvest indicated by number in parentheses following crop.

APPLE (0), CRABAPPLE (0), PEAR (0) QUINCE (0): European Red Mite, Two-spotted Spider Mite, McDaniel Mite, Canadensis Mite, Schoenherr Mite—Use 1/2 lb. per 100 gals. Spray trees thoroughly. Apply at pink and petal fall and during cover sprays as necessary to maintain control. Do not make more than 4 applications after petal fall per season and not more than 500 gals. of spray per acre per application. Do not feed pomace from treated apples or crabapples to livestock.

APRICOT (0), CHERRY (0), NECTARINE (0), PEACH (0), PLUM (0), PRUNE (0): European Red Mite, McDaniel Mite, Two-spotted Spider Mite—Use 1/2 lb. per 100 gals of water immediately after bloom. Apply during cover sprays as necessary to maintain control. Use 1/2 lb. per 100 gals. of water in spray.

Do not make more than two applications to apricot, cherry, nectarine, peach or three applications to plums or prunes.

RUSHBERRIES (Blackberries, Boysenberries, Blueberries, Gooseberries, Loganberries, Raspberries): Two-spotted Mite—Use 2 lbs. per acre. APPLY AS A POST-HARVEST APPLICATION OR WHEN NO FRUIT IS PRESENT.

CITRUS CITRUS (0), GRAPEFRUIT (0), LEMONS (0), LIMES (0), ORANGES (0), TANGELOS (0), TANGERINES (0): Citrus Red Mite, Six-spotted Mite, Texas Citrus Mite—Use 1/2 lb. per 100 gals. of water, 4 to 5 lbs. per acre. DO NOT APPLY MORE THAN ONCE PER SEASON WHILE FRUIT IS PRESENT.

GRAPES (0): Pacific Mite, Two-spotted Mite, Willamette Mite—Use 2 lbs. per acre. Do not make more than 3 applications during fruiting period.

FIGS (14): Pacific Mite, Two-spotted Mite—Use 2 1/2 to 5 lbs. per acre. Do not make more than 2 applications during the fruiting period.

NURSERY STOCK (NON-BEARING TREES): Citrus Red Mite, European Red Mite, Two-spotted Mite—Use 1/2 lb. per 100 gallons of water. Spray thoroughly when mites appear and repeat as necessary. Thorough coverage of the top and underside of foliage is essential for effective mite control.

ROW CROPS

Apply when mites first appear and repeat as required to maintain control. Unless otherwise stated, apply in sufficient water for thorough coverage. Spray plants thoroughly. If mite control is required beyond the following use pattern, supplement the mite control program with other suitable miticides. Observe days interval between last application and harvest indicated by number in parentheses following crop.

CUCUMBERS (0): Two-spotted Mite—Use 1 lb. per acre for field use, (1/2 lb. per 100 gal. for greenhouse use.) Do not apply more than 2 applications during the fruiting season.

MELONS (0): Two-spotted Mite—Use 1 lb. per acre.

PUMPKINS (0): Two-spotted Mite—Use 1 lb. per acre.

SQUASH, WINTER (0): Two-spotted Mite—Use 1 lb. per acre.

STRAWBERRIES (3): Two-spotted Mite—Use 2 lbs. per acre. During the fruiting period, make 3 applications at intervals less than 25 days.

TOMATOES (0): Tropical Mite, Two-spotted Mite—Use 1/2 lb. per acre. Make 2 applications during the fruiting period.

EVERGREENS: Spruce Mite—Use 1/2 lb. per 100 gals. of water.

HOPS (10): Two-spotted Mite—Use 1/2 lb. per 100 gal. of water per acre. Do not make more than 10 applications during the fruiting period.

ORNAMENTALS (0): Two-spotted Mite—Use 1/2 lb. per 100 gals. of water. Apply in sufficient water for thorough coverage. Do not apply more than 2 applications during the fruiting period.

As a complete fumigant for soil-borne mites, Tedion 50 Wettable Powder can be used in conjunction with steam heating. Apply to the soil in a 1/2 inch layer, 2 to 3 inches deep, and cover with a 1/2 inch layer of soil. Use a 1/2 inch layer of soil, and cover with a 1/2 inch layer of soil. Use a 1/2 inch layer of soil, and cover with a 1/2 inch layer of soil.

At quitting time turn steam into treated pipes, leaving vents closed all night. Avoid soil temperatures above 100°F. Do not use Tedion 50 Wettable Powder in conjunction with steam heating. Make six treatments at 10 to 15 day intervals. Do not use Tedion 50 Wettable Powder in conjunction with steam heating. Make six treatments at 10 to 15 day intervals. Do not use Tedion 50 Wettable Powder in conjunction with steam heating.

Make six treatments at 10 to 15 day intervals. Do not use Tedion 50 Wettable Powder in conjunction with steam heating. Make six treatments at 10 to 15 day intervals. Do not use Tedion 50 Wettable Powder in conjunction with steam heating.

Make six treatments at 10 to 15 day intervals. Do not use Tedion 50 Wettable Powder in conjunction with steam heating. Make six treatments at 10 to 15 day intervals. Do not use Tedion 50 Wettable Powder in conjunction with steam heating.

Make six treatments at 10 to 15 day intervals. Do not use Tedion 50 Wettable Powder in conjunction with steam heating. Make six treatments at 10 to 15 day intervals. Do not use Tedion 50 Wettable Powder in conjunction with steam heating.

Make six treatments at 10 to 15 day intervals. Do not use Tedion 50 Wettable Powder in conjunction with steam heating. Make six treatments at 10 to 15 day intervals. Do not use Tedion 50 Wettable Powder in conjunction with steam heating.

Make six treatments at 10 to 15 day intervals. Do not use Tedion 50 Wettable Powder in conjunction with steam heating. Make six treatments at 10 to 15 day intervals. Do not use Tedion 50 Wettable Powder in conjunction with steam heating.

Make six treatments at 10 to 15 day intervals. Do not use Tedion 50 Wettable Powder in conjunction with steam heating. Make six treatments at 10 to 15 day intervals. Do not use Tedion 50 Wettable Powder in conjunction with steam heating.