

PELLETED SNAIL BAIT

DIRECTIONS FOR USE

① **BUSH AND VINE FRUITS** (Blackberries, Raspberries, Loganberries, Raspberries, Strawberries), **NURSERIES**, **ORNAMENTAL GREENHOUSES**, **VEGETABLE CROPS** (Beans, Broccoli, Brussels Sprouts, Cabbage, Carrots, Cauliflower, Celery, Collards, Corn, Cucumbers, Eggplant, Kale, Kohlrabi, Melons, Peppers, Pumpkins, Rutabagas, Spinach, Squash, Tomatoes, Turnips): To control Slugs, Snails and Cutworms use 100-300 pounds per acre. (2½-7½ pounds per 1000 square feet). Apply to soil surfaces around plants. If the ground is dry, sprinkle lightly with water. Repeat as necessary. Do not apply to foliage. Do not apply after edible parts start to form.

1-5 **AVOCADOS, GRAPEFRUIT, LEMONS, LIMES, ORANGES, TANGELOS AND TANGERINES**: To control Slugs, Snails and Cutworms use up to 300 pounds per acre as a broadcast under trees. Do not apply to tree foliage, other plant parts, or edible parts.

② **STRAWBERRY ROOT WEEVIL** — Place 3 to 5 pellets around the base of each strawberry plant, and around azaleas, camellias, rhododendrons and primroses. Apply when 75% to 90% of pupae change to adult stage in soil (about April or May). Do not apply to foliage or other plants parts. Do not contaminate edible parts. Do not apply more than 120 pounds per acre.

INSECTICIDE

EPA REGISTRATION NO. 279-2302 AA

| | |
|---------------------------------|----------------|
| ACTIVE INGREDIENTS: | BY WT. |
| Calcium Arsenate | 5.00% |
| Metaldahyde | 3.15% |
| INERT INGREDIENTS: | 91.85% |
| | 100.00% |

Total arsenic expressed as metallic1.88%
 Water soluble arsenic (As) max.1.25%

 **DANGER - POISON** 
KEEP OUT OF REACH OF CHILDREN

DANGER: Avoid breathing dust. Poisonous If Swallowed. Avoid contact with skin, eyes or clothing. Avoid contaminating feed and foodstuffs. Keep away from domestic animals. Field workers should use protective gloves when handling and broadcasting Pelleted Snail Bait. Wash thoroughly with soap and water after handling.

This pesticide may be fatal to children, and dogs or other pets if eaten. Keep children and pets out of treated area.

FIRST AID TREATMENT

Antidote For Arsenic: Give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Then give 2 tablespoonfuls of Epsom salt or Milk of Magnesia in water, and plenty of milk and water. Call A Physician Immediately! Have victim lie down and keep quiet.

USAGE CAUTIONS

Not for use or storage in or around the home.

Do not apply on food crops or directly to foliage or against the base of plants. Injury may result from bait becoming lodged in plant crevices.

Birds and other wildlife feeding on treated areas may be killed. Keep out of lakes, ponds and streams. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on the label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not apply when weather conditions favor drift from the areas treated. Do not apply when run-off is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes.

DISPOSE OF WASTES by burying in non-crop land away from water supplies. Empty bags should be buried with wastes or burned. Keep out of smoke.

For Emergency Assistance Call
716-735-3765


279-2302

023W Revised



Printed in U.S.A.

ADDITIONAL INFORMATION APPEARING ON BAGS TO WHICH THIS LABEL WILL BE ATTACHED.



Dealers Should Sell in Original Packages Only.

DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROPS OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS BECAUSE ANIMAL, PLANT OR CROP INJURY, ILLEGAL RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR.

TERMS OF SALE OR USE: On purchase or use of this product buyer and user agree to the following conditions:

WARRANTY: FMC warrants that this product, as of the time of sale by FMC, (1) conforms to the ingredient statement on the label, and (2) is reasonably fit for the purposes set forth in the Directions for Use. **EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS. FMC MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.**

DAMAGES: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages. Notice of any claim must be reported to FMC within 10 days of discovery by buyer or user, failing which buyer or user waives any claim for such damage.

EXCUSE FROM NON-CONTROLLABLE CONDITIONS: Because of certain critical conditions created or incurred by buyer or user or over which FMC has no control, buyer and user, as a condition of purchase or use, assume responsibility for and release FMC from, liability arising out of the handling or use of the product attributable to such causes. No person is authorized to vary or waive any statement contained herein.

Manufactured in U.S.A. by:
FMC CORPORATION
AGRICULTURAL CHEMICAL DIVISION
Middleport, N.Y. 14105

FMC
Chemicals