

Code: 30830

Net Contents
Gallons

TEDION®* E.C.

Tedion® E.C. is most effective against hatching Mite Eggs and young Mites. Kill of adult Mites is not rapid and several days may elapse before full effect is noted, however, Tedion® E.C. has long residual activity.

DIRECTIONS FOR USE

Apply when Mites first appear and repeat as necessary. Use the following recommended rates in sufficient water to thoroughly cover one acre. By ground, apply a minimum of 200 gallons per acre on stone fruits and nuts; a minimum of 300 gallons per acre on pome fruits; a minimum of 200 gallons per acre on bush and vine fruits and a minimum of 10 gallons per acre on vegetable and row crops. By air, apply a minimum of 10 gallons per acre on vegetable and field crops and a minimum of 20 gallons per acre on small fruits and tree crops. Observe days interval between application and harvest indicated by number in () following the crop.

APPLES (0), CRAB APPLES (0), PEARS (0), QUINCE (0) - West Coast: European Red Mite, Twospotted Spider Mite, McDaniel Spider Mite, Canadensis Mite, Schoene Spider Mite - Use 5 to 10 quarts per acre. Do not apply more than 3 times after petal fall. Do not feed treated apple and crab apple pomace to livestock.

APRICOTS (0), CHERRIES (0), NECTARINES (0), PEACHES (0): European Red Mite, Twospotted Spider Mite, McDaniel Spider Mite - Use 5 to 8 quarts per acre. Apply when a majority of overwintering Mite Eggs have hatched and repeat as necessary to maintain control. Do not apply more than once after shuck-split.

CITRUS (0), GRAPEFRUIT (0), LEMONS (0), LIMES (0), ORANGES (0), TANGELOES (0), TANGERINES (0): Citrus Red Mite, Sixspotted Mite, Texas Citrus Mite - Use 5 to 10 quarts per acre in sufficient water to thoroughly cover the fruit and foliage. Apply only once per season. Pulp from citrus treated in accordance with label directions may be used in the preparation of feed.

CUCUMBERS (0): Twospotted Spider Mites - Use 2 quarts per acre. Do not apply more than 3 times during the fruiting period.

0647


MANUFACTURED IN U.S.A. BY

FMC CORPORATION

AGRICULTURAL CHEMICAL DIVISION

MIDDLEPORT, N.Y.
FRESNO, CA

YAKIMA, WA
RICHMOND, VA



MITICIDE

EPA REGISTRATION NO. 279-2387 AA

ACTIVE INGREDIENTS:

*Tetradifon; 4-chlorophenyl 2,4,5-trichlorophenyl sulfone

Petroleum Hydrocarbons

INERT INGREDIENTS:

*Tedion - U. S. Patent No. 2,812,281
Contains 1.0 pound of Tedion per gallon.

BY WT.

12.30%

32.50%

55.20%

100.00%

CAUTION

KEEP OUT OF REACH OF CHILDREN

Avoid breathing of vapor. Avoid contact with skin. Harmful if swallowed. Use with adequate ventilation.

DO NOT STORE BELOW 35° F. Do not use, pour, spill or store near heat or open flame.

Do not use in a manner or at a time other than in accordance with label directions because animal, plant or crop injury, illegal residues or other undesirable results may occur.

For Emergency Assistance Call
716-735-3765

Dealer Should Sell in Original Packages Only.

WARRANTY: FMC warrants that this product, as of the time of sale by FMC, conforms to the ingredient statement on the label, and is reasonably fit for the purposes set forth in the Directions for Use EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS. MAINTAIN OTHER WARRANTY EXPRESS OR IMPLIED.

GRAPES (0): Pacific Spider Mite, Twospotted Spider Mite, Willamette Mite - Use 4 quarts per acre. Do not apply more than 3 times during the fruiting period.

HOPS (14): Twospotted Spider Mite - Use 4 quarts per acre. Do not apply more than twice during the fruiting period. Do not feed treated straw to livestock.

MELONS (0): Twospotted Spider Mite - Use 2 quarts per acre.

PLUMS (0), PRUNES (0): European Red Mite, Twospotted Spider Mite, McDaniel Spider Mite - Use 5 to 10 quarts per acre. Apply when a majority of overwintering Mite Eggs have hatched and repeat as necessary for control. Do not apply more than 3 times during the fruiting period.

SPERMINT (10), PEPPERMINT (10): Twospotted Spider Mite - Use 5 1/2 pints per acre. Do not apply more than twice per season. Do not feed mint hay or spent mint hay to livestock.

STRAWBERRIES (3): Twospotted Spider Mite - Use 4 quarts per acre. During the fruiting period do not apply at intervals of less than 35 days.

TOMATOES (0): Tropical Spider Mite, Twospotted Spider Mite - Use 2 quarts per acre. Do not apply more than 3 times during the fruiting period.

ENVIRONMENTAL AND DISPOSAL CAUTIONS

This product is toxic to fish. Keep out of lakes, ponds and streams. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from the areas treated. Do not apply where runoff is likely to occur. Apply this product only as specified on this label.

Do not reuse empty 1 and 5 gallon containers. Destroy them by perforating or crushing. Return 25 gallon drum to recycler or destroy it by perforating or crushing. Bury or discard in a safe place away from water supplies.

FMC Chemicals

EPA EST. 279-CA-1 Printed in U.S.A.