

CODE
2676

NET WEIGHT
50 LBS.

CHLORDANE 25 GRANULAR

INSECTICIDE

DIRECTIONS FOR USE

CORN SOIL INSECT CONTROL: Corn Rootworm, Seed Corn Beetle - Apply 4 pounds per acre in the row at planting time or apply 8 pounds per acre as a broadcast treatment to soil surface before planting. Disc into top 3 or 4 inches of soil. White Grubs, White-fringed Beetle, Wireworms - Apply 10 to 20 pounds per acre as a broadcast treatment to soil surface before planting. Disc into top 3 or 4 inches of soil.

POTATOES: Japanese Beetle Larvae, White-fringed Beetle Larvae, White Grubs, Wireworms - Apply 16 to 20 pounds per acre. Make broadcast application prior to or at time of planting and work into the top 3 or 4 inches of soil. Increase dosage to 35 to 40 pounds per acre on peat or muck soils.

FPA Reg. No. 279-2543

ACTIVE INGREDIENT:

*Technical Chlordane	25.0%
INERT INGREDIENT:	75.0%
	<hr/> 100.0%

*Equivalent to 15% Octachloro-4,7-methano-tetrahydroindane and 10% related compounds.

CAUTION

KEEP OUT OF REACH OF CHILDREN.

For Emergency Assistance Call
716-735-3765

Avoid breathing dust. Avoid contact with skin. Wash with soap and water after exposure. Avoid contamination of feed and foodstuffs.

USAGE CAUTIONS

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, ponds and streams. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

DISPOSE OF WASTES by burying in non-crop lands away from water supplies. Container should be disposed of by burying with wastes or by burning. Keep out of smoke.

TURF, NURSERY AND GREENHOUSE SOILS (Such as Golf Courses, Athletic Fields, Large Estate Nurseries, Cemeteries, Park Sod, also Nursery and Greenhouse soil containing ornamental plants), **HOME LAWNS:** Chinch Bug, Sod Web worm, Japanese Beetle Grubs, White Grubs, White-fringed Beetle, Rose Chafer, Wireworms, Ants, Armyworms, Boxelder Bugs, Crickets, European Earwig, Grasshoppers, Chiggers, Spiders, Ticks - Apply 20 pounds per acre (1/2 pound per 1,000 square feet). Cover surface uniformly using a lawn fertilizer spreader. Apply in the spring or when insects are observed to be troublesome. Do not treat soil when it is wet. Apply before rain or sprinkling. For HOME LAWN use, immediately after treatment, drench lawn with water or sprinkle heavily to make treatment more effective and to reduce the hazard to children and pets.

CAUTION: Keep children and domestic pets off treated lawns until the insecticide has been washed off the grass into the soil by drenching or sprinkling with water and grass has completely dried.

ACCEPTED