Code: 31145

50 Pounds Net Weig

SULPHUR 10 THIODAN® 3 DUST

DIRECTIONS FOR USE

Apply when insects or disease first appear and repeat as necessary. May be applied with ground or airplane application equipment.

GRAPES to control 3-ape Leafhopper, Grape Leaffolder (Calif. Only) use 35 pounds per acre. For Leaffolder control repeat applications maybe necessary. Do not use on Concord Variety of grapes as severe injury is likely to occur. Do not apply within 7 days of harwest. Apply with ground application equipment only.

BROCCOLI, CABBAGE to control Cabbage Aphids, Cabbage Looper, Diamondback Hoth Larvae, Flea Beetle, Harlequin Bugs, Stink Bugs, Imported Cabbago Worm use 33 pounds per acre. Do not apply within 7 days of

CAULIFLOWER to control Cabbage Looper, Flea Beetle, Harlegiun Bug, Stink Bug, Imported Cabbage worm, Diamond-back Moth Larvae use 25 pounds per acre. Do not apply after edible parts begin to form. Close trim outer leaves prior to marketing.

COTTON to control Cetton Stink Bug, Lygus Bugs, Bollworm, Cabbage Looper, Cotton Leaf perforator use 35 pounds per acre. Do not apply after bolls begin to open. Do not graze meat or dairy animals in treated fields.

HEAD LETTUCE to control Cabbage Looper, Diamond-back Moth Larvae, Green Peach Aphid, Imported Cabbage Worm use 33 pounds per acre. Do not apply more than 3 applications after thinning. Do not apply within 14 days of harvest. Remove outer leaves at harvest. Do not feed crop refuse to livestock.

PEPPERS to control Flea Beetle, Green Peach Aphid, Pepper Maggot and Hornworm use 33 pounds per acre. Do met apply within 1 day of har-

POTATOES to control Aphid, Leaftopper, Flea Beetle, Colorado Potato Beetle, Potato Tuberworm, Green Stinkbug, Southern Armyworm, Potato PayTfid, False Chinch Bug use 30-35 pounds per acre. No time "limitation.

STRAWBERRIES to control Powdery Hildew, Meadow Spittlebug and Strawberry Aphid use 39 pounds per acre. Do not apply within 4 days of picking. For multiple applications during the fruiting period, do not apply at intervals of less than 15 days or more than twice during any 35 day period.

<u>TOMATOES</u> to control Aphid, Flea Beetle, Hornworm, Colorado Potato Bootle, Tometo Psyllid, Whitefly, Stinkbug, Tomato Fruitworm, Cabbage Looper, Yellow Striped Armyworm and Tomato Russet Mite use 33 pounds per acre. Do not apply within 1 day of harvest.

DRIED PEARS to control Black Bean Aphid, Bean Leaf Skeletonizer and Mexican Bean Beetle use 93 pounds per acre. Do not apply within 40 days of harvest. Do not feed treated threshings to livestock. Sulphur may injure certain varieties of beans, wheck sulphur tolerance prior to use.

SUCCULENT BEARS to control Cucumber Beetle use 33 pounds per acre. Do not apply after pode begin to form. Do not feed treated threshings to livestock. Sulphur may injure ceptain varieties of beans, check sulphur tolerance prior to use.

USDA RET. NO. 279-

BY WT. ACTIVE INGREDIENTS: Endouglifan (Hexachlorohexahydromethano-2,4,3benzadioxathiepin oxide) ---- 3.00% Sulphur - - - - - - - - - - - - 10.00% INERT INGREDIENTS ----- 87.00% 100.00%

Thiodan is a registered trademark of Farbwerke Hoechst A.G. - U.S. Patent No. 2.799.685

CAUTION KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing dust. Avoid contact with skin, eyes, or clothing. In case of contact with skin or eyes, flush with plenty of water; for eyes, get medical attention. Wash with soap and water after handling and before eating or smoking; wear clean clothing. Avoid contamination of feed and foodstuffs. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals. Workers entering treated areas on the day of application should wear protective clothing.




USAGE CAUTION

Application of materials containing ruprur stould rut be made prior to or during periods of excess rely ligh tempera or within 10 days to 2 weeks of the scrae.

Do not use on or allow to drift onto me and, shash, cusum apricots or other plants which well not to enate outpour.

This product is poisonous to fish and wild! fer Yeer Lit or ! pends and streams. Do not apply in any mamerinate the field (labe!.

Rinse equipment and dispose of wastes to Linking the cross away from water supplies. Bag should be a single of the bur Keep Out of Smoke .



279-2548

USAGE CAUTION. Do not alion this male in the " onto neighboring crop or non-crop areas or Line 1. 5 %% "." time other than in accordance with direct is in the alan injury, excess we residues or other a delication occur.

Dealers Should Self in Original Packages On v

NOTICE Niagara and Sciller warrant, to the extent of the pin hate. In this the product will conform to the ingredient statement set from no the wide Conformation no other warranty, express or implied covering the crist of the her is also as in the ershall be held responsible in any manner for personal or or other and then loss of any kind resulting from the handling storage in use of the choice whether or not in accordance with directions, regardless of neg penies in tall 1911 Nogari 31 Saller Bover assumes all risk and liability therefrom and errents or to see the control of on these conditions. No person is authorized to vary or waive arising in the estimated

Printed in U.S.A.

Camerauen s Not. toor Copy

OHEM

September 1

50 Pounds Net Weight

SULPHUR 10 THIODAN® 3 DUST

INECTIONS FOR USE sease first appear and repeat as necessary, and or airplane application equipment.

Leafhopper, Grape Leaffolder (Calif. Only) re. For Leaffolder control repeat appliable not use on Concord Variety of grapes as o occur. Do not apply within 7 days of harapplication equipment only.

rol Cabbage Aphids, Cabbage Looper, Diamondetle, Harlequin Bugs, Stink Bugs, Imported a per acre. Do not apply within 7 days of

abbage Looper, Flea Beetle, Harlegiun Bug, age worm, Diamond-back Moth Larvae use 25 apply after edible parts begin to form. prior to marketing.

Stink Bug, Lygus Bugs, Bollworm, Cabbage rator use 35 pounds per acre. Do not apply in. Do not graze meat or dairy animals in

Cabbage Looper, Diamond-back Moth Larvae, led Cabbage Worm use 33 pounds per acre. Do lications after thinning. Do not apply with emove suter leaves at harvest. Do not feed

eetle, Green Peach Aphid, Pepper Maggot and er acre. Do met apply within 1 day of har-

L. Leafhopper, Flea Beetle, Colorado Potato Green Stinkbug, Southern Armyworm, Potato Bug ùse 30-35 pounds per acre. No time

owdery Mildew, Meadow Spittlebug and Straw— To per acre. Do not apply within 4 days of plications during the fruiting period, do f less than 15 days or more than twice dur—

d. Flea Beetle, Hornworm, Colorado Potato Mitefly, Stinkbug, Tomato Fruitworm, Cabeed Armyworm and Tomato Russet Mite use 33 apply within 1 day of harvest.

lack Bean Aphid, Bean Leaf Skeletonizer and \$3 pounds per acre. Do not apply within 40 feed treated threshings to livestock. Sul-arieties of beans, check sulphur tolerance.

of Cucumber Beetle use 33 pounds per acre. begin to form. Do not feed treated threshour may injure ceptain varieties of beans, prior to use.

USDA REG. NO. 270-


ACTIVE INGREDIENTS:	BY	WT.
Endosulfan (Hexachlorohexahydromethano-2,4,3-benzodioxathiepin oxide)	3	00%
Sulphur	10	.005
INERT INGREDIENTS	87	.00%

Thiodan is a registered trademark of Farbwerke Hoechst A.G. - U.S. Patent No. 2,799,685

100.00%

CAUTION KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing dust. Avoid contact with skin, eyes, or clothing. In case of contact with skin or eyes, flush with plenty of water; for eyes, get medical attention. Wash with soap and water after handling and before eating or smoking; wear clean clothing. Avoid contamination of feed and foodstuffs. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals. Morkers entering treated areas on the day of application should wear protective clothing.



• 1.


USAGE CAUTION

Application of materials containing is stur should not be made just prior to or during periods of miss is vely night temperatures or within 10 days to 2 weeks of sprain.

Do not use on or allow to drift onto me on, shaesh, cloumbers, apricots or other plants which will not to erate outpour.

This product is possonous to fish and we do fee Mees let or lakes, ponds and streams. Do not apply in any man en not let fied on the label.

Rinse equipment and dispose of wastes to our, represents away from water supplies. Bag should be a series of the burning. Keep Out of Smoke .



279-25-48

USAGE CALTION. Do not allow this make is it in it onto neighboring crop or non-crop areas or which is man on or at a time other than in accordance with direct is a substantingury, excessive residues or other use its in its may occur.

Dealers Should Sell in Original Packages Only

NOTICE hisagars and Salter warrant, to the extent of the prohase of the product will conform to the ingredient statement set from on the late to the notice and the product will conform to the ingredient statement set from on the late to the next of the her late will be not the sponsible in any manner for personal more product the next value of the loss of any kind resulting from the handling storage of select the product whether or not in accordance with directions, regardless of neg gence of the total his again or Salter Buyer assumes all risk and liability therefrom and accepts and sees the product on these conditions. No person is authorized to vary or waive and to a risk and this notice.

Printed in USA

Cameraden s Not.

r A