
Code 2760

EPN 2 Granular



50 Lbs. Net Weight

Insecticide

EPA Reg. No. 279-2626 EPA Est., 279-IL-1

WARNING KEEP OUT OF REACH OF CHILDREN.

WARNING: May be fatal if swallowed, inhaled or absorbed through skin! Rapidly absorbed through skin. Do not get in eyes, on skin or on clothing. Wear natural rubber gloves, protective clothing and goggles. In case of contact, wash immediately with soap and water. Wear a mask or respirator of a type passed by the U.S. Bureau of Mines for EPN protection. Keep all unprotected persons and beneficial animals out of operating area or vicinity where there may be danger of drift. Vacated areas should not be re-entered until drifting insecticide and volatile residues have dissipated. Do not contaminate feed and foodstuffs. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water befor re-use.

Note to Physician: Atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine. For Emergency Assistance Call 716-735-3765.

Directions for Use NOT FOR USE OR STORAGE IN OR AROUND THE HOME

Apply at the rate of 10 to 25 pounds per acre. Use standard aerial or ground granular insecticide application equipment. Observe days interval between last application and harvest or feeding to livestock as indicated by number in () following crop.

For the control of European Corn Borer

Field Corn (14) Apply 10 days after eggs first hatch or when 75% of the plants show evidence of feeding in leaf whorls. Repeat 7 to 10 days later if new evidence of leaf feeding is present. For control of second brood borers, apply when eggs begin to hatch.

Corn Grown For Seed (14) Apply when 25% of the plants show first brood whorl feeding. Retreat in 7 days if feeding continues. Treat for second brood at pollen shedding time and while the plants are still in the green silk stage.

Sweet Corn (14) Make first application when 25% of the plants show evidence of leaf feeding and continue at 5 day intervals as long as leaf feeding is evident. From one to three treatments may be required. For control of second brood borers, apply when eggs begin to hatch

Store in a cool, dry place.

Environmental and Disposal Cautions

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, ponds and streams. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Dispose of wastes by burying in non-crop lands away from water supplies. Container should be disposed of by burying with wastes or by burning. Keep out of smoke.

Dealers should sell in original packages only.

Terms of sale or use On purchase or use of this product buyer and user agree to the following conditions:

Warranty FMC warrants that this product, as of the time of sale by FMC, (1) conforms to the ingredient statement on the label, and (2) is reasonably fit for the purposes set forth in the Directions for Use. Except as so warranted the product is sold as is. FMC makes no other warranty express or implied.

Damages Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages. Notice of any claim must be reported to FMC within 10 days of discovery by buyer or user, failing which buyer or user waives any claim for such damage.

Excuse from non-controllable conditions Because of certain critical conditions created or incurred by buyer or user or over which FMC has no control, buyer and user, as a condition of purchase or use, assume responsibility for and release FMC from, liability arising out of the handling or use of the product attributable to such causes. No person is authorized to vary or waive any statement contained herein

FMC

FMC Corporation
Agricultural Chemical Division
Middleport New York 14105