

Code: 30895

Net Weight
50 Pounds

[279-2699]

ETHION 4 SEVIN®* 10 DUST

DIRECTIONS FOR USE

Apply when pests first appear and repeat as necessary to maintain control. May be applied by ground or air. Observe days interval between last application and harvest indicated by number in () following the crop.

ALMONDS (30): Codling Moth, Filbert Worm, Fruittree Leafminer, Brown Mite, European Red Mite, Twospotted Spider Mite — Use 40 to 50 pounds per acre. Do not apply more than once during the fruiting period. A rate of 51 to 60 pounds per acre may also be used if no application is made after husk split.

BEANS (2): Mexican Bean Beetle, Bean Leaf Beetle, Cucumber Beetles, Flea Beetles, Leafhoppers, Corn Earworm, Stink Bug, Tarnished Plant Bug, Thrips, Twospotted Spider Mite — Use 20 pounds per acre. Do not feed treated foliage to livestock.

COTTON: ~~Thrips~~, Flea Beetles, Cotton Fleahopper, Cotton Leafworm, Striped Blister Beetle, Soil Weevil, Bollworm, Fall Armyworm, Cotton Leafperforator, Leafhoppers, Leafhoppers, Tarnished Plant Bug, Lygus Bug, Aphids, ~~Professional~~ Pink Bollworm, Grasshoppers, Stink Bug and Saltmarsh Caterpillar — Use 25 pounds per acre. Do not apply after bolls begin to open. Cotton that has been treated should not be grazed or fed to dairy animals or other livestock. In California workers entering treated fields within 24 hours following application should wear protective clothing.

GRAPES: Grape Leafhopper, Twospotted Spider Mite, Willamette Mite, Grape Leafhopper, Grape Leaf Skeletonizer — Use 20 to 30 pounds per acre. Make no more than 2 applications nor apply within 15 days of harvest, or make no more than 3 applications nor apply within 30 days of harvest. Do not apply to Salvador, Muscat or Febré-type varieties of grapes as injury may occur. Ground application preferred.

NECTARINES (30), PEACHES (30): Oriental Fruit Moth, Peach Twig Borer, Periodical Cicada, Codling Moth, Tussock Moth, European Red Mite, Twospotted Spider Mite, Brown Mite — Use 40 to 60 pounds per acre. Do not apply more than twice during the fruiting period.

PLUMS, PRUNES: Peach Twig Borer, Leafhoppers, Codling Moth, Orange Tortrix, Western Tussock Moth, Brown Mite, European Red Mite, Twospotted Spider Mite — Use 40 to 60 pounds per acre. At the 40 to 43 pounds per acre rate, do not make more than 3 applications during the fruiting period nor apply within 21 days of harvest. At the rate of 44 to 60 pounds per acre, do not make more than 2 applications during the fruiting period nor apply within 45 days of harvest.

TOMATOES (2): Colorado Potato Beetle, Flea Beetles, Leafhoppers, Fall Armyworm, Tomato Fruitworms, Tomato Hornworms, Tarnished Plant Bugs, Stink Bugs, Tropical Spider Mites, Twospotted Spider Mites, Leafminers — Use 20 pounds per acre.

INSECTICIDE — MITICIDE

EPA REG. NO. 279-2699 AA

ACTIVE INGREDIENTS	BY WT.
Ethion; 0,0,0'-0'-Tetraethyl S,S'-methylene bisphosphorodithioate	4.00%
*Carbaryl; 1-naphthyl N-methylcarbamate	10.00%
INERT INGREDIENTS	86.00%
	100.00%

S. Patent No. 2,873,228

ENVIRONMENTAL AND DISPOSAL CAUTIONS

This product is toxic to fish and wildlife. Keep out of lakes, ponds, and streams. Do not apply when weather conditions favor drift from the areas treated. Do not apply where run-off is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DISPOSE OF WASTES by burying in non-crop land away from water supplies. Empty bags should be buried with wastes or burned. Keep Out Of Smoke.

CAUTION

For Emergency Assistance Call:
716-735-3765

KEEP OUT OF REACH OF CHILDREN

® Trademark of Union Carbide Corporation.

Harmful if swallowed, inhaled or absorbed through skin. May be absorbed through skin. Avoid breathing dust. Avoid contact with eyes or skin. Wash thoroughly after handling. Wash all contaminated clothing with soap and water before re-use. Avoid storage near or contamination of feed and food-stuffs.

NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

Some phytotoxicity may occur on tender foliage in presence of rain or high humidity of several days duration following application.

279-2699

ADDITIONAL INFORMATION APPEARING ON BAGS TO WHICH THIS LABEL WILL BE ATTACHED.



Dealers Should Sell in Original Packages Only.


DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROPS OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS BECAUSE ANIMAL, PLANT OR CROP INJURY, ILLEGAL RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR.

TERMS OF SALE OR USE: On purchase of this product, buyer and user agree to the following conditions.

WARRANTY: FMC warrants that this product, as of the time of sale by FMC (1) conforms to the ingredient statement on the label, and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED, THE PRODUCT IS SOLD AS IS. FMC MAKES NO OTHER WARRANTY, EXPRESS OR IMPLIED.

DAMAGES: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damage, not exceeding the purchase price paid and shall not include incidental or consequential damages. Notice of any claim must be reported to FMC within 10 days of discovery by buyer or user, failing which buyer or user waives any claim for such damage.

EXCUSE FROM NON-CONTROLLABLE CONDITIONS: Because of certain critical conditions created or incurred by buyer or user, or over which FMC has no control, buyer and user, as a condition of purchase or use, assume responsibility for and release FMC from liability arising out of the handling or use of the product attributable to such causes. No person is authorized to vary or waive any statement contained herein.



Am...
F...
...
...

FMC
F...
...