





Code: 30975

50

# KELTHANE<sup>®</sup> 5 TRITHION<sup>®</sup> 3 DUST

## DIRECTIONS FOR USE

Apply when pests first appear and repeat as necessary to maintain control. May be applied with ground or airplane application equipment. Observe days interval between last application and harvest indicated by number in ( ) following the crop.

**APPLES (30), PEARS (30), CABAPPLES (30), QUINCE (30):** Apple Aphid, European Red Mite, Rosy Apple Aphid, Woolly Apple Aphid, Two-Spotted Mite, Codling Moth, Schoone Mite, McDaniel Mite, Yellow (cargini) Mite, Pacific Mite, Willamette Mite, Apple Rust Mite - Use 40 pounds per acre.

**APRICOTS (30), CHERRIES (30), PEACHES (30), NECTARINES (30):** Aphids, European Red Mite, Clover (Brown Almond), Two-Spotted, Schoone, McDaniel, Pacific, Willamette, Peach Silver, Plum Nursery and Yellow Mites - Use 40-60 pounds per acre. On cherries do not repeat applications within 30 days.

**CANTALOUPE (5), WATERMELON (5):** Melon Aphid, Serpentine Leafminer, Potato Flea Beetle, Two-Spotted and Pacific Mites - Use 25-30 pounds per acre.

**CITRUS (ORANGES, GRAPEFRUIT, LEMON, LIME, TANGELOS, TANGELINES) (7):** Citrus Red, Citrus Flat, Citrus Rust, Six-Spotted, Texas Citrus and Yuma Mites - Use 50-80 pounds per acre. Do not repeat applications within 30 days.

**CUCUMBERS (7), SUMMER SQUASH (7):** Melon Aphid, Potato Leafminer, Two-Spotted and Pacific Mites - Use 25-30 pounds per acre.

**FIGS (7), PLUMS (30), PRUNES (30):** European Red Mite, Clover (Brown Almond), Two-Spotted, McDaniel, Pacific, Willamette, Peach Silver, Rust Mite, Plum Nursery and Yellow Mites - Use 40 pounds per acre. On figs, plums, and prunes do not repeat application within 30 days.

**GRAPES (30):** Grape Leafhopper, Two-Spotted Mite, Pacific Mite, Willamette Mite - Use 30 pounds per acre. Ground application equipment preferred.

## INSECTICIDE-MITICIDE

U.S.D.A. REGISTRATION NO. 279-2701

ACTIVE INGREDIENTS:	BY WT.
Carbophenothion (S-[(p-chlorophenylthio)methyl] 0,0-diethyl phosphorodithioate)	3.00%
1,1-bis (p-chlorophenyl)-2,2,2-trichloroethanol	5.00%
INERT INGREDIENTS:	92.00%
	100.00%

\*Trademark Rohm & Haas Co.

Trithion is a registered trademark of Stauffer Chemical Company for an insecticide-miticide. U.S. Patent No. 2,793,224.

## WARNING

### KEEP OUT OF REACH OF CHILDREN

Hazardous if Swallowed. Avoid breathing dust. Do not get in eyes, on skin or on clothing. Wear dust respirator and goggles. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Do not contaminate feed and foodstuffs. Wash all contaminated clothing with soap and hot water before re-use.

Trithion, upon repeated, prolonged or careless use, may cause cholinesterase inhibition. Atropine is Antidotal. Keep supply of Atropine 1/100 grain tablets on hand.

**PEPPERS (7):** Serpentine Leafminer, hopper, Tropical Mite, Pacific Mite - Use 20-30 pounds per acre.

**SNAP BEANS (7), LIMA BEANS (7):** Serpentine Leaf Miner, Potato Leaf 20-30 pounds per acre. Do not feed

**STRAWBERRY (3):** Strawberry Aphid, leafhopper, and Six-Spotted Mites - Use

**TOMATOES (7):** Serpentine Leafminer, Beetle, Potato Leafhopper, Tomato Spotted Mite - Use 20-30 pounds

## USAGE C

This product is toxic to fish and streams. Birds feeding on this product only as specified on

This product is toxic to bees and are actively visiting the area.

DISPOSE OF WASTES by burying in supplies. Empty bags should be out of smoke.

279-2701

ACCEPTED  
Dept. 1/15  
WHEN THE FEDERAL GOVERNMENT AND ROYALTY FOR ECONOMIC POISON IS WITHIN NO. TO AFFECTED COMMERCE

50 Pounds Net Weight

# KELTHANE® 5 TRITHION® 3 DUST

## USES FOR USE

and repeat as necessary to maintain ground or airplane application equipment between last application and harvest during the crop.

**APPLES (30), QUINCE (30):** Apple, Apple Aphid, Woolly Apple Aphid, B. Schoone Mite, McDaniel Mite, Willamette Mite, Apple Rust Mite

**PEACHES (30), NECTARINES (30):** Clover (Brown Almond), Two-Spotted, Yamette, Peach Silver, Plum Nursery pounds per acre. On cherries do not repeat.

**MELON (5):** Melon Aphid, Serpentine Leafhopper, and Pacific Mites - Use 25-30

**LEMON, LIME, TANGELOS, TANGERS:** Flat, Citrus Rust, Six-Spotted, Texas 10-80 pounds per acre. Do not repeat

**CASH (7):** Melon Aphid, Potato Leafhopper - Use 25-30 pounds per acre.

**(10):** European Red Mite, Clover (Brown Almond), Pacific, Willamette, Peach Silver, Yellow Mites - Use 40 pounds per acre. Do not repeat application within

per, Two-Spotted Mite, Pacific Mite, pounds per acre. Ground application

## INSECTICIDE-MITICIDE

U.S.D.A. REGISTRATION NO. 279-2701

ACTIVE INGREDIENTS:	BY WT.
Carbophenothion (S- [(p-chlorophenylthio)methyl] 0,0-diethyl phosphorodithioate)	3.00%
1,1-bis (p-chlorophenyl)-2,2,2-trichloroethanol	5.00%
INERT INGREDIENTS:	92.00%
	100.00%

\*Trademark Rohm & Haas Co.

Trithion is a registered trademark of Stauffer Chemical Company for an insecticide-miticide. U.S. Patent No. 2,793,224.

## WARNING

### KEEP OUT OF REACH OF CHILDREN

Hazardous if Swallowed. Avoid breathing dust. Do not get in eyes, on skin or on clothing. Wear dust respirator and goggles. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Do not contaminate feed and foodstuffs. Wash all contaminated clothing with soap and hot water before re-use.

Trithion, upon repeated, prolonged or careless use, may cause cholinesterase inhibition. Atropine is Antidotal. Keep supply of Atropine 1/100 grain tablets on hand.

**PEPPERS (7):** Serpentine Leafminer, Potato Flea Beetle, Potato Leafhopper, Tropical Mite, Pacific Mite, Two-Spotted Mite - Use 25-30 pounds per acre.

**SNAP BEANS (7), LIMA BEANS (7):** Bean Aphid, Mexican Bean Beetle, Serpentine Leaf Miner, Potato Leafhopper, Two-Spotted Mite - Use 20-30 pounds per acre. Do not feed treated forage to livestock.

**STRAWBERRY (3):** Strawberry Aphid, Two-Spotted, Strawberry (Atlantic), and Six-Spotted Mites - Use 25-40 pounds per acre.


**TOMATOES (7):** Serpentine Leafminer, Potato Psyllid, Potato Flea Beetle, Potato Leafhopper, Tomato Russet Mite, Tropical Mite, Two-Spotted Mite - Use 20-30 pounds per acre.

## USAGE CAUTIONS

This product is toxic to fish and wildlife. Keep out of lakes, ponds and streams. Birds feeding on treated areas may be killed. Apply this product only as specified on the label.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

DISPOSE OF WASTES by burying in non-crop lands away from water supplies. Empty bags should be buried with wastes or burned. Keep out of smoke.



30975

# KELTHANE\* 5 TRITHION® 3

## DIRECTIONS FOR USE

Apply when pests first appear and repeat as necessary to maintain control. May be applied with ground or airplane application equipment. Observe days interval between last application and harvest indicated by number in ( ) following the crop.

**APPLES (30), PEARS (30), CABAPPLES (30), QUINCE (30):** Apple Aphid, European Red Mite, Rosy Apple Aphid, Woolly Apple A Two-Spotted Mite, Codling Moth, Schoene Mite, McDaniel, Yellow (carpini) Mite, Pacific Mite, Willamette Mite, Apple Rus. Mite — Use 40 pounds per acre.

**APRICOTS (30), CHERRIES (30), PEACHES (30), NECTARINES (30):** Aphids, European Red Mite, Clover (Brown Almond), Two-Spotted, Schoene, McDaniel, Pacific, Willamette, Peach Silver, Plum Nursery and Yellow Mites — Use 40-60 pounds per acre. On cherries do not repeat applications within 30 days.

**CANTALOUPE (5), WATERMELON (5):** Melon Aphid, Serpentine Leaf-miner, Potato Flea Beetle, Two-Spotted, and Pacific Mites — Use 25-30 pounds per acre.

**CITRUS (ORANGES, GRAPEFRUIT, LEMON, LIME, TANGELOES, TANGERINES) (7):** Citrus Red, Citrus Flat, Citrus Rust, Six-Spotted, Texas Citrus and Yuma Mites — Use 50-80 pounds per acre. Do not repeat applications within 30 days.

**CUCUMBERS (7), SUMMER SQUASH (7):** Melon Aphid, Potato Leaf-miner, Two-Spotted and Pacific Mites — Use 25-30 pounds per acre.

**FIGS (7), PLUMS (30), PRUNES (30):** European Red Mite, Clover (Brown Almond), Two-Spotted, McDaniel, Pacific, Willamette, Peach Silver, Rust Mite, Plum Nursery and Yellow Mites — Use 40 pounds per acre. On figs, plums, and prunes do not repeat application within 30 days.

**GRAPES (30):** Grape Leafhopper, Two-Spotted Mite, Pacific Mite, Willamette Mite — Use 30 pounds per acre. Ground application equipment preferred.

## INSECTICIDE-MITICIDE

U.S.D.A. REGISTRATION NO. 279-2701

ACTIVE INGREDIENTS:	BY WT.
Carbophenothion (S- [(p-chlorophenylthio) methyl] 0,0-diethyl phosphorodithioate) .....	3.00%
1,1-bis (p-chlorophenyl)-2,2,2-trichloroethanol .....	5.00%
INERT INGREDIENTS: .....	92.00%
	100.00%

\*Trademark Rohm & Haas Co.

Trithion is a registered trademark of Stauffer Chemical Company for an insecticide-miticide. U.S. Patent No. 2,793,224.

## WARNING

### KEEP OUT OF REACH OF CHILDREN

Hazardous if Swallowed. Avoid breathing dust. Do not get in eyes, on skin or on clothing. Wear dust respirator and goggles. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Do not contaminate feed and foodstuffs. Wash all contaminated clothing with soap and hot water before re-use.

Trithion, upon repeated, prolonged or careless use, may cause cholinesterase inhibition. Atropine is Antidotal. Keep supply of Atropine 1/100 grain tablets on hand.

50 Pounds Net Weight

# KELTHANE\* 5 TRITHION® 3 DUST

### USE FOR USE

Repeat as necessary to maintain and or airplane application equipment between last application and harvest of the crop.

**APPLES (30), QUINCE (30):** Apple Aphid, Woolly Apple Aphid, Shore Mite, McDaniel Mite, Willamette Mite, Apple Rust Mite

**PEACHES (30), NECTARINES (30):** Brown Almond, Two Spotted Mite, Peach Silver, Plum Nursery Mites per acre. On cherries do not repeat.

**MELON (7):** Melon Aphid, Serpentine Leafminer, and Pacific Mites - Use 25-30

**LEMON, LIME, TANGELOES, TANGERS:** Citrus Rust, Six-Spotted Mite, Texas Mite - Use 25-30 pounds per acre. Do not repeat

**POTATO (7):** Melon Aphid, Potato Leafminer - Use 25-30 pounds per acre.

European Red Mite, Clover (Brown) Mite, Pacific Willamette, Peach Silver, Two Spotted Mites - Use 40 pounds per acre. Do not repeat application within

Two Spotted Mite, Pacific Mite, Shore Mite per acre. Ground application

### INSECTICIDE-MITICIDE

U.S.D.A. REGISTRATION NO. 279-2701

ACTIVE INGREDIENTS:	BY WT.
Carbophenothion (S-[(p-chlorophenylthio)methyl] 0,0-diethyl phosphorodithioate) .....	3.00%
1,1-bis (p-chlorophenyl)-2,2,2-trichloroethanol .....	5.00%
INERT INGREDIENTS: .....	92.00%
	100.00%

\*Trademark Rohm & Haas Co.

Trithion is a registered trademark of Stauffer Chemical Company for an insecticide-miticide. U.S. Patent No. 2,793,224.

## WARNING

### KEEP OUT OF REACH OF CHILDREN

Hazardous if Swallowed. Avoid breathing dust. Do not get in eyes, on skin or on clothing. Wear dust respirator and goggles. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Do not contaminate feed and foodstuffs. Wash all contaminated clothing with soap and hot water before re-use.

Trithion, upon repeated, prolonged or careless use, may cause cholinesterase inhibition. Atropine is Antidotal. Keep supply of Atropine 1/100 grain tablets on hand.

**PEPPERS (7):** Serpentine Leafminer, Potato Flea Beetle, Potato Leafhopper, Tropical Mite, Pacific Mite, Two Spotted Mite - Use 25-30 pounds per acre.

**SNAP BEANS (7), LIMA BEANS (7):** Bean Aphid, Mexican Bean Beetle, Serpentine Leaf Miner, Potato Leafhopper, Two Spotted Mites - Use 20-30 pounds per acre. Do not feed treated forage to livestock.

**STRAWBERRY (3):** Strawberry Aphid, Two Spotted, Strawberry (Atlantic), and Six Spotted Mites - Use 25-40 pounds per acre.

**TOMATOES (7):** Serpentine Leafminer, Potato Psyllid, Potato Flea Beetle, Potato Leafhopper, Tomato Russet Mite, Tropical Mite, Two Spotted Mite - Use 20-30 pounds per acre.

### USAGE CAUTIONS

This product is toxic to fish and wildlife. Keep out of lakes, ponds and streams. Birds feeding on treated areas may be killed. Apply this product only as specified on the label.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

DISPOSE OF WASTES by burying in non-crop lands away from water supplies. Empty bags should be buried with wastes or burned. Keep out of smoke.

