

Code: 30965

[279-2706]

50 Po

MANUFACTURED:

DIBROM[®] 4 ETHION 4 DUST

DIRECTIONS FOR USE

Apply this product when insects or mites first appear and repeat as necessary. May be applied with ground or airplane application equipment. Observe days interval between last application and harvest indicated by number in () following the crop.

BEANS (4): Bean Leaf Beetle, Mexican Bean Beetle, Two-Spotted Spider Mite — Use 20-40 pounds per acre. Lygus Bugs, Aphids, Thrips, Leafhoppers, Leafminers, Cabbage Loopers — Use 40 pounds per acre. Do not feed treated foliage to livestock.

GRAPEFRUIT (1), LEMONS (21), ORANGES (1) (California and Arizona): Thrips, Aphids, Red Spider Mite, Citrus Cutworms — Use 40-100 pounds per acre. Do not apply more than once per season on lemons; on other listed fruits do not apply within 90 days of previous applications.

GRAPES: Grape Leafhopper, Two-Spotted Spider Mite, Willamette Mite, Thrips (Calif. Only), Drosophila (Calif. Only) — Use 30 pounds per acre. Make no more than 2 applications per season nor apply within 15 days of harvest; or make no more than 3 applications per season nor apply within 30 days of harvest. Do not apply to Salvador, Muscat or Fehr-Szago varieties as injury may occur. Ground application equipment preferred.

MELONS (Cantaloupes, Honeydews, Muskmelons, Watermelons) (1): Melon Leafhopper, Melon Leafminer, Serpentine Leafminer, Two-Spotted Spider Mite — Use 30-40 pounds per acre. Aphids, Armyworms, Cabbage Loopers, Lygus Bugs — Use 40 pounds per acre.

STRAWBERRIES (4): Two-Spotted Spider Mite — Use 20 pounds per acre.

TOMATOES (2): Leafminer, Tropical Spider Mite, Two-Spotted Spider Mite — Use 20 pounds per acre.

INSECTICIDE-MITICIDE

EPA REGISTRATION NO. 279-2706 AA	
ACTIVE INGREDIENTS:	BY WT.
*Naled (1,2-dibromo-2,2-dichloroethyl dimethyl phosphate)	4.00%
**Ethion	4.00%
INERT INGREDIENTS:	92.00%
	100.00%
**0,0,0',0'-Tetraethyl S,S'-methylene bisphosphorodithioate. U.S. Pat. No. 2,873,228.	

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. May be absorbed through skin. May cause irritation of eyes, nose and throat. Avoid breathing dust. Avoid contact with skin, eyes and clothing. Wash thoroughly after using or handling. Avoid contamination of or storage near feed and foodstuffs.

NOTE TO PHYSICIAN: Atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with Atropine.

USAGE CAUTIONS

This product is toxic to fish and wild birds and streams. Birds feeding on treated crops may contaminate water by cleaning of equipment. Apply this product only as specified on label.

This product is highly toxic to bees and other beneficial insects. Protective information should be given to your Cooperative Agricultural Extension Agent.

Do not apply when weather conditions are unfavorable. Do not apply when run-off is likely.


STORE IN COOL, DRY PLACE.

May corrode metal dust equipment; clean after use.

DISPOSE OF WASTES by burying in soil or water supplies. Empty bags should be destroyed. Do not re-use.

For Emergency Assistance
716-735-3700

*DIBROM is a registered trademark of D. I. D. U.S. Patent No. 2,971,882.



7067



50 Pounds Net Weight

DIBROM® 4 ETHION 4 DUST

GUARANTEE EXPIRES:

Repeat and repeat as airplane application. First application and re-crop.

Colorado Beetle, Two-Spotted Spider, Greenbugs, Aphids, etc. — Use 40 pounds per acre.

California and Arizona Armyworms — Use 40 pounds per acre once per season within 90 days of emergence.

Colorado Mite, Willamette Mite — Use 30 pounds per acre per season nor apply more than 3 applications per season. Do not apply to Salvage crops. Ground application.

Watermelons (1): Leafminer, Two-spotted Spider, Aphids, Armyworms — Use 20 pounds per acre.

Two-Spotted Spider

INSECTICIDE-MITICIDE

EPA REGISTRATION NO. 279-2706 AA

ACTIVE INGREDIENTS:	BY WT.
*Naled (1,2-dibromo-2,2-dichloroethyl dimethyl phosphate)	4.00%
**Ethion	4.00%
INERT INGREDIENTS:	92.00%
	<u>100.00%</u>


**0,0,0',0'-Tetraethyl S,S'-methylene bisphosphorodithioate. U.S. Pat. No. 2,873,228.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. May be absorbed through skin. May cause irritation of eyes, nose and throat. Avoid breathing dust. Avoid contact with skin, eyes and clothing. Wash thoroughly after using or handling. Avoid contamination of or storage near feed and foodstuffs.

NOTE TO PHYSICIAN: Atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with Atropine.



USAGE CAUTIONS

This product is toxic to fish and wildlife. Keep out of lakes, ponds and streams. Birds feeding on treated areas may be killed. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on the label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not apply when weather conditions favor drift from the areas treated. Do not apply when run-off is likely to occur.

STORE IN COOL, DRY PLACE.

May corrode metal dust equipment; clean equipment after use.

DISPOSE OF WASTES by burying in non-crop lands away from water supplies. Empty bags should be buried with wastes or burned. Do not re-use.

For Emergency Assistance Call 716-735-3765

*DIBROM is a registered trademark of Chevron Chemical Company. U.S. Patent No. 2,971,882.



Code: 30965

[279-2706]

MANUFACTURED:

DIBROM[®] 4 ETH

DIRECTIONS FOR USE

Apply this product when insects or mites first appear and repeat as necessary. May be applied with ground or airplane application equipment. Observe days interval between last application and harvest indicated by number in () following the crop.

BEANS (4): Bean Leaf Beetle, Mexican Bean Beetle, Two-Spotted Spider Mite — Use 20-40 pounds per acre. Lygus Bugs, Aphids, Thrips, Leafhoppers, Leafminers, Cabbage Loopers — Use 40 pounds per acre. Do not feed treated foliage to livestock.

GRAPEFRUIT (1), LEMONS (21), ORANGES (1) (California and Arizona): Thrips, Aphids, Red Spider Mite, Citrus Cutworms — Use 40-100 pounds per acre. Do not apply more than once per season on lemons; on other listed fruits do not apply within 90 days of previous applications.

GRAPES: Grape Leafhopper, Two-Spotted Spider Mite, Willamette Mite, Thrips (Calif. Only), Drosophila (Calif. Only) — Use 30 pounds per acre. Make no more than 2 applications per season nor apply within 15 days of harvest; or make no more than 3 applications per season nor apply within 30 days of harvest. Do not apply to Salvador, Muscat or Fehr-Szago varieties as injury may occur. Ground application equipment preferred.

MELONS (Cantaloupes, Honeydews, Muskmelons, Watermelons) (1): Melon Leafhopper, Melon Leafminer, Serpentine Leafminer, Two-Spotted Spider Mite — Use 30-40 pounds per acre. Aphids, Armyworms, Cabbage Loopers, Lygus Bugs — Use 40 pounds per acre.

STRAWBERRIES (4): Two-Spotted Spider Mite — Use 20 pounds per acre.

TOMATOES (2): Leafminer, Tropical Spider Mite, Two-Spotted Spider Mite — Use 20 pounds per acre.

INSECTICIDE

EPA REGISTRATION
ACTIVE INGREDIENTS:

*Naled (1,2-dibromo-2,2-dimethyl phosphate) ...

**Ethion

INERT INGREDIENTS: ...

**0,0,0',0'-Tetraethyl S,S'-methyldithioate. U.S. Pat. No. 2,87

CAUTION

KEEP OUT OF REACH

Harmful if swallowed. May cause irritation of eyes, nose, and throat. Avoid breathing dust. Avoid contact with skin. Wash thoroughly after use. Do not contaminate food or storage near food.

NOTE TO PHYSICIAN: Atropine is also antidotal and may be used with Atropine.

AC