

[279-2847]

CODE  
JDL-1



NET WEIGHT  
**50** LBS.

# PYRENONE<sup>®</sup> FLEXI-DUST

## INSECTICIDE

### DIRECTIONS FOR USE

Apply at the rate of 30 pounds of dust per acre. Treat only when insects are present and repeat as required to maintain control. Apply dust uniformly to upper and lower leaf surfaces.

<u>CROP</u>	<u>TO CONTROL</u>
ASPARAGUS	Asparagus Beetle
BEANS	Aphids, Leafhoppers, White Fly, Mexican Bean Beetles
BROCCOLI CABBAGE BRUSSELS SPROUTS CAULIFLOWER	Aphids, Cabbage Looper, Cross-striped Cabbageworm, Diamondback Moth Larvae, Flea Beetles, Harlequin Bug, Imported Cabbageworm
CELERY	Green Peach Aphid, Cabbage Looper, Leaf Tier
CRANBERRIES	Fireworms, Leafhoppers
EGGPLANTS	Colorado Potato Beetle, Flea Beetles, Green Peach Aphid
LETTUCE	Cabbage Looper, Green Peach Aphid, Diamondback Moth Larvae, Imported Cabbageworm
PEPPERS	Green Peach Aphid
POTATOES	Aphids, Colorado Potato Beetle, Flea Beetles, Leafhoppers
RADISH	Aphids, Flea Beetles
SPINACH	Aphids, Cabbage Looper, Webworm
TOMATOES	Green Peach Aphid, Colorado Potato Beetle, Flea Beetles
ORNAMENTALS	Aphids, Flea Beetles, Leafhoppers on: (African Violets, Asters, Azaleas, Begonias, Camellias, Carnations, Chrysanthemums, Dahlias, Dogwood, Geraniums, Gladioli, Marigolds, Roses, etc.)

**FRUIT FLY CONTROL OF FRUITS AND VEGETABLES:** To control fruit flies (*Drosophila*) on apples, blackberries, boysenberries, currants, dewberries, grapes, loganberries, peaches, pears, plums, prunes,

USDA Reg. No. 279-2847

### ACTIVE INGREDIENTS:

\*Technical Piperonyl Butoxide . . . . . 1.00%  
Pyrethrins . . . . . 0.10%

INERT INGREDIENTS: . . . . . 98.90%

\*Equivalent to 0.80% (butylcarbityl) (6-propylpiperonyl) ether and 0.20% of other related products.

## CAUTION

### KEEP OUT OF REACH OF CHILDREN.

AVOID INHALATION OF DUST.

PYRENONE FLEXI-DUST is intended for use as a contact insecticide during the growing season or as a pre-harvest treatment where other insecticides cannot be used because of residue restrictions and use limitations.

### USAGE CAUTIONS

Destroy empty container by burning.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

### Dealers Should Sell in Original Packages Only.

**TERMS OF SALE OR USE:** On purchase or use of this product buyer and user agree to the following conditions:  
**WARRANTY:** Niagara warrants that this product, as of the time of sale by Niagara, (1) conforms to the ingredient statement on the label, and (2) is reasonably fit for the purposes set forth in the Directions for Use. **EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS. NIAGARA MAKES NO OTHER WARRANTY, EXPRESS OR IMPLIED.**



**NIAGARA CHEMICAL DIVISION**  
**FMC CORPORATION**  
**MIDDLEPORT, N. Y. 14105**

melons and tomatoes apply as a POST-HARVEST application. Dust fruits or vegetables immediately after picking while in containers in the field. Remove crops from field as soon as they are picked and have been dusted. Do not hold picked crops in field overnight. If the crops are held at the processing plants more than 20 hours, a repeat application should be made. Dust lightly but thoroughly.

**STORED GRAIN PROTECTANT:** For use on corn, barley, rice, rye, buckwheat, birdseed, and other seeds in storage.

Pyrenone Flexi-Dust kills accessible stages of many grain insects particularly rice weevils and granary weevils, and protects against many insects in the bran-bug group including flour beetles and grain beetles. One treatment is usually adequate for a full storage season.

Treated grains require only normal cleaning before processing for human food and may be fed to livestock without hazard. Shelled corn may require cleaning if subject to inspection for grading purposes.

Whole grains and seeds to be treated should be in condition for normal storage. Treatment should be applied when the grain goes into storage and before serious infestation has developed.

### HOW MUCH PYRENONE FLEXI-DUST TO APPLY:

Application on grains and seeds should be made at the rate of 80 to 100 pounds per 1,000 bushels (8/10 to 1 pound per 10 bushels). The dust should be sufficiently well mixed with the whole grain or seeds to assure even distribution. It can be applied to a moving stream of grain in the usual or typical conveyor. Small lots of grain or seeds can be treated by hand with a shovel or other suitable equipment.

Pyrenone Flexi-Dust on ear corn including snapped corn should be applied at the same rate specified above, and should be sufficiently well mixed with the corn to assure even distribution. One method is by dusting the protectant over alternate shallow layers of the corn.

Snapped (unshucked) corn should be shucked before treatment for best results. Otherwise the benefit of the treatment is primarily for the prevention of insect migration from infested ears. The amount of ear corn that will produce a bushel of shelled corn is figured as an "ear-corn bushel".