



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

July 13, 2020

Bonnie Bieber
FMC Corporation
c/o FMC Stine Research Center
1090 Elkton Road
Newark, Delaware 19711

Subject: Registration Review Label Mitigation for Metsulfuron Methyl and Tribenuron Methyl
Product Name: Amathon Herbicide (with TotalSol Soluble Granules)
Application Date: 12/12/2017
EPA Registration Number: 279-9618
Decision Number: 540736; 564567

Dear Ms. Bieber:

The Agency, in accordance with the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, has completed reviewing all of the information submitted with your application to support the Registration Review of the above referenced product in connection with the 22 Sulfonylurea (SU) Herbicides Interim Decision, and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

A copy of your label stamped "Accepted" is enclosed. Products shipped after 12 months from the date of this amendment must bear the new revised label. Your release for shipment of the product bearing the amended label constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6.

Page 2 of 2
EPA Reg. No. 279-9618
Decision No. 540736; 564567

If you have any questions about this letter, please contact Erik Kraft by phone at 703-308-9358, or via email at kraft.erik@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read "Erik Kraft". The signature is written in a cursive style with a large initial "E" and a long, sweeping tail.

Erik Kraft, Product Manager 24
Fungicide and Herbicide Branch
Registration Division (7505P)
Office of Pesticide Programs

Enclosure

AMATHON™

HERBICIDE
(WITH TOTALSOL® SOLUBLE GRANULES)

ACCEPTED

07/13/2020

Under the Federal Insecticide, Fungicide
and Rodenticide Act as amended, for the
pesticide registered under
EPA Reg. No. 279-9618

Tribenuron Methyl	GROUP	2	Herbicide
Metsulfuron Methyl	GROUP	2	Herbicide

Soluble Granule

For Use on Cereals, Fallow and as a Pre-plant or Post-harvest Burndown Herbicide

Active Ingredients	By Weight
Tribenuron methyl	45%
Metsulfuron Methyl	6%
Other Ingredients	49%
TOTAL	100%

Contains 0.45 lb Tribenuron Methyl and 0.06 lb Metsulfuron Methyl per pound

EPA Reg. No. 279-9618

Nonrefillable Container

Refillable Container

EPA Est. No. _____

Net: _____

OR

Net: _____

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

IF ON SKIN: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

You may also contact 1-800-331-3148 for emergency medical treatment information.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing dust or spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. For medical emergencies involving this product, call toll free 1-800-331-3148.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

Long-sleeved shirt and long pants.

Chemical resistant gloves made of any waterproof material including polyethylene or polyvinyl chloride. Shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exists, use detergent and hot water. Keep and wash PPE separately from other laundry.

Sold By



FMC Corporation
2929 Walnut Street
Philadelphia, PA 19104

USER SAFETY RECOMMENDATIONS

USERS SHOULD:

Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

DO NOT apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when cleaning equipment or disposing of equipment.

Groundwater Advisory

This product is known to leach through soil into groundwater under certain conditions as a result of label use. This product may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several weeks or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features including ponds, streams, and springs will reduce the potential loading of this product from runoff water and sediment. Runoff of this product will be greatly reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

Windblown Soil Particles Advisory

This product has the potential to move off-site due to wind erosion. Soils that are subject to wind erosion usually have a high silt and/or fine to very fine sand fractions and low organic matter content. Other factors which can affect the movement of windblown soil include the intensity and direction of prevailing winds, vegetative cover, site slope, rainfall, and drainage patterns. Avoid applying this product if prevailing local conditions may be expected to result in off-site movement.

Non-target Organism Advisory

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated area. Protect the forage and habitat of non-target organisms by minimizing spray drift. For further guidance and instructions on how to minimize spray drift, refer to the Spray Drift Management section of this label

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, including plants, soil, or water is:

- Coveralls.
- Chemical resistant gloves made of any waterproof material.
- Shoes plus socks.

AMATHON™ herbicide (with TotalSol® soluble granules), referred to below as AMATHON herbicide, must be used only in accordance with instructions on this label or as otherwise permitted by FIFRA.

To the extent consistent with applicable law, FMC will not be responsible for losses or damages resulting from the use of this product in any manner not specifically directed by FMC. Check with your state extension service or Department of Agriculture before use, to be certain AMATHON herbicide is registered in your state.

PRODUCT INFORMATION

AMATHON herbicide is a water soluble granule that is used for selective postemergence weed control in wheat (including durum), barley, triticale and for post-harvest burndown, fallow, and pre-plant burndown weed control. The best control is obtained when AMATHON herbicide is applied to young, actively growing weeds. The use rate will depend on weed spectrum and size of weed at time of application. The degree and duration of control may depend on the weed spectrum and infestation intensity, weed size at application, and environmental conditions at and following treatment.

AMATHON herbicide is noncorrosive, nonflammable, nonvolatile, and does not freeze. AMATHON herbicide must be mixed in water and applied as a uniform broadcast spray.

ENVIRONMENTAL CONDITIONS AND BIOLOGICAL ACTIVITY

AMATHON herbicide is absorbed through the foliage of broadleaf weeds, rapidly inhibiting their growth. Leaves of susceptible plants appear chlorotic from 1 to 3 weeks after application and the growing point subsequently dies.

AMATHON herbicide provides the best control in vigorously growing crops that shade competitive weeds. Weed control in areas of thin crop stand or seeding skips may not be as satisfactory. However, a crop canopy that is too dense at application can intercept spray and reduce weed control.

AMATHON herbicide may injure crops that are stressed from adverse environmental conditions (including extreme temperatures or moisture), abnormal soil conditions, or cultural practices. In addition, different varieties of the crop may have differing levels of sensitivity to treatment with AMATHON herbicide under otherwise normal conditions.

Treatment of sensitive crop varieties may injure crops. To reduce the potential of crop injury to cereals, tank mix AMATHON herbicide with 2,4-D (ester formulations perform best—see the Tank Mixtures section of this label) and apply after the crop is in the tillering stage of growth.

In warm, moist conditions, the expression of herbicide symptoms is accelerated in weeds; in cold, dry conditions, expression of herbicide symptoms is delayed. In addition, weeds hardened-off by drought stress are less susceptible to AMATHON herbicide.

AMATHON herbicide is rainfast in 4 hours.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. If the instructions on the tank mix partner label conflicts with this AMATHON herbicide label, **DO NOT** use in a tank mixture with AMATHON herbicide.

RESTRICTIONS

- **DO NOT** apply this product through any type of irrigation system.
- **DO NOT** discharge excess material on the soil at a single spot in the field, grove, or mixing/loading station.
- **DO NOT** apply to wheat, barley or triticale underseeded with another crop.
- **DO NOT** apply, drain or flush equipment on or near desirable trees or other plants or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.
- **DO NOT** use on lawns, walks, driveways, tennis courts. Prevent drift of spray to desirable plants.
- **DO NOT** use on grasses grown for seed.
- **DO NOT** apply to irrigated land where tailwater will be used to irrigate crops other than wheat and barley.
- **DO NOT** apply to frozen ground as surface runoff may occur.
- **DO NOT** apply to snow-covered ground.
- When using AMATHON herbicide in tank mixes or sequential applications with other products containing tribenuron methyl and metsulfuron methyl, **DO NOT** exceed the following limits.

Use	Active Ingredient	Maximum lb ai per Single Application	Maximum lb ai per Year
Cereals Burndown	Tribenuron methyl	0.0094	0.016
	Metsulfuron methyl	0.0013	0.0037

PRECAUTIONS

- Calibrate sprayers only with clean water away from well sites.
- Make scheduled checks of spray equipment.
- Ensure that all operation employees accurately measure pesticides.
- Mix only enough product for the job at hand.
- Avoid overfilling of spray tank.
- Avoid storage of pesticides near well sites.
- When triple-rinsing the pesticide container, be sure to add the rinsate to the spray mix.
- Dilute and agitate excess solution and apply at labeled rates or uses.
- To reduce the potential for movement of treated soil due to wind erosion, do not apply to powdery dry or light sandy soils until they have been stabilized by rainfall, trashy mulch, reduced tillage, or other cultural practices. Injury to immediately adjacent crops may occur when treated soil is blown onto land used to produce crops other than cereal grains or pasture/rangeland.
- Injury to or loss of adjacent sensitive crops and vegetation may result from failure to observe the following:
 - Take all necessary precautions to avoid all direct or indirect contact (including spray drift) with non-target plants or areas.
 - Carefully observe all sprayer cleanup instructions both prior to and after using this product, as spray tank residue may damage crops other than wheat or barley.
- Varieties of wheat (including durum), barley and triticale may differ in their response to various herbicides. FMC advises that you first consult your state experiment station, university, or extension agent as to crop sensitivity to any herbicide. If no information is available, limit the initial use to a small area.
- The combined treatment effects of AMATHON herbicide postemergence preceded by preemergence wild oat herbicides may cause crop injury to spring wheat when crop stress (soil crusting, planting too deep, prolonged cold weather, or drought) causes poor seedling vigor.
- Under certain conditions including heavy rainfall, prolonged cold weather, or wide fluctuations in day/night temperatures prior to or soon after AMATHON herbicide application, temporary discoloration and/or crop injury may occur. To reduce the potential of crop injury, tank mix AMATHON herbicide with 2,4-D (ester formulations perform best - see the "TANK MIXTURES" section of this label) and apply after the crop is in the tillering stage of growth.
- In the Pacific Northwest, to prevent cold weather-related crop injury, avoid making applications during winter months when weather conditions are unpredictable and can be severe.
- AMATHON herbicide must not be applied to wheat, barley or triticale that is stressed by severe weather conditions, drought, low fertility, water-saturated soil, disease, or insect damage, as crop injury may result. Risk of injury is greatest when crop is in the 2 to 5- leaf stage. Severe winter stress, drought, disease, or insect damage following application also may result in crop injury.
- Dry, dusty field conditions may result in reduced control in wheel track areas.

WEED RESISTANCE MANAGEMENT

AMATHON herbicide which contains the active ingredients tribenuron methyl and metsulfuron methyl is a Group 2 herbicide based on the mode of action classification system of the Weed Science Society of America. Proactively implementing diversified weed control strategies to minimize selection for weed populations resistant to one or more herbicides is a best practice. A diversified weed management program may include the use of multiple herbicides with different sites of action and overlapping weed spectrum with or without tillage operations and/or other cultural practices. Research has demonstrated that using the labeled rate and directions for use is important to delay the selection for resistance.

The continued effectiveness of this product depends on the successful implementation of a weed resistance management program.

To aid in the prevention of developing weeds resistant to this product, users must follow as many of the following herbicide resistance management practices as is practical:

- Scout fields before application to ensure herbicides and rates will be appropriate for the weed species and weed sizes present.
- Start with a clean field, using either a burndown herbicide application or tillage.
- Control weeds early when they are relatively small (less than 4 inches).
- Apply full rates of AMATHON herbicide for the most difficult to control weed in the field at the specified time (correct weed size) to minimize weed escapes.
- Scout fields after application to detect weed escapes or shifts in control of weed species.
- Control weed escapes before they reproduce by seed or proliferate vegetatively.

- Report any incidence of non-performance of this product against a particular weed to your FMC representative, local retailer, or county extension agent.
- Contact your FMC representative, crop advisor, or extension agent to find out if suspected resistant weeds to this MOA have been found in your region. If resistant biotypes of target weeds have been reported, use the application rates of this product specified for your local conditions. Tank mix products so that there are multiple effective sites of actions for each target weed.
- If resistance is suspected, treat weed escapes with an herbicide having a site of action other than Group 2 and/or use nonchemical methods to remove escapes, as practical, with the goal of preventing further seed production.
- Suspected herbicide-resistant weeds may be identified by these indicators:
 - Failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;
 - A spreading patch of non-controlled plants of a particular weed species; and
 - Surviving plants mixed with controlled individuals of the same species.
- Use a broad spectrum soil-applied herbicide with other sites of action as a foundation in a weed control program.
- Utilize sequential applications of herbicides with alternative sites of action.
- Rotate the use of this product with non-Group 2 herbicides.
- Avoid making more than two applications of AMATHON herbicide and any other Group 2 herbicides within a single growing season unless mixed with an herbicide with a different site of action with an overlapping spectrum for the difficult-to-control weeds.
- Incorporate non-chemical weed control practices, including mechanical cultivation, crop rotation, cover crops and weed-free crop seeds, as part of an integrated weed control program.
- Use good agronomic principles that enhance crop development and crop competitiveness.
- Thoroughly clean plant residues from equipment before leaving fields suspected to contain resistant weeds.
- Manage weeds in and around fields, during and after harvest to reduce weed seed production.

INTEGRATED PEST MANAGEMENT

This product may be used as part of an Integrated Pest Management (IPM) program that can include biological, cultural, and genetic practices aimed at preventing economic pest damage. IPM principles and practices include field scouting or other detection methods, correct target pest identification, population monitoring, and treating when target pest populations reach locally determined action thresholds. Consult your state cooperative extension service, professional consultants or other qualified authorities to determine appropriate action treatment threshold levels for treating specific pest/crop systems in your area.

APPLICATION INFORMATION

WHEAT, BARLEY AND TRITICALE

APPLICATION TIMING

Apply after the crop is in the 2-leaf stage, but before the flag leaf is visible on wheat (except durum varieties of spring wheat), barley and triticale.

In durum varieties of spring wheat, apply only with 2,4-D and make applications after the crop is tillering but before the crop reaches the boot stage.

Apply AMATHON herbicide when all or most of the weeds have germinated. Annual broadleaf weeds must be past the cotyledon stage, actively growing, and less than 4" tall or wide.

CEREALS USE RATE

Use 0.33 oz (0.0094 lb ai/A Tribenuron methyl/0.0013 lb ai/A Metsulfuron methyl) AMATHON herbicide for those weeds listed under the "WEEDS CONTROLLED" section of this label. Make only one application of AMATHON herbicide per year.

Restriction:

DO NOT harvest within 45 days of the last application.

Use Area	Active Ingredient (AI)	Application Timing	Maximum Oz/A of Product per Single Application	Maximum lb ai/A per Single Application	Maximum Number of Applications per Year	Maximum Oz/A of Product per Year	Maximum lb ai/A of Product per Year	Pre-Harvest Interval, Days
Wheat (except durum varieties of spring wheat), barley and triticale	Tribenuron methyl	Apply in the 2-leaf stage but before the flag is visible	0.33	0.0094	1	0.33	0.0094	45
	Metsulfuron methyl			0.0013			0.0013	
Durum varieties of spring wheat	Tribenuron methyl	Apply only with 2,4-D and make applications after the crop is tillering but before the crop reaches the boot stage	0.33	0.0094	1	0.33	0.0094	45
	Metsulfuron methyl			0.0013			0.0013	

BURNDOWN - POST HARVEST, FALLOW, PRE-PLANT

APPLICATION TIMING

AMATHON herbicide may be used as a burndown treatment when the majority of weeds have emerged and are actively growing. AMATHON herbicide may be applied to crop stubble, as a fallow treatment, or as a pre-plant burndown prior to planting any crop. See "CROP ROTATION" for the minimum interval allowed between the burndown application and when a crop may be planted.

BURNDOWN USE RATE

Apply 0.33 oz (0.0094 lb ai/A Tribenuron methyl/0.0013 lb ai/A Metsulfuron methyl) AMATHON herbicide per acre as a burndown treatment prior to or shortly after planting wheat (including durum), barley or triticale (prior to emergence).

Make only one application of AMATHON herbicide per year.

See "CROP ROTATION" for the minimum interval allowed between the burndown application and when a crop may be planted.

RESTRICTIONS

Use Area	Active Ingredient (AI)	Application Timing	Maximum oz/A of Product per Single Application	Maximum lb ai/A per Single Application	Maximum Number of Applications per Year	Maximum oz/A of Product per Year	Maximum lb ai/A of Product per Year	Pre-Harvest Interval, Days
Burndown Post-Harvest, Fallow, Pre-Plant	Tribenuron methyl	Refer to the Crop Rotation Section	0.33	0.0093	1	0.33	0.0093	45
	Metsulfuron methyl			0.0012			0.0012	

SPRAY ADJUVANTS - ALL CROPS OR USES

Include a spray adjuvant with applications of AMATHON herbicide. In addition, an ammonium nitrogen fertilizer may be used.

Consult your Ag dealer or applicator, local FMC fact sheets and technical bulletins prior to using an adjuvant system. If another herbicide is tank mixed with AMATHON herbicide, select adjuvants authorized for use with both products. Products must contain only EPA-exempt ingredients.

NONIONIC SURFACTANT (NIS)

- Apply 0.06 to 0.5% volume/volume (0.5 pt to 4 pt per 100 gal of spray solution).
- Surfactant products must contain at least 60% nonionic surfactant with a hydrophilic/lipophilic balance (HLB) greater than 12.

CROP OIL CONCENTRATE (COC) - PETROLEUM OR MODIFIED SEED OIL (MSO)

- Apply at 1% v/v (1 gal per 100 gal spray solution) or 2% under arid conditions. MSO adjuvants may be used at 0.5% v/v if specified on local FMC product literature or service policies.
- Oil adjuvants must contain at least 80% high quality, petroleum (mineral) or modified vegetable seed oil with at least 15% surfactant emulsifiers.

SPECIAL ADJUVANT TYPES

- Combination adjuvant products may be used at doses that provide the required amount of NIS, COC, MSO and/or ammonium nitrogen fertilizer. Consult product literature for use rates and restrictions.
- In addition to the adjuvants specified above, other adjuvant types may be used if they provide the same functionality and have been evaluated and approved by FMC product management. Consult separate FMC technical bulletins for detailed information before using adjuvant types not specified on this label.

AMMONIUM NITROGEN FERTILIZER

- Use 2 qt/acre of a high-quality urea ammonium nitrate (UAN), including 28%N or 32%N, or 2 lb/acre of a spray-grade ammonium sulfate (AMS). Use 4 qt/acre UAN or 4 lb/acre AMS under arid conditions.
- See TANK MIXTURES With Liquid Nitrogen Fertilizer for instructions on using fertilizer as a carrier in place of water.

WEED CONTROL INFORMATION

WEEDS CONTROLLED

AMATHON herbicide effectively controls the following weeds when used according to label directions:

Black mustard	Hairy buttercup
Blue/Purple mustard	London Rocket
Bushy wallflower /Treacle mustard†	Marestail***†
Canola, volunteer (except Clearfield)**	Marshelder†
Canada thistle	Mayweed chamomile/Stinking chamomile/dog fennel (<i>Anthemis cotula L.</i>) ***†
Coast fiddleneck	Miners lettuce
Common Chickweed†	Narrowleaf hawksbeard** ***
Common Groundsel	Nightflowering catchfly
Common Lambsquarters†	Pineappleweed
Common Purslane	Poison hemlock***
Corn, Gromwell ***	Prickly lettuce**†
Corn spurry	Puncturevine
Cowcockle	Redroot pigweed†
Cressleaf groundsel *** (butterweed)	Russian thistle**†
Curly Dock ***	Shepherd's-purse
Dandelion	Slimleaf lambsquarters
Early whitlowgrass	Smallseed falseflax†
False chamomile/ Wild chamomile/Scentless chamomile (<i>Matricaria maritima L.</i>)	Tansymustard
Field pennycress	Tarweed fiddleneck
Flixweed†	Tumble/Jim Hill mustard ***
	Wild mustard†
	Wild parsnip***

WEEDS PARTIALLY CONTROLLED*

AMATHON herbicide partially controls the following weeds when used according to label directions:

Annual sowthistle	Pennsylvania smartweed
Cleavers, false	Purslane speedwell
Common cocklebur†	Prostrate knotweed
Common sunflower (volunteer)**†	Redmaids
Common vetch**	Redstem filaree***
Deadnettle	Small-flower buttercup
Eastern black nightshade†	Tumble pigweed
Hairy nightshade	White cockle
Hairy vetch***	Wild buckwheat
Henbit	Wild carrot
Narrowleaf hawksbeard	Wild garlic
	Wild radish**

* Partially controlled weeds exhibit a visual reduction in numbers as well as a significant loss of vigor. For better results, include a tankmix partner including 2,4-D, MCPA, bromoxynil or dicamba. See the "TANK MIXTURES" section of this label.

** See the Specific Weed Instructions section of this label for more information.

***2,4-D LVE addition required.

† Naturally occurring resistant biotypes are known to occur.

EXTENDED WEED CONTROL:

Used as directed, AMATHON herbicide will deliver extended control or suppression of cleavers, dandelion, narrow leaved hawk's beard and volunteer canola (including glyphosate resistant canola) beyond the control obtained with postemergence activity alone. Degree and duration of extended control will depend on environmental conditions at and following treatment and weed infestation levels.

SPECIFIC WEED INSTRUCTIONS

Narrowleaf hawksbeard: During the post harvest, fallow, and/or pre-plant burndown period, apply AMATHON herbicide in a tank mix with 1 to 2 pints of glyphosate per acre at labeled rates for postemergence control of narrowleaf hawksbeard. For wheat, apply AMATHON herbicide in a tank mix with 2,4-D for postemergence control of narrowleaf hawksbeard. Add 2,4-D at labeled rates. Apply this tank mix only in the spring when the wheat is fully tillered and before the jointing stage.

Canola, volunteer: In-crop applications of AMATHON herbicide will not control Clearfield varieties of volunteer canola. However, burndown applications made prior to crop emergence will provide effective control when tank mixed with glyphosate.

Russian thistle, Prickly lettuce: For best results, apply AMATHON herbicide in a tank mix with dicamba (including "Clarity") and 2,4-D or MCPA (ester or amine), or bromoxynil containing products (including "Bison"). Apply AMATHON herbicide in the spring when Russian thistle, and prickly lettuce are less than 2" tall or 2" across and are actively growing (refer to the Tank Mixtures section of this label).

Wild radish: For best results, apply AMATHON herbicide plus MCPA at labeled rates, plus 0.25% v/v nonionic surfactant (1 qt per 100 gal of spray solution) to wild radish rosettes less than 6" diameter, either in the fall or spring. Applications made later than 30 days after weed emergence will result in partial control. Fall applications must be made before plants harden-off.

Volunteer Sunflowers: Varieties resistant to SU and IMI products (like AMATHON herbicide, "Beyond", "Pursuit", "Raptor") are under development. For best results, use AMATHON herbicide in a tank mix with "Starane", "Colt + Salvo", "Colt + Sword", dicamba (including "Clarity") and 2,4-D or MCPA (ester or amine), or bromoxynil containing products (including "Bison").

TANK MIXTURES

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

AMATHON herbicide may be tank mixed with full or reduced rates of other herbicide, insecticide, and fungicide products registered for use in the specified crops. Consult tank mix partner labeling for rate and crop rotation restrictions. Read and follow all manufactures label instructions for the companion herbicide(s). **DO NOT** use a tank mix partner product if its label conflicts with the AMATHON herbicide label. Ensure the tank mix product is labeled for the same timing, method of application, adjuvants, and use restrictions as AMATHON herbicide, as well as other products used in the tank mixture. Read and follow all applicable use directions, precautions, and limitations specified on the respective product labels. Weed control and crop safety resulting from the use of tank mixtures not specifically noted on this label are the responsibility of the user.

WHEAT, BARLEY AND TRITICALE

With 2,4-D (amine or ester) or MCPA (amine or ester)

Tank mix AMATHON herbicide with 2,4-D and MCPA (preferably ester formulations) herbicides for use on wheat, barley and triticale. For best results, add 2,4-D or MCPA herbicides to the tank at labeled rates. Add 1 to 2 pt of nonionic surfactant per 100 gal of spray solution to the 3 way mixture, where necessary, as deemed by local guidance. Higher rates of 2,4-D or MCPA may be used, but **DO NOT** exceed the highest rate allowed by those respective labels. At higher rates, use of additional nonionic surfactant may not be needed, unless specified otherwise in the 2,4-D or MCPA label, or local guidance.

With 2,4-D or MCPA (amine or ester) and Dicamba (including "Clarity")

AMATHON herbicide may be applied in a 3-way tank mix with formulations of dicamba (including "Clarity") and 2,4-D or MCPA. Make applications of AMATHON herbicide + dicamba (including "Clarity") + 2,4-D or MCPA (ester or amine) at labeled rates. Use higher rates when weed infestation is heavy. Add 1 to 2 pt of nonionic surfactant per 100 gal of spray solution to the 3 way mixture, where necessary, as deemed by local guidance. Use of additional nonionic surfactant may not be needed with the higher phenoxy rates and ester phenoxy formulations. Consult the specific 2,4-D or MCPA and dicamba labels, or local guidance for more information. Apply this 3-way combination to winter wheat after the crop is tillering and prior to jointing (first node). In spring wheat (including durum), apply after the crop is tillering and before it exceeds the 5-leaf stage.

With Bromoxynil containing products (including, "Bison")

AMATHON herbicide may be tank mixed with bromoxynil containing herbicides registered for use on wheat, barley or triticale. For best results, add bromoxynil containing herbicides (including "Bison") to the tank at labeled rates. Tank mixes of AMATHON herbicide plus bromoxynil may result in reduced control of Canada thistle.

With fluroxypyr (including "Starane" brands)

AMATHON herbicide may be tank mixed with fluroxypyr containing herbicides for improved control of Kochia (2-4" tall) and other broadleaf weeds. For best results, add fluroxypyr containing herbicides (including "Starane") to the tank at labeled rates. 2,4-D and MCPA herbicides (preferably ester formulations) may be tank mixed with AMATHON herbicide plus "Starane".

With Postemergence Grass Herbicides

When used in tank mixture with "GoldSky", "Everest", or "Rimfire Max", AMATHON herbicide will result in improved control of yellow and green foxtail. Consult tank mix partner labeling for any adjuvant, rate, and grass weed height limitations, as reduced grass control may result when using tank mixtures with some WSSA Group 1 (ACCase) herbicides.

With Insecticides

AMATHON herbicide may be tank mixed or used sequentially with insecticides registered for use on cereal crops. However, under certain conditions (drought stress, or if the crop is in the 2 to 4 leaf stage), tank mixes or sequential applications of AMATHON herbicide with organophosphate insecticides (including "Lorsban") may produce temporary crop yellowing or, in severe cases, crop injury. The potential for crop injury is greatest when wide fluctuations in day/night temperatures occur just prior to or soon after application. Test these mixtures in a small area before treating large areas.

DO NOT apply AMATHON herbicide within 60 days of crop emergence where an organophosphate insecticide has been applied as an in-furrow treatment because crop injury may result.

DO NOT use AMATHON herbicide plus Malathion because crop injury may result.

GRAZING

Allow at least 7 days between application and grazing of treated forage. In addition, allow at least 7 days between application and feeding of forage (green chop) from treated areas to livestock. Allow at least 30 days between application and feeding of hay from treated areas to livestock. Allow at least 45 days between application and harvesting of grain. Harvested straw may be used for bedding and/or feed.

CROP ROTATION

Before using AMATHON herbicide carefully consider your crop rotation plans and options. For rotational flexibility, **DO NOT** treat all of your acres at the same time.

MINIMUM ROTATIONAL INTERVALS

Minimum rotation intervals* are determined by the rate of breakdown of AMATHON herbicide applied. AMATHON herbicide breakdown in the soil is affected by soil pH, presence of soil microorganisms, soil temperature, and soil moisture. Low soil pH, high soil temperature, and high soil moisture increase AMATHON herbicide breakdown in soil, while high soil pH, low soil temperature, and low soil moisture slow AMATHON herbicide breakdown. Of these 3 factors, only soil pH remains relatively constant. Soil temperature, and to a greater extent, soil moisture, can vary significantly from year to year and from area to area. For this reason, monitor soil temperatures and soil moisture regularly when considering crop rotations.

* The minimum rotation interval represents the period of time from the last application to the anticipated date of the next planting. Minimum rotation intervals must be extended 1 crop season if drought conditions prevail after application and before the rotational crop is planted.

SOIL pH LIMITATIONS

DO NOT use AMATHON herbicide on soils having a pH above 7.9, as extended soil residual activity could extend crop rotation intervals beyond normal. Under certain conditions, AMATHON herbicide could remain in the soil for 34 months or more, injuring wheat, barley or triticale. In addition, other crops planted in high-pH soils can be extremely sensitive to low concentrations of AMATHON herbicide.

CHECKING SOIL pH

Before using AMATHON herbicide, determine the soil pH of the areas of intended use. To obtain a representative pH value for the test area, take several 0” to 4” samples from different areas of the field and analyze them separately. Consult local extension publications for additional information on suggested soil sampling procedures.

TIME INTERVAL BEFORE PLANTING (MONTHS AFTER TREATMENT WITH AMATHON HERBICIDE)

Crop	Soil pH	Minimum Rotation Interval (months)
Sorghum, Grain	7.9 or lower	11
Peas, Dry/Green	7.9 or lower	11
Canola	7.9 or lower	11
Flax	7.9 or lower	11
Lentils	6.8 or lower 6.9 to 7.9	11 22
Alfalfa	6.8 or lower 6.9 to 7.9	11 22
Beans, Dry	6.8 or lower 6.9 to 7.9	11 22
Sunflower	7.9 or lower	11
Field Corn	7.9 or lower	12
Soybean	7.9 or lower	12
Wheat (spring, durum or winter), triticale or spring barley	7.9 or lower	1 day

Rotation Intervals for crops not covered above - The minimum rotation interval is 34 months with at least 28” of cumulative precipitation during the period:

- to any major field crop not listed (See the Rotation Intervals table)
- if the soil pH is not in the specified range
- if the use rate applied is not specified in the table
- or if the minimum cumulative precipitation has not occurred since application. A field bioassay must be successfully completed before rotation to any minor crops (as determined by the USDA criteria). See section on Field Bioassay for further information.

FIELD BIOASSAY

A field bioassay is necessary if crops other than wheat, barley or those listed on this label are to be planted on land previously treated with AMATHON herbicide. To conduct a field bioassay, grow test strips of the crop or crops you plan to grow the following year in fields previously treated with AMATHON herbicide. Crop response to the bioassay will indicate whether or not to rotate to the crop(s) grown in the test strips.

If a field bioassay is planned, check with your local FMC representative for information detailing field bioassay procedure.

APPLICATION INFORMATION

PRODUCT MEASUREMENT

AMATHON herbicide may be measured using the AMATHON herbicide volumetric measuring cylinder provided by FMC. The degree of accuracy of this cylinder varies by $\pm 7.5\%$. For more precise measurement, use scales calibrated in ounces.

MIXING INSTRUCTIONS

1. Fill the tank 1/4 to 1/3 full of water.
2. While agitating, add the required amount of AMATHON herbicide.
3. Continue agitation until the AMATHON herbicide is fully dispersed, at least 5 minutes.
4. Once the AMATHON herbicide is fully dispersed, maintain agitation and continue filling tank with water. Thoroughly mix AMATHON herbicide with water before adding any other material.
5. As the tank is filling, add tank mix partners (if desired) then add the required volume of spray adjuvant. Always add spray adjuvant last. Antifoaming agents may be used. **DO NOT** use with spray additives that alter the pH of the spray solution below pH 6.0 as rapid product degradation can occur. Spray solutions of pH 7.0 and higher allow for optimum stability of AMATHON herbicide.
6. If the mixture is not continuously agitated, settling will occur. If settling occurs, thoroughly re-agitate before using.
7. Apply AMATHON herbicide spray mixture within 24 hours of mixing to avoid product degradation.
8. If AMATHON herbicide and a tank mix partner are to be applied in multiple loads, pre-slurry the AMATHON herbicide in clean water prior to adding to the tank. This will prevent the tank mix partner from interfering with the dissolution of the AMATHON herbicide.

APPLICATION METHOD

GROUND APPLICATION

For optimum spray distribution and thorough coverage, use flat-fan or low-volume flood nozzles.

- For flat-fan nozzles, use a spray volume of at least 5 gal per acre (GPA).
- For flood nozzles on 30" spacing, use flood nozzles no larger than TK10 (or the equivalent), a pressure of at least 30 psi and a spray volume of at least 10 GPA only. For 40" nozzle spacing, use at least 13 GPA; for 60" spacing use at least 20 GPA. It is essential to overlap the nozzles 100% for all spacings.
- **DO NOT** use "Raindrop RA" nozzles for AMATHON herbicide applications, as weed control performance may be reduced.
- Use screens that are 50-mesh or larger.

AERIAL APPLICATION

Use at least 2 GPA. In Idaho, Oregon and Utah use at least 3 GPA. **DO NOT** apply AMATHON herbicide by air in the state of New York.

See the **Spray Drift Management** section of this label.

APPLICATIONS WITH LIQUID NITROGEN SOLUTION FERTILIZER

Liquid nitrogen fertilizer solutions may be used as a carrier in place of water. Run a tank mix compatibility test before mixing AMATHON herbicide in fertilizer solution. AMATHON herbicide must first be slurried with water and then added to liquid nitrogen solutions (e.g., 28-0-0, 32-0-0). Ensure that the agitator is running while the AMATHON herbicide is added. Use of this mixture may result in temporary crop yellowing and stunting.

If using low rates of liquid nitrogen fertilizer in the spray solution (less than 50% of the spray solution volume), the addition of surfactant is necessary. Add surfactant at 0.5 pt - 1 qt per 100 gal of spray solution (0.06 -0.25% v/v) based on local guidance.

When using high rates of liquid nitrogen fertilizer solution in the spray solution, adding surfactant increases the risk of crop injury. If 2,4-D or MCPA is included with AMATHON herbicide and fertilizer mixture, ester formulations tend to be more

compatible (see manufacturer's label). Additional surfactant may not be needed when using AMATHON herbicide in tank mix with 2,4-D ester or MCPA ester and liquid nitrogen fertilizer solutions. Consult your agricultural dealer, consultant, field advisor, or FMC representative for guidance before adding an adjuvant to these tank mixtures.

Note: In certain areas east of the Mississippi river unacceptable crop response may occur with use of straight or dilute nitrogen fertilizer carrier solutions where cold temperatures or widely fluctuating day/night temperatures exist. In these areas consult your agricultural dealer, consultant, field advisor, or FMC representative for guidance before using nitrogen fertilizer carrier solutions.

DO NOT use low rates of liquid nitrogen fertilizer solution as a substitute for a surfactant. Liquid nitrogen fertilizer solutions that contain sulfur can increase crop response.

DO NOT use with liquid fertilizer solutions with a pH less than 3.0.

SPRAY EQUIPMENT

Be sure to calibrate air or ground equipment properly before application. Select a spray volume and delivery system that will ensure thorough coverage and a uniform spray pattern with minimum drift. Use higher spray volumes to obtain better coverage when crop canopy is dense. Avoid swath overlapping, and shut off spray booms while starting, turning, slowing, or stopping, to avoid injury to the crop.

DO NOT make applications using equipment and/or spray volumes or during weather conditions that might cause spray to drift onto nontarget sites. For additional information on spray drift refer to Spray Drift Management section of label.

Continuous agitation is not required to keep AMATHON herbicide in suspension but may be required to keep tank mix partners in solution or suspension. Refer to tank mix partner labels for additional information.

BEFORE SPRAYING AMATHON HERBICIDE

The spray equipment must be clean before AMATHON herbicide is sprayed. Follow the cleanup procedures specified on the labels of the previously applied products. If no directions are provided, follow the six steps outlined in the After Spraying AMATHON herbicide section of this label.

AT THE END OF THE DAY

At the end of each day of spraying, when multiple loads of AMATHON herbicide are applied, rinse and then partially fill the interior of the tank with fresh water, and flush the boom and hoses. This will prevent the buildup of dried pesticide deposits which can accumulate in the application equipment.

AFTER SPRAYING AMATHON herbicide AND BEFORE SPRAYING CROPS OTHER THAN WHEAT, BARLEY AND TRITICALE

To avoid subsequent injury to desirable crops, thoroughly clean all mixing and spray equipment immediately following applications of AMATHON herbicide as follows:

1. Empty the tank and drain the sump completely.
2. Spray the tank walls with clean water using a minimum volume of 10% of the tank volume. Circulate the water through the lines, including all by-pass lines, for at least two minutes. Flush the boom well and empty the sprayer. Completely drain the sump.
3. Repeat step 2.
4. Remove the nozzles and screens and clean separately in a bucket containing water. The rinsate solution may be applied back to the crop(s) specified on this label. **DO NOT** exceed the maximum-labeled use rate. If cleaners are used, consult the cleaner label for rinsate disposal instructions. If no instructions are given, dispose of the rinsate on site or at an approved waste disposal facility.

Notes:

1. Steam-cleaning aerial spray tanks is advised to facilitate the removal of any caked deposits.
2. When AMATHON herbicide is tank mixed with other pesticides, examine all cleanout procedures for each product and follow the most rigorous procedure.
3. Follow any pre-cleanout guidelines directed on other product labels.

MANDATORY SPRAY DRIFT MANAGEMENT

Ground Boom Applications:

- Apply with the nozzle height advised by the manufacturer, but no more than 3 feet above the ground or crop canopy.
- For applications prior to the emergence of crops and target weeds, applicators are required to use a Coarse or coarser droplet size (ASABE S572.1).
- For all other applications, applicators are required to use a Medium or coarser droplet size (ASABE S572.1).
- **DO NOT** apply when wind speeds exceed 10 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

Aerial Applications:

- **DO NOT** release spray at a height greater than 10 feet above the vegetative canopy, unless a greater application height is necessary for pilot safety.
- For applications prior to the emergence of crops and target weeds, applicators are required to use a Coarse or coarser droplet size (ASABE S572.1).
- For all other applications, applicators are required to use a Medium or coarser droplet size (ASABE S572.1).
- The boom length must not exceed 65% of the wingspan for airplanes or 75% of the rotor blade diameter for helicopters.
- Applicators must use one-half swath displacement upwind at the downwind edge of the field.
- Nozzles must be oriented so the spray is directed toward the back of the aircraft.
- **DO NOT** apply when wind speeds exceed 10 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

Boom-less Ground Applications:

- Applicators are required to use a Medium or coarser droplet size (ASABE S572.1) for all applications.
- **DO NOT** apply when wind speeds exceed 10 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

SPRAY DRIFT MANAGEMENT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.

BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size – Ground Boom

- Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure - Use the lowest advised spray pressure for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size – Aircraft

- Adjust Nozzles - Follow nozzle manufacturer's directions for setting up nozzles. Generally, to reduce fine droplets, nozzles must be oriented parallel with the airflow in flight.

BOOM HEIGHT – Ground Boom

Use the lowest boom height that is compatible with the spray nozzles that will provide uniform coverage. For ground equipment, the boom must remain level with the crop and have minimal bounce.

RELEASE HEIGHT - Aircraft

Higher release heights increase the potential for spray drift. When applying aurally to crops, do not release spray at a height greater than 10 ft above the crop canopy, unless a greater application height is necessary for pilot safety.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.

Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

HANDHELD TECHNOLOGY APPLICATIONS:

- Take precautions to minimize spray drift.

BOOM-LESS GROUND APPLICATIONS:

- **Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.**

DRIFT CONTROL ADDITIVES

Using product compatible drift control additives can reduce drift potential. When a drift control additive is used, read and carefully observe cautionary statements and all other information on the additive's label. If using an additive that increases viscosity, ensure that the nozzles and other application equipment will function properly with a viscous spray solution. Preferred drift control additives have been certified by the Council of Producers & Distributors of Agrotechnology (CPDA).

IDENTIFICATION INFORMATION FOR PRODUCTS REFERENCED IN THIS LABEL**REGISTERED PRODUCTS REFERENCED IN THIS LABEL FOR TANK MIXTURES
OR MENTIONED FOR OTHER REASONS**

Product Name	Active Ingredient(s)	EPA Registration Number
Beyond® herbicide	Imazamox	241-441
Bison® herbicide	MCPA + Bromoxynil	9779-347
Clarity® Herbicide	Dicamba	7969-137
Colt®+Salvo® Herbicide	2,4-D + Fluroxypyr	34704-1010
Colt®+Sword® Herbicide	MCPA + Fluroxypyr	34704-1011
Discover® NG Herbicide	Clodinafop-Propargyl	100-1173
Everest® 3.0 AG	Flucarbazone-Sodium	66330-433
Everest® 3.0 Herbicide	Flucarbazone-Sodium	66330-429
Goldsky® herbicide	florasulam, fluroxypyr, pyroxsulam	62719-582
Lorsban® Advanced Insecticide	Chlorpyrifos	62719-591
Lorsban® 15G Granular Insecticide	Chlorpyrifos	62719-34
Lorsban® 50W in Water Soluble Packets Insecticide	Chlorpyrifos	62719-221
Lorsban®-4E Insecticide	Chlorpyrifos	62719-220
Pursuit® herbicide	Imazethapyr	241-310
Raptor® herbicide	Imazamox	241-379
Rimfire® Max herbicide	propoxycarbazone-sodium, mesosulfuron-methyl	264-1099
Salvo® herbicide	2,4-D	34704-609
Starane® Flex Herbicide	Florasulam + Fluroxypyr	62719-604
Starane® NXT Herbicide	Fluroxypyr + Bromoxynil	62719-557
Starane® Ultra Herbicide	fluroxypyr	62719-577

PESTICIDE STORAGE AND DISPOSAL

Pesticide Storage: Store the product in original container only. **DO NOT** contaminate water, other pesticides, fertilizer, food, or feed in storage. Store in a cool, dry place.

Product Disposal: **DO NOT** contaminate water, food, or feed by disposal. Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING:

Refer to the Net Contents section of this product's labeling for the applicable "Nonrefillable Container" or "Refillable Container" designation.

Nonrefillable Plastic and Metal Containers (Capacity Equal to or Less Than 50 Pounds): Nonrefillable container. **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. **DO NOT** burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities

Nonrefillable Plastic and Metal Containers (Capacity Greater Than 50 Pounds): Nonrefillable container. **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. **DO NOT** burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Nonrefillable Plastic and Metal Containers, e.g., Intermediate Bulk Containers [IBC] (Size or Shape Too Large to be Tipped, Rolled or Turned Upside Down): Nonrefillable container. **DO NOT** reuse or refill this container. Clean container promptly after emptying the contents from this container into application equipment or mix tank and before final disposal using the following pressure rinsing procedure. Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that the water spray thoroughly covers the top, bottom and all sides inside the container. The nozzle manufacturer generally provides instructions for the appropriate spray pressure, spray duration and/or spray volume. If the manufacturer's instructions are not available, pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume. Drain, pour or pump rinsate into application equipment or rinsate collection system. Repeat this pressure rinsing procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Nonrefillable Paper or Plastic Bags, Fiber Sacks including Flexible Intermediate Bulk Containers (FIBC) or Fiber Drums With Liners: Nonrefillable container. **DO NOT** reuse or refill this container. Completely empty paper or plastic bag, fiber sack or drum liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. Then offer for recycling if available or dispose of empty paper or plastic bag, fiber sack or fiber drum and liner in a sanitary landfill, or by incineration. **DO NOT** burn, unless allowed by state and local ordinances.

Refillable Fiber Drums With Liners: Refillable container (fiber drum only). Refilling Fiber Drum: Refill this fiber drum with AMATHON herbicide (with TOTALSOL® soluble granules) containing metsulfuron methyl and tribenuron methyl only. **DO NOT** reuse this fiber drum for any other purpose. Cleaning before refilling is the responsibility of the refiller. Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. Disposing of Fiber Drum and/or Liner: **DO NOT** reuse this fiber drum for any other purpose other than refilling (see preceding). Cleaning the container (liner and/or fiber drum) before final disposal is the responsibility of the person disposing of the container. Offer the liner for recycling if available or dispose of liner in a sanitary landfill, or by incineration. **DO NOT** burn, unless allowed by state and local ordinances. If drum is contaminated and cannot be reused, dispose of it in the manner required for its liner. To clean the fiber drum before final disposal, completely empty the fiber drum by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. Then offer the fiber drum for recycling if available or dispose of in a sanitary landfill, or by incineration. **DO NOT** burn, unless allowed by state and local ordinances.

All Other Refillable Containers: Refillable container. Refilling Container: Refill this container with AMATHON herbicide (with TOTALSOL® soluble granules) containing metsulfuron methyl and tribenuron methyl only. **DO NOT** reuse this container for any other purpose. Cleaning before refilling is the responsibility of the refiller. Prior to refilling, inspect carefully for damage including cracks, punctures, abrasions, worn out threads and closure devices. If damage is found, **DO NOT** use the container, contact FMC at the number below for instructions. Check for leaks after refilling and before transporting. If leaks are found, **DO NOT** reuse or transport container, contact FMC at the number below for instructions. Disposing of Container: **DO NOT** reuse this container for any other purpose other than refilling (see preceding). Cleaning the container before final disposal is the responsibility of the person disposing of the container. To clean the container before final disposal, use the following pressure rinsing procedure. Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that the water spray thoroughly covers the top, bottom and all sides inside the container. The nozzle manufacturer generally provides instructions for the appropriate spray pressure, spray duration and/or spray volume. If the manufacturer's instructions are not available, pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume. Drain, pour or pump rinsate into application equipment or rinsate collection system. Repeat this pressure rinsing procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. **DO NOT** burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Outer Foil Pouches of Water Soluble Packets (WSP): Nonrefillable container. **DO NOT** reuse or refill this container. Offer for recycling if available or, dispose of the empty outer foil pouch in the trash as long as WSP is unbroken. If the outer pouch contacts the formulated product in any way, the pouch must be triple rinsed with clean water. Add the rinsate to the spray tank and dispose of the outer pouch as described previously.

DO NOT transport if this container is damaged or leaking. If the container is damaged, leaking or obsolete, or in the event of a major spill, fire or other emergency, contact CHEMTREC (Transportation and Spills) at 1-800-424-9300, day or night.

NOTICE TO BUYER: Purchase of this material does not confer any rights under patents of countries outside of the United States.

FMC, Amathon™, and TotalSol®, are trademarks or registered trademarks of FMC Corporation or an affiliate.

"Pursuit", "Raptor", "Beyond" , and "Clarity are registered trademarks of BASF Corporation.

Rimfire is a registered trademark of Bayer Corporation.

"Raindrop RA" is a registered trademark of Delavan

Everest is a registered trademark of UPL NA Inc..

Goldsky, Lorsban and Starane are registered trademarks of Corteva Agriscience

Salvo is a registered trademark of Loveland Products, Inc.

Bison is a registered trademark of WinfieldUnited

'Colt® + Salvo®' and Colt® + Sword ® are registered trademarks of Loveland Products

D – 4471 071020

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

Notice: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions beyond the control of FMC or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and, to the extent consistent with applicable law, Buyer and User agree to hold FMC and Seller harmless for any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the Directions for Use when used in accordance with the directions under normal conditions of use. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, FMC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE, NOR ANY OTHER EXPRESS OR IMPLIED WARRANTIES WITH RESPECT TO THE SELECTION, PURCHASE, OR USE OF THIS PRODUCT. Any warranties, express or implied, having been made are inapplicable if this product has been used contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to (or beyond the control of) Seller or FMC, and, to the extent permitted by applicable law, Buyer assumes the risk of any such use.

To the extent consistent with applicable law, FMC or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF FMC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF FMC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

This Condition of Sale and Limitation of Warranty and Liability may not be amended by any oral or written agreement.