



HOGS: To control Lice, Sarcoptic Mange, Mix 1 gallon with 150 gallons of water. Spray or wash animals thoroughly. Do not apply within 28 days of slaughter. Do not use on animals under 3 months old. For Mange control, repeat treatment at 7-10 day intervals for two or more applications, as needed. In cases of light infestation one application may be adequate. Repeat as necessary for lice.

WARNING

MAY BE FATAL IF SWALLOWED. Do not breathe spray mist. Do not get in eyes, on skin or clothing. Wash thoroughly after using. Do not contaminate feed or food-stuffs. Do not use or store near heat or open flame.

Toxic to fish and wildlife. Do not contaminate streams, lakes, or ponds with this material. Destroy empty containers and wastes. Do not contaminate water by cleaning of equipment or disposal of wastes.

ANTIDOTE: Give one tablespoonful of salt in a glass of warm water to induce vomiting. Repeat until vomit fluid is clear. Call a physician immediately.

Avoid spraying and dipping animals during cold, stormy weather. Avoid treating sick or overheated animals. Feed and water animals before treating. Spraying is recommended for young animals. If young animals must be dipped, individual hand immersion is essential to avoid swallowing of dip solution.

Do not use on dogs, cats, or other household pets or on poultry, or in poultry houses. Use only as directed on this label.

Stock Tax forms a stable emulsion with water, but some settling may occur, especially in hard water. Thoroughly agitate mixture before dipping or spraying.

27115

E.P.A. Reg. No. 299-57



STOCK TOX

TOXAPHENE EMULSIFIABLE CONCENTRATE

FOR USE AS A LIVESTOCK SPRAY OR DIP
TO CONTROL

SCABIES (PSOROPTIC, CHLORIOPTIC AND SARCOPTIC MANGE)
HORNFLIES, LICE, TICKS

CONTENTS: 1 GALLON

WARNING: Keep Out of Reach of Children.

SEE BACK PANEL FOR OTHER WARNINGS.

ACTIVE INGREDIENTS:

* Toxaphene	62.0%
Petroleum Distillate	32.0%
INERT INGREDIENTS:	6.0%

* Technical Chlorinated Camphene (Chlorine content 67% to 69%)

DIRECTIONS

BEEF CATTLE AND HORSES: To control Scabies (Psoroptic, Sarcoptic and Chlorioptic Mange), Lone Star Ticks, Cattle Ticks, Winter Ticks, Gulf Coast Ticks, Hornflies, Cattle Lice, Ear Ticks: Mix 1 gallon with 150 gallons of water. Thoroughly wet animals by spraying or dipping. For ear ticks, use mist sprayer and apply into each ear and to head and neck. Repeat treatment at 7 to 10 day intervals for two or more applications as needed for scabies control. Repeat as necessary for control of other insects. Do not apply within 28 days of slaughter. Do not use on dairy cattle. Do not use on animals under 3 months old. Do not dip animals under 6 months old. If calves between 3 and 6 months old must be treated, apply only as a spray.

BACK RUBBERS: As an aid in the control of Hornflies and Lice on beef cattle: Mix 1 quart with 5 gallons of kerosene or diesel fuel. Fill tank of back-rubber device or soak burlap roll with solution so it is thoroughly wet but not to point of dripping. Recharge as necessary. Do not permit animals access to treatment within 28 days of slaughter. Do not permit access to backrubbers by dairy animals or calves under 6 months old.

SHEEP AND GOATS: To control Psoroptic Sheep Scab, Fleeceworms, Hornflies, Lone Star Ticks, Ear Ticks: Mix 1 gallon with 150 gallons of water. Thoroughly wet animals by spraying or dipping. For ear ticks, use a mist sprayer and apply into each ear and to head and neck. Repeat treatment at 7 to 10 day intervals for two or more applications as needed for scabies control. Repeat as necessary for other insects.

To control Lice and Sheep Ticks (Keds): As a dip, use 1 gallon with 800 gallons of water. For spraying, 1 gallon with 200 gallons of water will give best results. Thoroughly wet animals. Repeat as necessary.

Do not apply within 28 days of slaughter. Do not use on dairy goats. Do not use on animals under 3 months old.

606 W. Main St.

C. J. Martin Company

Nacogdoches, Texas 75961