



Longer control is obtained if MARTIN'S BEEF CATTLE DUST is rubbed into the hair rather than being thrown onto the animal.

For the best results start your fly control program before the hornfly population reaches great numbers and maintain applications as necessary.

**WARNING:** Harmful if swallowed. May be absorbed through the skin. Avoid breathing of dust. Avoid contact with eyes, skin, and clothing. Wash thoroughly after using.

**CAUTION:** Do not apply to dairy animals. Do not store near feed and food products. Do not apply within 30 days of slaughter. Do not treat animals under 3 months of age. Do not use on dogs, cats, or other household pets. Do not use on poultry nor in poultry houses. Do not treat sows within 2 weeks of farrowing nor for at least 3 weeks thereafter.

Toxic to fish and wildlife; do not contaminate streams, lakes, or ponds with this material. Destroy empty containers and spillage.

U.S.D.A. Reg. No. 299-118

76138

# BEEF CATTLE DUST

For Use on

**Beef Cattle, Sheep, Goats, and Hogs**

CONTENTS: 2 POUNDS

**CAUTION: Keep Out of Reach of Children**

SEE BACK PANEL FOR OTHER CAUTIONS.

## DIRECTIONS:

**BEEF CATTLE:** For control of hornflies, ticks, and lice on beef cattle treat animals by rubbing dust lightly into hair. Treat animals thoroughly. Repeat treatment as needed.

**SWINE:** For control of hog lice apply dust to hogs, pens, and litter. Treat animals thoroughly. One treatment may be sufficient. Repeat in 10 days or thereafter as needed.

**SHEEP AND GOATS:** For control of lice and keas on sheep and goats, treat animals thoroughly, preferably following shearing. Repeat as needed.

**HORN FLY CONTROL:** Normal hornfly control periods to be expected with MARTIN'S BEEF CATTLE DUST are 1 1/2-4 weeks depending upon the severity of the existing infestation and the weather. When hornflies are allowed to reach great numbers before control measures are taken, it may become necessary to make a repeat application a few days after the first application.

**ACTIVE INGREDIENTS:**

Teraphene (Chlorinated Camphene containing on to 69 percent chlorine) . . . . . 5%  
Lindane (Gamma Isomer of Benzene Hexachloride) . . . . . 1%

**INERT INGREDIENTS:**

100%

299

**Austin, Texas**

*E. J. Martin & Sons* incorporated

**Austin, Texas**



Longer control is obtained if MARTIN'S BEEF CATTLE DUST is rubbed into the hair rather than being thrown onto the animal.

For the best results start your fly control program before the hornfly population reaches great numbers and maintain applications as necessary.

**WARNING:** Harmful if swallowed. May be absorbed through the skin. Avoid breathing of dust. Avoid contact with eyes, skin, and clothing. Wash thoroughly after using.

**CAUTION:** Do not apply to dairy animals. Do not store near feed and food products. Do not apply within 30 days of slaughter. Do not treat animals under 3 months of age. Do not use on dogs, cats, or other household pets. Do not use on poultry nor in poultry houses. Do not treat sows within 2 weeks of farrowing nor for at least 3 weeks thereafter.

Toxic to fish and wildlife; do not contaminate streams, lakes, or ponds with this material. Destroy empty containers and spillage.

U.S.D.A. Reg. No. 299-118

76138

# BEEF CATTLE DUST

For Use on

**Beef Cattle, Sheep, Goats, and Hogs**

CONTENTS: 2 POUNDS

**CAUTION: Keep Out of Reach of Children**

SEE BACK PANEL FOR OTHER CAUTIONS.

## DIRECTIONS:

**BEEF CATTLE:** For control of hornflies, ticks, and lice on beef cattle treat animals by rubbing dust lightly into hair. Treat animals thoroughly. Repeat treatment as needed.

**SWINE:** For control of hog lice apply dust to hogs, pens, and litter. Treat animals thoroughly. One treatment may be sufficient. Repeat in 10 days or thereafter as needed.

**SHEEP AND GOATS:** For control of lice and keds on sheep and goats, treat animals thoroughly, preferably following shearing. Repeat as needed.

**HORN FLY CONTROL:** Normal hornfly control periods to be expected with MARTIN'S BEEF CATTLE DUST are 1 1/2-4 weeks depending upon the severity of the existing infestation and the weather. When hornflies are allowed to reach great numbers before control measures are taken, it may become necessary to make a repeat application a few days after the first application.

ACTIVE INGREDIENTS:  
Toxaphene (Chlorinated Camphene containing 67 to 69 percent chlorine) 57.15  
Lindane (Gamma Isomer of Benzene Hexachloride) 949.  
INERT INGREDIENTS: 1007.

Austin, Texas

*E. J. Martin & Sons* incorporated

Austin, Texas