



7 1/2 %

SEVIN DUST

Ready To Use In:
POULTRY HOUSE, LITTER,
OR BY DIRECT HAND APPLICATION
TO INDIVIDUAL BIRDS

See Direction Panel

CAUTION: KEEP OUT OF REACH OF CHILDREN.

See Side Panels for Other Cautions.

NET WEIGHT 50 POUNDS

® Trademark of Union Carbide Corporation for 1-naphthyl N-methylcarbamate.

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methylcarbamate) 7.5%

INERT INGREDIENTS 92.5%

DIRECTIONS

POULTRY

Litter Treatment: Apply 1 pound to 60 square feet to control Northern Fowl Mite, Chicken Mite, Lice, Fleas and Bed Bugs. Use a scoop and scatter by hand, or apply by hand duster. Wear gloves when spreading by hand. Avoid contamination of nests, eggs, feeding and watering troughs. Do not house birds in treated houses within 7 days of slaughter.

HAND APPLICATION TO BIRDS

Use 10 ounces to treat 100 birds by means of a shaker can or hand duster to control Northern Fowl Mite, Lice, and Fleas. Direct dust to vent and fluff areas. Do not contaminate feed or drinking water.

The application of this material cannot be repeated within 4 weeks of each treatment. Hand application to birds is a supplement to litter treatment for control of these pests. Do not apply to poultry within 7 days of slaughter.

CAUTION: HARMFUL IF SWALLOWED OR INHALED. Do not take internally. Avoid breathing dust. Avoid contact with skin. Wash thoroughly after handling. Change contaminated clothing daily. Avoid contamination of feed and foodstuffs.

NOTE FOR PHYSICIAN: Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is an antidote.

Texas Livestock Remedy Registration
Certificate No. 91

0794

U.S.D.A. Reg. No. 299-155

17526

06 W. Main St.

C. J. Martin Company

Nacogdoches, Texas 75961