DIRECTIONS

TO CLEAN:


To provide direct contact, first remove the water from the bowl by forcing it over the trap with a swab. Hold the bottle over the bowl and apply two to four ounces to the swab. Clean the entire bowl thoroughly, especially up under the rim the water outlets, flush toilet and rinse out mop.

TO DISINFECT:

Pour 10 ounces directly into the water remaining in the clean bowl. Then swab the entire bowl and seat, and cover thoroughly with this solution. After disinfecting, rinse seat and bowl well.

TO CLEAN AND DISINFECT URINALS:

Saturate swab thoroughly, holding the bottle over the urinal. Clean the urinal by wetting the entire surface. Particular attention should be paid to the grooves along the sides, and the water outlets at the top. After the surface looks clean, remove screen at the bottom of the urinal and pour two to five ounces on the edge and into the drain. Allow to stand two to three minutes. Then swab the inside pipe thoroughly. Replace screen and flush well.



DANGER POISON: KEEP OUT OF REACH MAY BE FATAL IF SWALLOWED, CORROSIVE, DO NOT BREATHE VAPOR OR FUMES, PRODUCES CHEMICAL BURNS, DO NOT GET IN EYES, ON SKIN, OR CLOTHING, SEE ANTIDOTE STATEMENT ON SIDE PANEL.

NET 32 FLUID OZ. (1 QUART)

A CONCENTRATED SUDSING WHITE EMULSION THAT WILL MAKE TOILET BOWLS CLEAN AND SANITARY

BEFORE DISCARDING THIS CONTAINER RINSE THOROUGHLY WITH WATER.

ANTIDOTE:

INTERNAL—Call a physician immediately. Drink a teaspoonful or more of magnesia, chalk, or small pieces of soap, softened with water in milk or raw egg white.

EXTERNAL—Eyes. Wash with water for 15 minutes. Get prompt medical attention. Wipe off acid gently, immediately flood the surface with water, using soap freely, then cover with moist magnesia or baking soda.

USDA Reg. No. 334-286

MANUFACTURED BY
PRODUCTS COMPANY

CHICAGO, ILL. 60609

ACCEPTED