

PM 25
312-332
P 1 3

10% BROMACIL PELLETS WEED KILLER

ACTIVE INGREDIENT:
Bromacil (5-bromo-3-sec-butyl-6-methyluracil)

INERT INGREDIENTS
U.S. Patents 3,195,974 & 3,352,862

Under the Federal Insecticide, Fungicide, and Rodenticide Act
as amended for the period
ended under EPA Reg. No. 333-333
OCT - 4 1985
ACCEPTED

Keep out of reach of children
PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION! MAY IRRITATE EYES, NOSE, THROAT, AND SKIN.

Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling.

STATEMENT OF PRACTICAL TREATMENT

- If in eyes:** Flush with plenty of water. Get medical attention if irritation persists
- If on skin:** Wash with plenty of soap and water. Get medical attention if irritation persists

ENVIRONMENTAL HAZARDS

Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes.

IMPORTANT — Injury to or loss of desirable trees or other plants may result from failure to observe the following: do not apply (or clean equipment) on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. Do not use on lawns, walks, driveways, tennis courts, or similar areas. Prevent drift of product to desirable plants. Keep from contact with fertilizers, insecticides, fungicides and seeds. Thoroughly clean application equipment after use.

GENERAL INFORMATION — 10% Bromacil Pellets Weed Killer is a pelleted product for dry application to the soil for the control of brush and many annual and perennial weeds. Effects are slow to appear and will not become apparent until the chemical has been activated by moisture. The degree of control and duration of effect will vary with the amount of chemical applied, soil type, rainfall, and other conditions.

10% Bromacil Pellets is noncorrosive to equipment, nonvolatile, and nonflammable

NET 50 POUNDS

AG 697 0585 MADE FOR Made in U.S.A. Printed in U.S.A.
JOHN DOE CHEMICAL CO. • ANYPLACE IN THE U.S.A.

rip-
fon
der
ber-
ther
s as
y: of
n. In
nect
uch
s of
ther

com-
urer's

use of
teller.
se.

icrop-
strial
mer
ches.
tation
me.

irable
t use
gation
table
adja-

ch as
wild
mer
1/2 to 1
basal
spots
and is

to 96
20 to
berry

PANEL.

pg 2 of 2

NOTICE OF WARRANTY

Seller warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of the seller. In no case shall the seller be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the buyer. Seller makes no warranties of merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

10% Bromacil Pellets should be used only in accordance with recommendations on this label, or in separate published manufacturer's recommendations available through local dealers.

Seller will not be responsible for losses or damages resulting from use of this product in any manner not specifically recommended by seller. User assumes all risk associated with such nonrecommended use.

BRUSH CONTROL: To control undesirable woody plants on noncrop-land areas such as railroad right-of-ways, storage areas, industrial plant sites and drainage ditches: apply pellets in spring or early summer as a broadcast or basal (spot) treatment; for use on drainage ditches, apply as a basal (spot) treatment only. Growth of herbaceous vegetation will be prevented in the treated area for an extended period of time.

Note: For effective brush control and prevention of damage to desirable vegetation: do not apply to brush standing in water; do not use water from treated ditches for irrigation; do not use in irrigation ditches nor on right-of-ways or other sites where marketable timber or other desirable trees or shrubs are immediately adjacent to the treated area.

Basal (Spot) Treatment: To control undesirable woody plants such as cottonwood, hackberry, maple, oak, poplar, red bud, sweetgum, wild cherry, willow and winged elm: apply pellets in spring or early summer by spreading 1 to 2 tablespoonfuls on the ground (cover an area 1/2 to 1 square foot) at the base of each brush stem or clump 2 to 4 inches basal diameter; on larger clumps apply 1 tablespoonful to each of 3 to 4 spots around the base of the brush. Apply to uphill side of plant if ground is not level.

Broadcast Treatment: Apply in spring or early summer: use 60 to 96 pounds per acre to control oak, willow, sweetgum and pine; use 120 to 240 pounds per acre to control American elm, winged elm, hackberry and cottonwood.

CONTINUED ON RIGHT PANEL.

10% BI

ACTIVE INGREDIENT:
Bromacil (5-bromo-3-sec-
INERT INGREDIENTS

U.S. Patents 3,235,357 & 3,252,862

HAZ CAUTION

**Avoid contact with skin,
water after handling.**

If in eyes: Flush
If on skin: Wash

Do not apply directly to water
disposal of wastes.

IMPORTANT — Injury to plants, or on areas where washed or moved into courts, or similar areas. Do not use fertilizers, insecticides, etc.

GENERAL INFORMATION application to the soil for are slow to appear and moisture. The degree of applied, soil type, rainfall 10% Bromacil Pellets is

AG 697 0585

JOHN DOE CH

CONTINUED FROM LEFT PANEL

WEED CONTROL: To control most weeds for an extended period of time on noncropland areas such as equipment storage sites, industrial sites, ditch banks, around farm buildings, or other noncrop areas that are not immediately adjacent to cropland or desirable trees and shrubs: apply preemergence or early postemergence to weed growth by spreading 10% Bromacil Pellets uniformly over the surface of the ground to be treated. Apply at the following rates; use the higher levels of the dosage ranges on adsorptive soils (those high in organic matter or carbon).

Annual Weeds and Grasses such as foxtail, ryegrass, wild oats, crabgrass, cheatgrass, brome grass, ragweed, lambsquarters, puncturevine, and turkey mullein: use 30 to 60 pounds per acre.

Perennial Weeds and Grasses such as smooth brome, bahiagrass, bluegrass, redtop, purpletop, quackgrass, broomsedge, aster, dandelion, dogfennel, goldenrod, plantain, and wild carrot: use 60 to 120 pounds per acre. Higher rates of 120 to 240 pounds per acre are required for johnsongrass and other hard to-kill perennials such as bermudagrass, vaseygrass, dallisgrass, nutsedge, saltgrass, bouncingbet, dogbane, bracken fern and horsetail; apply these high rates only as spot treatments to localized infestations of weeds.

Retreatment: Apply 30 to 60 pounds per acre when annual weeds and grasses reappear on sites where weed growth has been controlled.

For Small Areas — One-half pound per 435 square feet is approximately 50 pounds per acre.

NOTICE TO BUYER: Purchase of this material does not confer any rights under patents of countries outside of the United States.

STORAGE AND DISPOSAL

Storage: Store product in original container only, away from other pesticides, fertilizer, food or feed.

Pesticide Disposal: Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles and empty residue into application equipment. Then dispose of liner and drum in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

PELLETS

10%

90%

EPA Reg No 352 332

US

NO SKIN.

roughly with soap and

ation persists.

non if irritation persists.

ring of equipment or

result from failure to
desirable trees or other
the chemical may be
lks, driveways, tennis
keep from contact with
ation equipment after

elleted product for dry
ennial weeds. Effects
has been activated by
amount of chemical

nonflammable.

Made in U.S.A. Printed in U.S.A.

MADE IN THE U.S.A.