



352-370

6-6-1979

# AGRICULTURAL BULLETIN



LANNATE® L INSECTICIDE  
COTTON  
EAST OF ROCKY MOUNTAINS

4/1

## RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

SUPPLEMENTAL LABELING  
EPA REG. NO. 352-370

LANNATE® L INSECTICIDE  
FOR USE ON COTTON  
EAST OF THE ROCKY MOUNTAINS

Du Pont Lannate® L Insecticide is recommended for control of cotton bollworm and tobacco budworm on cotton east of the Rocky Mountains. When Lannate® L is applied to eggs, control is from direct kill of eggs (ovicide) and newly hatched larvae (larvicide).

### HOW TO USE

Apply 1/2 to 1 pt. Lannate® L per acre in sufficient water (minimum 5 gals. per acre by air) to obtain thorough coverage. Initiate schedule of sprays when eggs and newly hatched larvae are present. Continue at 5- to 7-day intervals while eggs are present and larval control is adequate. If significant numbers of larvae survive the treatment, use higher rates of Lannate® L (as registered) or use other registered larvicides.


Periodic scouting of fields is required to insure proper timing of spray.

Note: Lannate® L may be applied up to 15 days before harvest. Do not graze or feed treated cotton to livestock. Use of Lannate® L may redden cotton foliage. If reddening is excessive, alternate with other insecticides or discontinue use.

### IMPORTANT

BEFORE USING LANNATE® L, READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on package label. Follow the instructions carefully.



5/29/79

