

ACCEPTED  
 8-7-67  
 UNITED STATES DEPARTMENT OF AGRICULTURE  
 FEDERAL AGRICULTURAL MECHANICAL SERVICE  
 OFFICE OF CHEMICAL REGISTRATION  
 WASHINGTON, D. C. 20250

grass usually requires two or more applications and if crab grass "yellows" and revives, it will be evident that additional treatments are needed and these should begin, before seed head formation, and continued at 5-7 day intervals.

**CAUTION:** Avoid contact with the skin or prolonged breathing of spray mist. Wash hands after using. Avoid contamination of feed and foodstuffs and keep beyond the reach of children and domestic animals. Keep livestock off treated areas.

**ANTIDOTE:** If taken internally induce vomiting and call a physician at once.

**GUARANTEE:** effective for purpose claimed.  
 Disodium Monomethylarsonate Hexahydrate.....50.0%

**ACTIVE INGREDIENT:**  
 Disodium Monomethylarsonate Hexahydrate.....50.0%

**INERT INGREDIENTS:**.....50.0%

Total.....100.0%

Arsenic, as elemental, all in water soluble form.....12.83%


Canadian Reg. No. P.C.P. Act.

LICENSED UNDER U.S. PAT. NO. 2,678,265

USDA Reg. No. 358-33

# SAVE YOUR LAWN

**NOTT'S**



**POWDER**  
**50%**  
**D. S. M. A.**

# Kill Crab Grass

**CAUTION:** Keep Out of Reach of Children  
 See other cautions on side panel.

**NOTT MANUFACTURING CO., INC.** Net Weight  
 Pleasant Valley, New York Made in U.S.A. **28 OUNCES**

## DIRECTIONS

Do not use when grasses are stunted or dormant, due to drought. On newly seeded lawns do not apply until after the 3rd mowing. Water lawn 1 day previous to applying ARTOX and water lightly again for 3 days, beginning 2 days after treatment. If soil is moist from rain no watering is required.

**WHEN TO APPLY:** Early control should be practiced, beginning in June when young crab grass seedlings are less than 2" tall. Repeat treatments, 5-7 days apart, will be necessary to control new germination. Use PRESSURE or HOSE END SPRAYER. Wet thoroughly the foliage of all crab grass plants. Rinse spraying equipment with plain water after use.

If discoloration occurs with first application, on lawns predominately Bent, Fescue or White Clover, reduce application rate by 50%. Do not use on St. Augustine grass.

**WHEN AND HOW TO USE:** At below 80°, or for early control of young crab grass plants, with PRESSURE or HOSE END SPRAYER use 1 oz. (2 tablespoonfuls) to 1 gal. water to cover 250 sq. ft. of lawn area, or use 4 ozs. (8 tablespoonfuls) to 4 gals. water to cover 1000 sq. ft. When temperature is 80° and above, and for the control of "mature" crab grass, use only 1/2 oz. (1 tablespoonful) to 1 gal. water to cover 250 sq. ft., or 4 oz. to 8 gals. water to cover 2000 sq. ft. Mature crab

ACCEPTED WITH COMMENTS