

McDaniel, Pacific, Willamette, Peach Silver, Plum, Nursery and Yellow Mites affecting PEACHES, APRICOTS, NECTARINES, CHERRIES, PLUMS, PRUNES, FIGS, apply at petal fall to first cover spray or whenever mites appear. If repeat application is necessary, wait 30 days to avoid excessive residues. Do not apply to Cherries, Plums, Prunes or Figs within 7 days of harvest, nor on other fruits listed above within 14 days of harvest.

**GRAPES, RASPBERRIES, MITS:** For control of two-spotted (Red Spider), European Red, Pacific, McDaniel, Willamette and Yellow Mites affecting Grapes, Raspberries, Dewberries, Loganberries, Boysenberries, Walnuts, Filberts, Pecans, Chestnuts, Hickory Nuts, apply just before bloom or whenever mites appear. Do not feed treated bean vines to meat or dairy animals. On Nuts do not apply within 14 days of harvest. Do not use on Grapes within 7 days of harvest, nor on Berries within 2 days of harvest.

**CLOVER MITES IN HOMES:** Spray walls, windowsills, foundations, also plants and ground at base of infested building. Outdoor application only.

**CAUTION:** Avoid skin contact, in case of skin contact wash with soap and water. Avoid inhalation of spray mist, do not take internally. Avoid contamination of feed and food-stuffs. Wash thoroughly after handling. Rinse empty container thoroughly with water and discard it.


**Caution: COMBUSTIBLE MIXTURE** — Do not use or store near heat or open flame.

**ACTIVE INGREDIENT:**

1,1,1-trichloro-2,2,2-trifluoroethane 18.5%

**INERT INGREDIENTS:** 81.5%

Total 100.0%



PLEASANT VALLEY, N. Y. 12569

EPA REG. NO. 358-157

\* T.M. Rohm & Haas Co.



# KELTHANE

**KILLS MITES ON PLANTS KILLS CLOVER MITES WHICH ENTER HOMES**

**CAUTION**

Keep Out Of Reach Of Children Contents  
See Directions and Cautions on Side Panels. 1 pint

Contents:

Contents  
1 Quart

1 Pint

8 fl. ozs.

4 OZS.

2 1/2  
1 1/2  
1 1/2

6 1/2  
7 5/8  
5

4 1/2  
8 1/2  
10 1/2

2 3/4  
2 1/2  
3 1/2

**TIME TO APPLY:** Applications should be made when the various mite species appear in large numbers. Repeat application as necessary.

**HOW TO APPLY:** Wet thoroughly all parts of the plant, especially the undersides of leaves. **AMOUNT TO USE:** 2 teaspoonfuls to 1 gal. water, or 6 oz. to 25 gals. water.

For control of Red Spider, Spruce and European Red Mites affecting ROSES, EVERGREENS, SHADE TREES, FLOWERS. Apply when mites are present and repeat as necessary.

**VEGETABLES:** For control of two-spotted Mites (Red Spider), Pacific, Tropical and Tomato Russet Mites on Beans, Cucumbers, Cantaloupes, Melons, Watermelons, Pumpkins, Winter and Summer Squash, Tomatoes, Peppers, Eggplant. Do not use on beans within 7 days of harvest, nor on other crops in this section within 2 days of harvest.

Do not feed treated bean vines to meat or dairy animals.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on the label.

**STRAWBERRIES:** For control of two-spotted Mites (Red Spider), Strawberry Mite (Atlantic Mite) and six-spotted Mite. For Cyclamen Mite use 2 to 4 teaspoons to 1 gal. water and drench plants thoroughly at 10 to 20 day intervals in Spring when new growth starts, or use as spot treatment during season. Do not apply within 2 days of harvest.

**FRUITS:** For control of European Red Mite, Clover Mite, two-spotted (Red Spider) Schuene, McDaniel, Willamette, Pacific yellow and Apple Rust Mites, affecting APPLES, PEARS, CRABAPPLES, QUINCES, apply in second or later cover sprays or whenever mites appear. Do not apply within 7 days of harvest.

For European Red Mite, two-spotted (Red Spider), Schuene.

305 157