

No Bunt "40" is recommended only for the control of Bunt on seed wheat.

DIRECTIONS FOR USE

DRY APPLICATION: 1 ounce per bushel (or 1/2 ounce per bushel, where this is a State recommendation). If equipment is not available, dry application may be made in a revolving barrel type of equipment or on a clean floor by mixing the seed and seed treatment together with a shovel. Mix thoroughly to give uniform coating on each seed.

SLURRY APPLICATION: Mix with water as below:

Amount No Bunt "40"	Cup Size for:	Cup Size for:	Amount No Bunt "40"
for 1 gal. slurry	1/2 oz. per bu.	1 oz. per bu.	for 10 gal. slurry
14 oz.	23 cc	46 cc.	8 3/4 lbs.

If machine is not equipped with 46 cc. cup to make application of 1 oz. per bushel, make the following slurry and use the 23 cc. cup:

28-oz.	23 cc.	17 1/2 lbs.
--------	--------	-------------

Apply in slurry treater using 23 cc. or 46 cc. metering cup, bucket or rotor (depending on machine used) and correct seed gate. Treatment can also be made by adding 1/3 pint of the low strength slurry to each bushel of seed to obtain 1/2 ounce rate, or by adding 2/3 pint of the same slurry to each bushel of seed for 1 ounce treatment, using a revolving barrel or similar equipment. Mix thoroughly to obtain uniform coverage of seed.

Local conditions may affect the use of this chemical. Consult State agricultural extension or experiment station specialists for specific recommendations and for information on possible lower dosages.

Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

PRODUCT NO.

508

10-8-71
359-293

NET WEIGHT
5 POUNDS

NO BUNT "40"

(SEED DISINFECTANT)

ACTIVE INGREDIENT:

Hexachlorobenzene 40.0%

INERT INGREDIENTS 60.0%

CAUTION: Keep Out of Reach of Children

Avoid contact with skin and eyes.

Reasonable care should be exercised in the use of this product. Avoid inhalation. Wash thoroughly after handling. Avoid contamination of feed and foodstuffs. Treated seed should not be used for feed, food or oil purposes.

Label treated seeds as follows:

"This seed treated with hexachlorobenzene, NO BUNT "40". Do not use for food feed or oil purposes."

Can be applied in standard slurry and dust treaters.

Do NOT reuse container. Destroy when empty.

Do NOT Store with Foodstuffs

Manufactured by

RHODIA INC.
CHIPMAN DIVISION

NEW BRUNSWICK, NEW JERSEY

Atlanta, Ga.	Bound Brook, N. J.	Fresno, Calif.	Hinsdale, Ill.
Pasadena, Tex.	Portland, Ore.	St. Paul, Minn.	

EPA Reg. No. 359-293 AB

Page 6C 3 371