

PRODUCT NO.

290

NET WEIGHT
50 POUNDS

CHIPMAN[®] META A PELLET

AN AGRICULTURAL POISON BAIT

For control of slugs, snails and cutworms in strawberry plantings [redacted] and in flower gardens.

CAUTION: Keep Out of Reach of Children

ACTIVE INGREDIENTS:

Tricalcium arsenate 5.16%
Metaldehyde 3.00%

INERT INGREDIENTS 91.84%

Arsenic expressed as metallic 1.94%
Arsenic in water soluble form expressed as metallic. Not more than 1.00%

WARNING: May Be Fatal If Swallowed! Avoid contact with skin, eyes or clothing. Wash thoroughly after using. Avoid breathing dust. Avoid contamination of feed and foodstuffs. Keep away from children and domestic animals. Field workers should use protective gloves when handling and broadcasting Chipman Meta A Pellets. Burn this bag when empty—Bury the ashes where they will not be a hazard to humans or animals. *add here in separate paragraph*

RECOMMENDATIONS

CHIPMAN META A PELLET is a special formulation which employs apple chop as carrier for the main toxicants, calcium arsenate and metaldehyde.

WHERE TO USE:

[redacted] In Strawberry plantings apply any time, but not within 14 days of hand picking. Apply so as not to contaminate fruit. In non-crop areas to protect flowers and other non-food plants.

This product is toxic to wildlife.

Created and exposed on soil surface Do NOT Store with Foodstuffs

will be hazardous to birds and other wildlife.

HOW TO APPLY: Apply uniformly by mechanical or power spreader or by hand. For hand application, broadcast liberally under trees, around irrigation standpipes, buildings, and other places where slugs and snails concentrate. Repeat application following irrigation.

WHEN TO APPLY: Slug control should be viewed as a year-round practice [redacted] Baits applied soon after the first rains will kill many slugs before they can lay eggs. Best results are obtained when material is applied in the evening. Avoid treatments during periods of heavy rain-fall. Repeat as needed.

AMOUNT TO USE: Apply at the rate of 10 to 20 pounds per acre. Repeat as needed.

CAUTION: Do not apply to foliage or other plant parts. Do not contaminate edible parts of plants. Do not apply against the base of tender plants or they may be damaged. Do not apply to strawberries within 14 days of picking.

Local conditions may affect the use of this chemical. [redacted] for specific recommendations as to dosage and time of application and for control of other pests.

Follow directions carefully. Timing and method of application, weather and crop condition, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

ACCEPTED
5-10-68

359-536

U.S.D.A. Reg.
No. 359-536

B B. 2M-1-28

Atlanta, Ga
Bound Brook, N. J.
Chicago, Ill
N. Kansas City, Mo

Manufactured by
RHODIA INC.
CHIPMAN DIVISION
NEW BRUNSWICK, NEW JERSEY

Palo Alto, Calif.
Pasadena, Tex.
Portland, Ore.
St. Paul, Minn.

