

ORNAMENTAL TURF USE ONLY

GENERAL INFORMATION

CHIPCO SPOT KLEEN is a wettable powder for use on golf course greens, tees, and fairways of Bentgrasses, Bluegrasses, Bermudagrasses, Fescues, Ryegrasses, St. Augustine grasses or their mixtures. CHIPCO SPOT KLEEN is not phytotoxic to any of the above mentioned grasses when used in accordance with the label. CHIPCO SPOT KLEEN is to be used for the prevention and control of the below mentioned diseases (provides both curative and protective action).

DIRECTIONS FOR USE

Apply with hand sprayer, power sprayer, or a hose-on proportioner. Add the recommended amount of CHIPCO SPOT KLEEN to a partially filled spray tank under agitation. Complete filling the tank; continuous agitation is necessary to insure even suspension of the wettable powder. Use spray mixture the same day it is prepared. Spray uniformly over the area to be treated. Apply after mowing or avoid mowing for twelve hours after application.

Apply recommended amounts in sufficient water to obtain thorough coverage, usually five gallons per 1,000 sq. ft. of turf area. When treating golf greens, always treat aprons and approaches. Use the highest recommended rate under conditions of severe disease pressure.

DOLLAR SPOT (*Sclerotinia homoeocarpa*), COPPER SPOT (*Gloeocercospora sorghi*) — Apply CHIPCO SPOT KLEEN at ¾ to 1½ oz. per 1,000 sq. ft., beginning when the disease first appears and continue at 10-14 day intervals through the season, or as needed.

LARGE BROWN PATCH (*Rhizoctonia solani*) — Apply CHIPCO SPOT KLEEN at ¾ to 1½ oz. per 1,000 sq. ft., beginning when the disease first appears and continue at 10-14 day intervals through the season or as needed. When conditions are unusually favorable for development of disease, reduce interval to 5-7 days.

FUSARIUM BLIGHT (*Fusarium roseum*) — Apply CHIPCO SPOT KLEEN at 4-8 oz. per 1,000 sq. ft. Water into the root zone by irrigating with one inch of water immediately after application. Apply two applications at 10-14 day intervals, beginning applications when the disease first appears.

STRIPE SMUT (*Ustilago striiformis*) — Apply CHIPCO SPOT KLEEN at 4-8 oz. per 1,000 sq. ft. Water into the root zone by irrigating with one inch of water immediately after application. Apply two applications at 14-21 day intervals. Begin applications when disease first appears, either in the spring or the fall.



ACCEPTED
August 21, 1973
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER-
ED UNDER NO. 359-665

“THE RIGHT APPROACH TO GOOD TURF”

CHIPCO SPOT KLEEN*

CHIPCO SPOT KLEEN*

TURF FUNGICIDE

ACTIVE INGREDIENT:		
Dimethyl 4,4' -o-phenylenebis [3-thioallophanate]	70%	
INERT INGREDIENTS	30%	
TOTAL	100%	


*TOPSIN M—Nippon Soda Company's trademark for Dimethyl 4,4' -o-phenylenebis [3-thioallophanate] licensed to Pennwalt Corporation.

U. S. Patent Pending

CAUTION: KEEP OUT OF REACH OF CHILDREN

See Specific Caution Statements on Side Panel

NET WEIGHT TWO POUNDS



May be hazardous
Avoid breathing
after handling
feed or food

Keep children
soil and air

Do not graze
livestock

External:
flush with

Internal:
CALL A PHYSICIAN

Burn or bury

EPA Reg. No.

and fairways of
grasses or their
grasses when
vention and con-
n).

Recommended amount
filling the tank;
powder. Use spray
apply after mow-

ally five gallons
approaches. Use

brghi) — Apply
se first appears

to 1 1/2 oz. per
intervals through
ent of disease.

per 1,000 sq. ft.
plication. Apply
st appears.

er 1,000 sq. ft.
plication. Apply
ears, either in



ACTIVE INGREDIENT:

Dimethyl 4,4' -o-phenylenebis [3-thioallophanate] 70%

INERT INGREDIENTS 30%

TOTAL 100%


*TOPSIN M—Nippon Soda Company's trademark for Dimethyl 4,4' -o-phenylenebis [3-thioallophanate] licensed to Pennwalt Corporation.

U. S. Patent Pending

CAUTION: KEEP OUT OF REACH OF CHILDREN

See Specific Caution Statements on Side Panel

NET WEIGHT TWO POUNDS



CAUTION

May be harmful if swallowed or inhaled. Avoid contact with eyes, skin, or clothing. Avoid breathing of dust or spray mist. Wash thoroughly with soap and water after handling. Wash contaminated clothes before reuse. Do not contaminate feed or food.

Keep children and pets off treated area until material has been washed into the soil and area is completely dry.

Do not graze animals on treated turf. Do not feed clippings from treated turf to livestock and poultry.

ANTIDOTE

External: If spilled on skin, wash off immediately with soap and water. If in eyes, flush with plenty of water.

Internal: Induce vomiting with a mixture of salt, soap, or mustard in warm water. **CALL A PHYSICIAN IMMEDIATELY.**

Burn or bury empty containers, stay out of smoke; never reuse them.

EPA Reg. No. 359-665-AA

PL-0573

PRODUCT NO. 331G

Manufactured for

CHIPMAN DIVISION of RHODIA INC.
NEW BRUNSWICK, NEW JERSEY

