



*CAPTAN 14.00%
 *N-trichloromethylthio-4-cyclohexene-1, 2-dicarboximide

INERT INGREDIENTS 61.40%

TOTAL 100.00%

CAUTION: Harmful if swallowed. Avoid breathing of spray mist. Avoid contact with skin; wash hands thoroughly after using. Avoid contamination of feed and foodstuffs. Keep out of reach of children.

USDA Reg. No. 370-16

GARDEN HOSE SPRAY CO., INC.
 CAMBRIDGE, MASSACHUSETTS

Printed in U.S.A.

Net Weight — 1½ Ozs.



MULTI-PURPOSESPRAY (Insecticide & Fungicide)

For "Arnold" Garden Hose Sprayer

CAUTION: Keep out of reach of children.



FOR COMPLETE SPRAYING INSTRUCTIONS SEE ENCLOSED WRAPPER
 ACTIVE INGREDIENTS:

*MALATHION 6.60%

* (0—0-dimethyl dithiophosphate of diethyl mercaptosuccinate)

*METHOXYCHLOR Technical 15.00%

*Equivalent to 13.2% 2, 2-bis (p-methoxyphenyl)-1,1,1-trichloroethane and 1.8% other isomers and reaction products.

"ARNOLD"

MULTI-PURPOSE SPRAY INSECTICIDE AND FUNGICIDE CARTRIDGE

FOR: FRUITS — VEGETABLES — ORNAMENTALS

Spraying Instructions

Use first application when new, small leaves first appear, and repeat every 7 to 10 days for at least eight applications. Consult your local extension service or Agricultural Experiment Station to determine the use of this spray under local weather or soil conditions.

FRUITS: Apples, Peaches, Pears, Cherries, Grapes, Raspberries, Strawberries, Black Spot, Sooty Blotch, Bitter Rot, Brown Rot, Leaf Spot, Rust, Spot and Fruit Rot.

Insects: Codling Moth, Plum Curculio, Apple Maggot, Tent Caterpillar, certain Fruit Moths, Pear Psylla, Plant Bug, Leaf Hoppers and Aphids.

VEGETABLES: Beans, Cucumbers, Eggplant, Melons, Peppers, Potatoes, Tomato, Squash and Tomatoes.

Diseases: Early and late blights of Potatoes and Tomatoes, Anthracnose of Tomatoes, Downy Mildew and Angular Leaf Spot of Cucumber and Melons.

Insects: Flea Beetles, Cabbage Worm, Cucumber Beetles, Fall Army Worm, Cutworms, Aphids, Flea Beetles, Leaf Hoppers, Mexican Bean Beetles, Mites, Colorado Potato Beetles and Squash Vine Borer.

ORNAMENTALS: Black Spot of Roses and Leaf Spot on Chrysanthemums.

Diseases: Black Spot of Roses and Leaf Spot on Chrysanthemums.

Insects: Aphids, certain Caterpillars, Flea Beetles, Mites, Rose Chafers and Tarnished Plant Bugs.

ACCEPTED

NOV 7 1969
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR ECONOMIC POISON REGISTERED
 UNDER NO 3116 SUBJECT
 TO ATTACHED COMMENTS.

"ARNOLD"

MULTI-PURPOSE SPRAY INSECTICIDE AND FUNGICIDE CARTRIDGE
FOR: FRUITS — VEGETABLES -- ORNAMENTALS

Spraying Instructions

For best application when new, small leaves first appear, and repeat every 7 days for at least eight applications. Consult your local extension agent or Agricultural Experiment Station to determine the use of this spray under local heat weather or soil conditions.

FRUITS: Apples, Peaches, Pears, Cherries, Grapes, Raspberries, Strawberries.

Diseases: Scab, Brooks Spot, Sooty Blotch, Bitter Rot, Brown Rot, Leaf Spot, Fruit Spot and ~~Fruit~~ Rot.

Insects: Codling Moth, Plum Curculio, Apple Maggot, Tent Caterpillar, Oriental Fruit Moth, Pear Psylla, Plant Bug, Leaf Hoppers and Aphids.

VEGETABLES: Beans, Cucumbers, Eggplant, Melons, Peppers, Potatoes, Pumpkin, Squash and Tomatoes.

Diseases: Early and late blights of Potatoes and Tomatoes, Anthracnose of Tomatoes, Downy Mildew and Angular Leaf Spot of Cucumbers, Squash and Melons.

Insects: Aphids, Cabbage Worms, Cucumber Beetles, Fall Army Worms, Flea Beetles, Leaf Hoppers, Mexican Bean Beetles, Mites, Colorado Potato Beetles and Squash Vine Borer.

ORNAMENTALS:

Diseases: Black Spot of Roses and Leaf Spot on Chrysanthemums.

Insects: Aphids, certain Caterpillars, Flea Beetles, Mites, Rose Chafer and Tarnished Plant Bugs.

CAUTION: Do not use on Raspberries after fruit begins to form. Do not use on Strawberries within 14 days of harvesting. Do not use on peaches within 21 days of harvest. Do not use on Pumpkins or Grapes within 14 days of harvest. Do not use on any other crops within 7 days of harvest.

WARNING: KEEP OUT OF REACH OF CHILDREN. Harmful if swallowed. Avoid contamination of feed and foodstuffs. Avoid inhalation of spray mist. Avoid contact with skin; wash hands thoroughly after using.

NON-WARRANTY: The seller guarantees the material sold to be true and sound. Seller's guarantee shall be limited to the terms of the label and in no event the buyer assumes any risk to persons or property arising from the use, handling and accepts the product on these conditions.

GARDEN HOSE SPRAY COMPANY, INC.

Cambridge • Massachusetts

USDA Reg. No. 370-16

Made in U. S. A.

68

For best results and for a more concentrated spraying mixture this cartridge must be replaced by a new cartridge when one-third of it has been used.

When cartridges are replaced the old cartridge may be saved and when several pieces have accumulated they may be used by putting enough pieces in the chamber to completely fill it.

If, instead of replacing this cartridge when one-third used, an additional piece of cartridge be inserted so that the cartridge chamber is practically full again, the same concentration will be obtained.

Satisfactory spraying results have been attained by the use of this cartridge without changing or keeping the chamber practically full at all times but remember, maximum efficiency is reached only when the cartridge chamber is not less than two-thirds full.

GARDEN HOSE SPRAY CO., INC. CAMBRIDGE 40, MASS.

PRINTED IN U.S.A.

68

ACCEPTED

NOV 7 1969

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
ED UNDER NO. 370-16 SUBJECT
TO ATTACHED COMMENTS.